

**ANTHELMINTIC POTENCY OF NEEM (*Azadirachta indica*) LEAF MEAL ON WEST AFRICAN DWARF (WAD) SHEEP**

**ABSTRACT**

A 90-day study was conducted to determine the response of semi intensively managed West African dwarf sheep to concentrate supplement containing varying levels of neem leaf meal (NLM). Twenty (20) West African Dwarf sheep aged 5 to 6 months with an average weight of 10kg were used in a Complete Randomized Design with animals grouped into four treatments of five replicates each balanced for weight. The animals were allowed to graze on natural pastures predominantly made up of *Panicum maximum* in the morning with a daily supplementation of 100g concentrate diet containing varying levels of neem leaf meal at 0, 5, 10 and 15%. Blood samples were taken from the animals before the commencement of the experiment and at the end of the experiment. At the start of the experiment, faecal samples were collected from each animal to determine the faecal egg count and this was repeated once in three weeks for the 90 day experimental period. There was significant ( $P < 0.05$ ) difference in the haematology indices studied with no definite pattern. The inclusion of NLM in the diets of West African Dwarf sheep significantly ( $P < 0.05$ ) reduced the faecal egg counts across the treatments with a percentage reduction range of 33.38 to 88.00% for sheep on 0% and 5% NLM, respectively. This study, however, concluded that neem leaf inclusion at 5% in West African dwarf sheep's diet had effects on the overall performance of the animals with a potential improvement in drastic reduction in faecal egg counts.

**key word:** Haematology, faecal egg counts, *A. indica*, West African Dwarf Sheep.

## 31 INTRODUCTION

32 Throughout the world, internal parasites pose one of the major health limitations for grazing animals.  
33 Although there are numerous internal parasites, only a few of them account for the majority of problems  
34 for grazing livestock.

35 Helminth infections in small ruminants are serious problems of the developing world, particularly where  
36 nutrition and sanitation are poor (Faye *et al.*, 2003). Helminthosis is a primary factor in the reduction of  
37 productivity of these animals through mortality and reduced weight gains (Gatongi *et al.*, 1997). While  
38 some studies have reported that goats are more susceptible than sheep to a similar challenge, others  
39 have reported that sheep usually suffer heavier worm burdens because of the difference in their grazing  
40 habits (Tinar *et al.*, 2005).

41 Economic losses are caused by gastrointestinal parasites in a variety of ways: they cause losses through  
42 lowered fertility, reduced work capacity, involuntary culling, a reduction in feed intake and lower weight  
43 gains, lower milk production, treatment costs, and mortality in heavily parasitized animals.

44 Prevention rather than cure is the philosophy used in developing control programs against gastrointestinal  
45 nematodes. It should be assumed that worms cannot be eradicated from the environment and livestock  
46 will continually be reinfected. However, infections can be limited to the extent that they will not cause  
47 economic loss to the producer. A combination of treatment and management is usually necessary to  
48 achieve control (David, 2010).

49 Sheep and goat farmers rely heavily on anti-parasitic drugs, or anthelmintics to control internal parasites  
50 in their small ruminant flocks. A wide variety of anthelmintics, covering the entire range of chemical  
51 groups, are used for the treatment of nematode parasites of sheep and goat. However, due to the serious  
52 problem of anthelmintic resistance (Chandrawathani *et al.*, 2004), there is growing demand for alternative  
53 methods of parasite control to reduce the dependence on these drugs.

54 In 1999, a survey of 39 sheep farms and 9 goat farms found that the majority had worm populations  
55 resistant to all classes of drugs (Chandrawathani *et al.*, 1999). From this investigation, it was clear that  
56 anthelmintic resistance was rapidly increasing.

57 Neem leaf (*Azadirachta indica*) is efficient as an antibiotic, anthelmintic and growth promoter when added  
58 to the feed of ruminant. Preliminary studies done by Chandrawathani *et al.*, (2000) showed that feeding  
59 Neem foliage is safe, eco-friendly, cheap and palatable to sheep. *Ad libitum* feeding of fresh Neem leaves  
60 produced 82% reduction in worm eggs of sheep and a further trial on a limited number of sheep showed  
61 that Neem produced a significant reduction in worm burdens (Chandrawathani *et al.*, 2002).

62 This study however investigate the effect of varying inclusion of Neem leaf meal in promoting growth and  
63 reducing helminth infections in West African Dwarf sheep grazing natural pasture.

64

## 65 MATERIALS AND METHOD

### 66 EXPERIMENTAL SITE

67 The experiment was carried out in the Sheep unit of Federal College of Forestry, Forestry Research  
68 Institute of Nigeria (FRIN), Jericho hill, Ibadan, Oyo State. It is located on the latitude 07<sup>0</sup>23'32"N and  
69 longitude 03<sup>0</sup>51'44"E with altitude 212m above sea level. The rainfall pattern is bimodal with peaks  
70 around June to July, and September to October. The mean annual rainfall is about 420mm in 109 days  
71 with mean maximum and minimum temperature of about 34<sup>0</sup>C and 24<sup>0</sup>C respectively. Mean relative  
72 humidity ranges from about 82% between June and September to approximately 60% between  
73 December and February (FRIN, 2014).

74

### 75 Experimental Animals

76 Twenty (20) growing West African Dwarf (WAD) sheep aged 5-6 months with average weight of 10kg  
77 were purchased from markets within Ibadan. The animals were quarantined for a period of 30 days. The  
78 experimental pens were disinfected with diazintol solution before the arrival of the animals. For the period  
79 of the experiment, the sheep were managed using semi intensive management system. They were  
80 allowed to graze on natural pastures which predominantly *Panicum maximum* in the morning from 8am  
81 and returned to their individual feeding pens after grazing for five hours.

82

### 83 Procurement and Processing of Experimental Material

84 Fresh Neem leaves samples were obtained from Neem trees in and around the Forestry Research  
85 Institute of Nigeria, Ibadan. The leaves were chopped for effective drying. The chopped leaves were sun  
86 dried for 3-4 days until they are crispy. The dry leaves were milled using a hammer mill to produce leaf  
87 meal before they were incorporated into the concentrate supplement at 0g, 5g, 10g, 15g Neem leaf/ 100g  
88 concentrate/animal/day respectively and fed to the animals before going out to graze for a period of 90  
89 days.

### 90 Animal Grouping and Treatment

91 The animals were grouped into four treatments of five replicates each balanced for weight namely;

92 Treatment 1: 0g of Neem leaf /100g concentrate/animal/day

93 Treatment 2: 5g of Neem leaf/100g concentrate/animal/day

94 Treatment 3: 10g of Neem leaf/100g concentrate/animal/day

95 Treatment 4: 15g of Neem leaf/100g concentrate/animal/day

96 The animals were supplemented daily with 100g concentrate experimental diet composed of maize,  
97 wheat offals, palm kernel cake, soyabean meal, bone meal with salt and premix, Neem leaf was added at  
98 varying levels (Table 1). Fresh, clean water was given to the animals *ad libitum*.

99

100 **Table 1: Composition of the Experimental Diets fed to sheep**

Ingredients (%)	Diets			
	0%NLM	5%NLM	10%NLM	15%NLM
Neem leaf meal	0	5	10	15
Maize	24	24	24	24
Palm kernel cake	20	15	10	5
Soyabean Meal	14	14	14	14
Wheat offals	38	38	38	38
Bone meal	2.5	2.5	2.5	2.5
Salt	1	1	1	1
Premix	0.5	0.5	0.5	0.5
Total	100	100	100	100

101

102 **DATA COLLECTION**

103 **Faecal collection**

104 Before the commencement of the experiment, faecal samples were collected from each animal to  
 105 determine the faecal egg count and this was repeated once in three weeks for the 90 day experimental  
 106 period. Hand gloves were used on the hands and the hand was dipped inside the rectum of the animals  
 107 to collect fresh faeces. Three grammes of each collected faecal samples were ground and mixed with  
 108 42ml of water. A saturated solution was poured into the mixture of faeces and water to float the eggs  
 109 following the modified McMaster method described by Miller *et al.*, (1998). A sample obtained from this  
 110 was collected and put into both compartments of McMaster counting chamber/slide and then viewed  
 111 under the microscope. The number of eggs within each viewed area was multiplied by 100 to get the  
 112 actual number of eggs per gram.

113

114 **Blood samples collection**

115 Blood samples were taken from the animals before the commencement of the experiment and at the end  
 116 of the experiment. Blood samples were collected via the jugular vein puncture using a 10ml hypodermic  
 117 syringe. Five milliliters of the blood was infused into collection bottles containing Ethylene Di-amine Tetra-  
 118 acetic acid (EDTA) for serum and the remaining 5ml into collection bottles without anti-coagulants for

119 plasma and taken to the laboratory for analysis. Blood parameters namely packed cell volume and  
120 haemoglobin concentration (HB) were determined following the procedure outlined by Schalm *et al.*,  
121 (1995). Red blood cell and total white blood cell were determined using haemocytometer (Dacie and  
122 Lewis, 1984). Serum biochemical parameters like serum urea nitrogen and serum total protein were  
123 determined by haemocytometer (Dacie and Lewis, 1984).

124

### 125 **Statistical Analysis**

126 All data collected were subjected to one way analysis of variance in a completely randomized design  
127 according to SAS (1999) and means were separated using the Duncan Multiple Range Test (Duncan,  
128 1955).

129

## 130 **RESULT AND DISCUSSION**

### 131 **Pre-haematology and serum indices of WAD sheep fed varying inclusion level of Neem** 132 **leaf meal**

133 Table 2 shows the pre-haematology values of the animals; urea nitrogen, packed cell volume,  
134 haemoglobin concentration, red blood cell, white blood cell and serum total protein.

135 The values for the urea nitrogen range between 32.15mg/dl to 47.50mg/dl while packed cell volume  
136 range between 27.25% to 32.75%. The haemoglobin concentration ranges between 8.75g/dl to 10.92  
137 g/dl.  $4.15 \times 10^6/\mu\text{l}$  to  $5.03 \times 10^6/\mu\text{l}$ ,  $3.53 \times 10^3/\mu\text{l}$  to  $5.42 \times 10^3/\mu\text{l}$  and 5.19g/dl to 5.51g/dl are the value range  
138 for red blood cell, white blood cell and serum total protein respectively.

139 The packed cell volume values obtained at the pre-haematology were within the physiological range of  
140 27.0 – 45.0 % given by Jain (1993), slightly higher than the range of 25–30% reported by Opara *et al.*,  
141 (2010). In contrast to this, Taiwo and Ogunsanmi (2003) reported higher values of 35.5% and 36.9% for  
142 clinically healthy West African dwarf sheep. The haemoglobin concentration ranges between 8.75 to  
143 10.92g/dl which falls within the range of 9–15 g/dl reported by (Kaneko, 1997; Patra *et al.*, 2003), but  
144 higher than the values of 5 to 6 g/dl obtained by Belewu and Ogunsola (2010) for goats. The red blood  
145 cell counts falls within the range of  $4.3 - 5.03 \times 10^6/\mu\text{l}$  the counts reported in this study fell below the range  
146 of  $10.25 - 12.85 \times 10^6/\mu\text{l}$  (Ajala *et al.*, 2000), 9.2–13.5 g/dl (Tambuwal *et al.*, 2002), 9.9–18.7 /dl by (Taiwo  
147 and Ogunsanmi, 2003). The white blood cell count falls between  $3.53 \times 10^3/\mu\text{l} - 5.42 \times 10^3/\mu\text{l}$ . The WBC  
148 counts were similar among the treatment groups and fell within the normal range (5 to 11g/dl) reported by  
149 Scott *et al.*, (2006) for sheep. The total serum protein of the animals falls between the range of 5.19-  
150 5.51mg/dl.

151

152

153

154

155 **TABLE 2: Pre-haematology and serum indices values of West African Dwarf sheep fed**  
 156 **concentrate containing varying inclusion levels of Neem leaf meal**

Parameters	0%NLM	5%NLM	10%NLM	15%NLM	±SEM
Packed Cell Volume (%)	29.75	28.25	27.25	32.75	1.61
Haemoglobin concentration (g/dl)	9.77	9.00	8.75	10.92	0.34
Red Blood Cell ( $\times 10^6/\mu\text{l}$ )	4.30	4.39	4.15	5.03	0.15
White Blood Cell ( $10^3/\mu\text{l}$ )	4.18	3.53	4.25	5.42	0.08
Urea Nitrogen (mg/dl)	34.81	39.28	47.5	32.15	1.83
Serum Total Protein(g/dl)	5.50	5.19	5.36	5.51	0.09

157 NLM- Neem leaf meal

158

159 **Post haematology and serum indices of WAD sheep fed concentrate supplement**  
 160 **containing varying inclusion level of Neem leaf meal**

161

162 Table 3 shows the post haematology values of WAD sheep fed varying inclusion levels of NLM; urea  
 163 nitrogen, packed cell volume, haemoglobin concentration, red blood cell, white blood cell and serum total  
 164 protein. Animals on the control (0% NLM) (25.62mg/dl) had the lowest urea nitrogen at the post  
 165 haematology while 15% NLM had the highest urea nitrogen (33.39mg/dl).

166 For the packed cell volume (PCV), the values are 26.25%, 31.25%, 24.25% and 27.25% for 0% NLM to  
 167 15% NLM respectively. 5% NLM (31.25%) had the significantly highest packed cell volume at the post  
 168 haematology. PCV was significantly higher at 5% inclusion level of NLM than other treatment groups.

169 For haemoglobin concentration, 10% NLM (10.42g/dl) had a significantly higher ( $P < 0.05$ ) value when  
 170 compared to other treatments. The values range between 7.59g/dl to 10.42g/dl from 0% NLM to 15%  
 171 NLM. For the red blood cell count, 10% NLM ( $8.80 \times 10^6/\mu\text{l}$ ) had the highest red blood cell count while 0%  
 172 NLM ( $7.59 \times 10^6/\mu\text{l}$ ) had the lowest red blood cell count, the values are  $7.59 \times 10^6/\mu\text{l}$ ,  $8.71 \times 10^6/\mu\text{l}$ ,  $8.80 \times$   
 173  $10^6/\mu\text{l}$  and  $8.43 \times 10^6/\mu\text{l}$  for 0% NLM to 15% NLM respectively. 15% NLM had the highest white blood cell  
 174 count ( $6.82 \times 10^3/\mu\text{l}$ ).

175 The post haematology values for all the parameters monitored differ among the dietary treatment. Urea  
 176 nitrogen at the post haematology decreased across the treatment compared to the pre haematology  
 177 except for treatment 4 which increased by 1.24. Although, Blood urea level was slightly higher for

178 treatments with NLM inclusion compared to the control (0% NLM), they were within the normal range.  
179 This could be due to the higher crude protein contents of NLM supplemented treatments, in which there  
180 was improvement in the crude protein content by the treatment materials confirming the observation by  
181 Coles (1986) that high dietary protein is associated with increase in urea level.

182 Sheep fed 5% NLM had a higher PCV at the post haematology compared to pre-haematology, it  
183 increased by 3.00 compared to other treatments that decreased at the post haematology. Although, there  
184 was reduction in the PCV of treatment1, 3 and 4 at post haematology, PCV of this work still falls within  
185 the range of 21-35% and 20.10-48.00% reported for West African Dwarf goats and Afec-Awassi sheep by  
186 Daramola *et al.*, (2005) and Jawasreh *et al.*, (2010) respectively. This indicated that the PCV has not  
187 been affected in all the treatments. It further showed that in all the treatments, animals did not suffer from  
188 anaemia or dehydration. This confirms the report of The Merck Veterinary Manual (1998) that a low PCV  
189 value was an indication of anaemia while sharp increase in PCV is most often caused by dehydration.

190 Sheep fed 5%NLM (10.42) had the highest Haemoglobin concentration. Animal fed 5% NLM had a higher  
191 value at the post haematology compared to the pre haematology, it increased by 1.42 while other  
192 treatments decreased as compared to the post haematology. The values reported in this study were  
193 within the range of 7-15 and 8.15-10.75 gL<sup>-1</sup> reported for West African Dwarf goats and West African  
194 Dwarf sheep by Daramola *et al.*, (2005) and Akinyemi *et al.*, (2010), respectively. Ogbuewu *et al.*, (2010)  
195 reported the highest haemoglobin concentration at 5% inclusion level of neem leaf meal in the diet of  
196 rabbits. The implication of the values obtained in this study is that the dietary proteins were of high quality  
197 (Abu *et al.*,1998).

198 The haemoglobin concentration (Hb) in the blood of the studied animals showed a similar pattern of  
199 variation as with PCV. Mean Hb concentration was higher in animals fed 5% NLM than in other  
200 treatments. With the relatively higher Hb concentration observed in 5% NLM, the dietary treatment  
201 seemed to be capable of supporting high oxygen carrying capacity blood in the sheep.

202 The post haematology values of the red blood cells increased across all the treatments compared to the  
203 pre haematology. The RBC counts reported in this study fell below the range of 10.25–12.85 × 10<sup>6</sup>/μl  
204 obtained by Ajala *et al.*, (2000), 9.2 – 13.5 ×10<sup>6</sup>/l reported by Tambuwal *et al.* (2002) and 9.9 – 18.7  
205 ×10<sup>6</sup>/μl by Taiwo and Ogunsanmi (2003).

206 The non significant value of red blood cells (RBC), packed cell volume (PCV) and hemoglobin (Hb) of the  
207 sheep on NLM diets relative to the control group is an indication that the animals were not anemic. The  
208 PCV and Hb values of sheep in the test diets were not different from the control group. This tends to  
209 confirm the report of Talebi *et al.* (2005) that nutrition affect the blood profiles of animal and this implies  
210 that up to 15% inclusion of NLM had a positive effect on the relative quantity of blood cell as well as total  
211 volume of blood.

212 Meanwhile, the white blood cell values at the post haematology are above the range of 2.23-3.48×10<sup>3</sup>/μl  
213 reported by Ukanwoko *et al.*, (2013). White blood cell in animal possesses phagocytic function (Campbell  
214 and Coles, 1986) and differential white blood cell counts were used as an indicator of stress response

215 and sensitive biomarkers crucial to immune function (Graczyk *et al.*, 2003). The white blood cell values at  
 216 5% NLM was the least in this study disagrees with the findings of Ososanya *et al.*, (2014) that recorded a  
 217 highest white blood cell value for WAD ewes fed water-washed neem fruit supplemented diet at 5%.

218 Sheep fed 5% NLM had an increase in the serum total protein in post haematology compared to the pre-  
 219 haematology while the 0% NLM, 10% NLM and 15% NLM had a decrease in the post haematology  
 220 compared to the pre haematology. Animals fed 5% NLM (5.32g/dl) had the highest serum total protein  
 221 while 0% NLM (4.61g/dl) had the lowest serum total protein. The values were within the range of 5.0-  
 222 12.3(g/dl) but lower than 6.3-8.5 (g/dl) reported for Afec-Awassi sheep and West African Dwarf goats by  
 223 Jawasreh *et al.*, (2010) and Daramola *et al.*, (2005), respectively. The implication of this result is that the  
 224 highest increase in total protein in the serum of the experimental animals in 5% NLM would suggest that  
 225 protein synthesis was efficient. The serum protein concentration indicates the balance between  
 226 anabolism and catabolism of protein in the body. The serum protein concentration at any given time in  
 227 turn is a function of hormonal balance, nutritional status, water balance and other factors affecting health  
 228 (Abdel Hameed *et al.*, 2011).

229  
 230 **TABLE 3: Post-Haematology and serum indices values of WAD sheep fed concentrate**  
 231 **supplement containing varying inclusion levels of Neem leaf meal**

Parameters	0%NLM	5%NLM	10%NLM	15%NLM	SEM
Packed Cell Volume (%)	26.25 <sup>b</sup>	31.25 <sup>a</sup>	24.25 <sup>b</sup>	27.25 <sup>ab</sup>	±0.97
Haemoglobin Concentration (g/dl)	8.78 <sup>b</sup>	10.42 <sup>a</sup>	8.10 <sup>b</sup>	9.10 <sup>ab</sup>	±0.32
Red Blood Cell (×10 <sup>6</sup> /µl)	7.59 <sup>c</sup>	8.71 <sup>ab</sup>	8.80 <sup>a</sup>	8.43 <sup>b</sup>	±0.24
White Blood Cell (×10 <sup>3</sup> /µl)	5.28 <sup>b</sup>	4.53 <sup>c</sup>	6.00 <sup>ab</sup>	6.82 <sup>a</sup>	±2.37
Urea Nitrogen (mg/dl)	25.62 <sup>b</sup>	32.31 <sup>a</sup>	31.94 <sup>b</sup>	33.39 <sup>ab</sup>	±2.42
Serum Total Protein (g/dl)	4.61 <sup>b</sup>	5.32 <sup>a</sup>	4.86 <sup>b</sup>	5.26 <sup>a</sup>	±0.19

232 <sup>a,b,c</sup> Mean values followed by the same letter in the same row are not significantly different (P≤ 0.05)

233

234



## 235 **FAECAL EGG COUNT**

236 Table 4 shows the faecal egg count (egg/gram) of sheep among the dietary treatments. The graphical  
237 presentations are obtained in Figure 1.

238 At the onset of the experiment, the faecal egg count of the animals were 0% NLM (800.00), 5% NLM  
239 (833.33), 10% NLM (533.33) and 15% NLM (533.33) which reduced ( $P<0.05$ ) at the end of week 12.

240 By the end of week 12, animals in 5% NLM (100), 10% NLM (133.33) and 15% NLM (113.33) showed a  
241 reduction in FEC; NLM administered in this study caused a significant reduction in the worm burden of the  
242 sheep while the animals in 0% NLM (533.33) which is the control diet were not effectively dewormed. The  
243 study showed that all the animals were naturally and heavily infested with worms at the beginning of the  
244 experiment. Administration of the neem leaf meal shows a significant reduction ( $P<0.05$ ) in the faecal egg  
245 counts of the animals.

246 At the end of this study, there was a significant reduction in FEC of animals supplemented with NLM  
247 based concentrate. The reduction in Faecal egg count of animals in this study corroborates earlier  
248 findings of Chandrawathani *et al.*, (2000) which reported 82% reduction in worm eggs in animals fed fresh  
249 neem leaves *ad libitum* and a further trial on a limited number of animals showed that neem produced a  
250 limited worm burdens (Chandrawathani *et al.*, 2002).

251 In another study, Chandrawathani *et al.*, (2006) evaluated the anthelmintic effect of Neem on nematode  
252 parasites of sheep, the result of study shows that for FEC there was significant difference between the  
253 control group and the treated group, worm burden estimations showed that the number of parasites was  
254 significantly higher in the control group compared to the treated group. This result indicated that feeding  
255 neem has an effect on the worm numbers of sheep. The result in this study contradicts the study  
256 conducted by Khadijah *et al.*, (2005) on the use of fresh Neem which showed no significant difference in  
257 faecal egg count compared with control sheep, although the control sheep had higher mean faecal egg  
258 counts.

259 This result may be affected by feeding systems such as free pasture grazing on contaminated pastures  
260 as the animals are constantly challenged with infective larvae from pasture, so faecal egg counts may  
261 increase.

262 Results of highly significant reduction in the EPG count in lambs fed Neem leaves were also reported by  
263 Arunachal *et al.*, (2002). However, Costa *et al.*, (2006) reported no anthelmintic activity while feeding the  
264 Neem leaves for three months to sheep. While Niezen *et al.*, (1998) observed reduction in EPG count of  
265 *Trichostrongylus species* by Sulla feeding in ewe lambs, Hordegen *et al.*, (2003) also observed a  
266 reduction in egg counts of *Haemonchus concortus* with the seeds of Neem. Kahiya *et al.*, (2003) also  
267 observed similar decrease in EPG counts feeding *Acacia karoo* diets. However, Pietrosevoli *et al.*,  
268 (1999) did not find any differences in EPG count by feeding Neem leaves up to 40% level as blocks in  
269 calves. Similar to the effectiveness of neem leaves in lowering the worm count (Niezen *et al.*, 1998).  
270 Hordegen *et al.*, (2003) also reported reductions in worm burden while feeding sulla and seeds of neem

271 respectively. The variability in faecal egg counts within the NLM fed group may be due to differences in  
272 terms of physiological conditions of each animal and its ability to utilize the medicinal properties in neem.

273 **TABLE 4: Faecal egg count (egg/gram) of West African Dwarf sheep fed concentrate**  
274 **supplement containing varying inclusion level of neem leaf meal**

Weeks	0% NLM	5% NLM	10% NLM	15% NLM	±SEM
0	800	833	533	533	
3	633 <sup>a</sup>	533 <sup>b</sup>	466 <sup>c</sup>	600 <sup>a</sup>	83.33
6	600 <sup>a</sup>	333 <sup>c</sup>	366 <sup>c</sup>	400 <sup>b</sup>	5.53
9	566 <sup>a</sup>	122 <sup>d</sup>	233 <sup>c</sup>	333 <sup>b</sup>	3.33
12	533 <sup>a</sup>	100 <sup>b</sup>	133 <sup>b</sup>	113 <sup>b</sup>	9.17
<b>% FEC Reduction</b>	33.38 <sup>b</sup>	88.00 <sup>a</sup>	75.05 <sup>a</sup>	78.80 <sup>a</sup>	3.21

275 <sup>a, b, c</sup> mean values followed by the same letter in the same row are not significantly different (P≤ 0.05)

276 NLM- Neem leaf meal; FEC; Faecal egg count

277

278


279

280

281

282

283



284

285 **Figure 1: The average initial and final faecal egg count of sheep fed neem leaf meal**

286

## 287 **CONCLUSION**

288 Animals supplemented with NLM based concentrate also had a significant reduction in their faecal egg  
 289 count compared to the control treatment. However, animals on 5% NLM had the highest % faecal egg  
 290 count reduction value. Sheep on 5% NLM had the best haematological values for packed cell volume,  
 291 haemoglobin concentration and red blood cell, at the post haematology than other diets. Instead of  
 292 farmers using anti-parasitic drugs or anthelmintics to control internal parasites in their small ruminant  
 293 flocks which have residual effect on the populace consuming the meat of these animals, possible  
 294 anthelmintic potential of medicinal plants such as Neem tree (*Azadirachta indica*) can be exploited.

## 295 **REFERENCES**

296 Abdel Hameed, A. A., Salid, A. M., Fadel Elseed, A. M and Amasab, E. O. Effect of Feeding Untreated or  
 297 Urea Treated Groundnut Hull Supplemented with Different Protein Sources on Blood Parameters of  
 298 Sudan Desert Lambs. 2011; *Online Journal of Animal and Feed Research*. Pp 40-46.

299

300 Abu, O. A., Rabo, J. S., Onifade, A. A and Danny, C. B. Blood Composition and Histological Changes in  
 301 the Gastro-intestinal Tract of Rabbits Fed Untreated Rice Diets. 1998; *Proceedings of Annual Conference*  
 302 *of Nigeria Society Animal Production*, 23: 144-147.

303 Ajala O. O, Oyeyemi M. O, Oke A. O and Alakpa C. O. Haematological and biochemical parameters in  
304 West African dwarf (WAD) bucks fed diets containing *Milletia thonningii*. 2000; *African Journal of*  
305 *Biomedical Research*, 3; 121–124.

306

307 Akinyemi, A. F., Alokun, J. A and Fajemisin, A. N. Digestibility, Nitrogen Balance and Haematological  
308 profile of West African Dwarf sheep fed Dietary Levels of *Moringa oleifera* Supplement to *Panicum*  
309 *maximum*. 2010; *Journal of Animal Sciences*. 6: 634-643.

310 Arunachal, P.K., K. Karunanithi and R. Narendrababu, Comparative study on anthelmintic efficacy of  
311 neem products and praziquantel in sheep. 2002; *Indian Journal of Small Ruminant*. 8: 131-132.

312 Belewu, M.A and Ogunsola, F.O. Haematological and Serum Indices of goat fed Fungi Treated *Jatropha*  
313 *curcas* Kernel Cake in a mixed ration. 2010 ; *Journal of Agricultural Biotechnology and sustainable*  
314 *development*, 2(3) :035-038.

315 Campbell, T.W., and Coles, E.H. Avian Clinical Pathology.p. 279. In coles, E.H (ed.) *Veterinary Clinical*  
316 *Pathology*1986; 4<sup>th</sup> ed. W.B. Saunders Co., Philadelphia, Pennsylvania, USA.

317

318 Chandrawathani, P., Adnan, M. and Zaini, C. M. Preliminary Studies on Neem (*Azadirachta indica*) as an  
319 Alternative Anthelmintic for Sheep. 2000; Proceedings of the 12<sup>th</sup> Veterinary Association Malaysia  
320 Scientific Congress, Malaysia, Kuantan.

321

322 Chandrawathani, P., Adnan, M. and Waller, P.J. Anthelmintic Resistance in Sheep and Goat Farms on  
323 Peninsular Malaysia. 1999; *Veterinary Parasitology*. 82:305-310.

324

325 Chandrawathani, P., Brelvi, D., Nor Fasihah, S., Adnan, M., Jammah, O., Sani, R. A., Hoglund, J., and  
326 Waller, P. J. Evaluation of the Neem Tree (*Azadirachta indica*) as a Herbal Anthelmintic for Nematode  
327 Parasite Control in Small Ruminants in Malaysia. 2002; *Tropical Biomedicine* 19 (1&2): 41-48.

328

329 Chandrawathani, P., Yusoff, N., Wan, L. C., Ham, A. and Waller, P. J. Total Anthelmintic Failure to Control  
330 Nematode Parasites of Small Ruminants on Government Breeding Farm in Sabah, East Malaysia. 2004;  
331 *Veterinary Research Communication* 28:203-214.

332

333 Chandrawathani, P., Chang, K.W., Nurulainmi, R., Waller, P.J., Adnan, M., Zaini, C.M., Jammah, O.,  
334 Khadijah, S. and Vincent, N. Daily feeding of fresh Neem leaves (*Azadirachta indica*) for worm control in  
335 sheep. 2006; *Tropical Biomedicals* 23(1):23-30.

336

337 Coles, E. H. *Veterinary Clinical Pathology* (4<sup>th</sup> Edition). W. B. Saunders Company, Harcourt Brace  
338 Jovanovidi, Inc. 1986.  
339

340 Costa, C.T.C., C.M.L. Bevilaqua, M.V. Maciel, A.L.F. Camurca-Vasconcelos and S.M. Morais *et al.*  
341 Anthelmintic activity of *Azadirachta indica* against sheep gastrointestinal nematodes. 2006;  
342 *Veterinary Parasitol.*, 137: 306-310.  
343

344 Dacie, J. V and Lewis, S. M. *Practical Haematology* 6<sup>th</sup> Edition, Church Livington Edinburgh. Melbourne.  
345 New York. 1984; Pp 24-26.  
346

347 Daramola, J. O., Adeloye, Fatobass, A. A. and Soladoye, A. O., Haematological and Biochemical  
348 Parameters of West African Dwarf Goats. 2005; *Livestock Research for Rural Development*, Vol.17 No 8.  
349

350 David, S. *Approaches to Managing Gastrointestinal Nematode Parasites*. 2010; Agricultural Research  
351 Service, USDA 1999-2012.  
352

353 Duncan, D.B. Multiple Range and Multiple F- Tests *Biometrics* 1955; 11:1-42.  
354

355 Faye, D., Leak, S., Nouala, S., Fall, A and Losson, B. Effects of Gastrointestinal Helminth Infections and  
356 Plane of Nutrition on the Health and Productivity of F1 (West African Dwarf Sahelian) goat Crosses in  
357 the Gambian 2003; *Small Ruminant Research*, 50:153-161.  
358

359 Gatongi, P. M., Scott, M. E., Ranyan, S., Gathuma, J. M., Munyan, W. K., Cheruiyot, H and Prichard, R.  
360 K.. Effects of three Nematode Anthelmintic Treatment Regimes on Flock Performance of Sheep and Goats  
361 under Extensive Management in Semi-arid Kenya. 1997; *Veterinary Parasitol*, 68:323-326.  
362

363 Graczyk, S., Pliszczak-Krol, A., Kotonski, B., Wilczek, J. and Chmielak,. Examination of Haematological  
364 and Metabolic Changes Mechanisms of Acute Stress in Turkeys. 2003; *Electronic Journal of Polish*  
365 *Agricultural Universities* 6:1-10.  
366

367 Hordegen, P., H. Hertzberg, J. Heilmann, W. Langhans and V. Maurer,. The anthelmintic efficacy of five  
368 plant products against gastro-intestinal trichostrongylids in artificially infected lambs. 2003; *Veterinary*  
369 *Parasitol.*, 117: 51-60.  
370

371 Jawasreh, K., Awawdeh, F., Bani Ismail, Z., Al-Rawashdeh, O and Al-Majali, A.. Normal Haematology  
372 and selected Biochemical Values in Different Genetic Lines of Awassi ewes in Jordan. 2010; *The Internet*  
373 *Journal of Veterinary medicine* 7(2).

374  
375 Kahiya, C., S. Mukaratirwa and S.M. Thamsborg,. Effects of *Acacia nilotica* and *Acacia karoo* diets on  
376 *Haemonchus contortus* infection in goats. 2003; *Veterinary Parasitol.*, 115: 265-274.  
377  
378 Kaneko, J.J. Clinical Biochemistry of Domestic Animals.1997; Fifth ed. Academic Press, New York, USA,  
379 pp 885-905.  
380 Khadijah, S., Wahab, A. R., Chandrawathani, P. and Wan Zahari, M.. Pen study of Neem Pellets  
381 (*Azadirachta indica*) for Helminth Control in Sheep. 2005; Proceeding of 41st Annual Scientific Seminar  
382 "Natural Products in the Management of Tropical Diseases". Kuala Lumpur.  
383  
384 Merck,. The Merck veterinary Manual. 5<sup>th</sup> edition. Siegmund, O. H. 1979; (ed), Merck and Co. slnc.  
385 Rahway, New Jersey, USA. 167Pp.  
386  
387 Miller, J. E .M., Bahirathan, L. S., Lamarie, G. F., Hembry, T. M., Kearney and Barras, R. S. Epidemiology  
388 of Gastrointestinal Nematode Parasitism in Suffolk and Gulf Coast Native Sheep with Special Emphasis  
389 on Relative Susceptibility to *Haemonchus concortus* Infection1998; *Veterinary Parasitol* 74:55-74.  
390  
391 Niezen, J.H., H.A. Robertson, G.C. Waghorn and W.A.G. Charleston, Production faecal egg counts and  
392 worm burdens of ewe lambs which grazed contrasting forages. 1998; *Veterinary Parasitol.*, 80: 15-27.  
393  
394 Ogbuewu, I. P., Uchegbu, M. C., Okoli, I. C and Iloeje, M. U.. Assessment of Blood Chemistry, Weight  
395 Gain and Linear Body Measurement of Pre-puberal Buck Rabbits Fed Different Levels of Neem  
396 (*Azadirachta indica*) Leaf Meals. 2010; *Chilean Journal of Agricultural Research* 70(3):515-520.  
397  
398 Ososanya, T.O., Adewumi, M.K and Faniyi, T.O. Influence of water-washed Neem fruits (*Azadirachta*  
399 *indica* A. juss) on haematology and serum biochemical indices of West African Dwarf ewes. 2014;  
400 *American Journal of Experimental Agriculture*, 4(2):225-232.  
401  
402 Patra, A.K., Sharma, K., Dutta, N and Pattanik, A.K. Response of gravid Does to Partial Replacement of  
403 Dietary Protein by a leaf meal mixture of Clonical Chemistry. 2003; 3<sup>rd</sup> ed. Philadelphia: W.B Saunders  
404 Company. P. 809-861.  
405  
406 Pietrosevoli, S., R. Olavez and T. Montilla, Use of neem *Azadirachta indica* A. juss leaves in gastro-  
407 intestinal nematode control in grazing bovines. 1999; *Revista-de-la-Facultad- de-Agronomia*, 16: 220-225.  
408  
409 SAS,. Statistical Analysis System Users Guide. 1999; Statistical Version 8<sup>th</sup> edition, Statistical Analysis  
410 System Institute INC, Cary .N.C.

411  
412 Scott, J.L., Ketheesan, N and Summers, P.M. Leucocyte Population Changes in the Reproductive Tract  
413 of the ewe in Response to Insemination. 2006; *Reproductive Fertility and Development*. 18:627-634.  
414  
415 Taiwo, V. O and Ogunsanmi A. O.. Haematology, Plasma, Whole Blood and Erythrocyte Biochemical  
416 Values of Clinically Healthy Captive-reared Grey Duiker (*Sylvicapra grimmia*) and West African Dwarf  
417 Sheep and Goats in Ibadan, Nigeria. 2003; *Israel Journal of Veterinary Medicine*. 58(2-3).  
418 Talebi, A.S., Asri-Rezaei, R. Rozeh-Chai, and Sahraei, R. Comparative Studies on Hematological Values  
419 of Broiler strains. 2005; *International Journal of Poultry Science* 4(8):573-579.  
420  
421 Tambuwal, F. M., Agaie, B. M. and Bangama, A. Haematological and Biomedical Values of Apparently  
422 Healthy Red Sokoto Goats. 2002, In: Proceedings of the 27<sup>th</sup> Annual Conference of the Nigerian Society  
423 for Animal Production (NSAP), 17-21 March 2002, Federal University of Technology, Akure, Nigeria. Pp  
424 50-53.  
425  
426 Tinar, R., Akyol, C. V., Cirak, V. Y., Senlik, B and Bauer, C. Investigations on the Seasonal Patterns of  
427 Strongyle Infections in Grazing Lambs, and the Occurrences of Anthelmintic Resistance on Sheep and  
428 Goat Farms in Western Anatolia, Turkey 2005; *Parasitol Research*, 96:18-23.  
429  
430 Ukanwoko, A. I., Ironkwe, M. O. and Nnecha, C. Growth Performance and Haematological Characteristics  
431 of West African Dwarf Goats fed Oil Palm Leaf Meal-Cassava Peel Based Diets. 2013; *Journal of Animal*  
432 *Production Advances*. 3(1):1-5.  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449