

Short Research Article

Olfactory attraction of banana tree (Coleoptera: Curculionidae) to banana genotypes inoculated with entomopathogenic fungus

ABSTRACT

The species: *Cosmopolites sordidus* (Germar) (Coleoptera: Curculionidae) stands out as one of the main pests of banana. The damage to the crop is caused by the larvae of this pest, when they feed on the plant tissues. The aim of this study was to investigate the olfactory responses of *C. sordidus* for different varieties and the possible olfactory interference after application of *B. bassiana* fungus on the crop. The research was conducted at the Phytosanitary Clinic in the Agriculture Sector of the Center for Humanities, Social and Agrarian Sciences of the Federal University of Paraíba, located in Bananeiras - PB, from January to July 2017. The experimental design was completely randomized, with four treatments (banana varieties) and 11 repetitions each. The bioassays were distributed in two stages; The first stage was an evaluation of the attractiveness of banana genotypes and rhizome and pseudostem tissues. In the second, the attractiveness of *C. sordidus* to the tissues contaminated with the fungus *B. bassiana* was investigated. The Prata banana plantation was less attractive to *C. sordidus*, the highest preference was to the Nanica banana plantation. The most attractive tissue was pseudostem. Given the conditions under which the study was conducted, the banana plantation and the plant tissues analyzed present an olfactory influence under *Cosmopolites sordidus*. The Nanica banana plantation is the most susceptible to insect attack. The application of the fungus *Beauveria bassiana* on banana baits does not interfere with the power exerted by the nanica banana plantation under *C. sordidus*.

Keywords: Plant resistance, Coleobrocas, Olfactometer, Beauveria bassiana

1. INTRODUCTION

The species beetle: *Cosmopolites sordidus* (Germar) (Coleoptera: Curculionidae) stands out as one of the main banana pests, occurring in practically every region of the planet [1]. Injuries to the crop are caused by the larvae of this pest, which when feeding on plant tissues build galleries causing the interruption of the sap and the rotting of the plant and thus leading to decreased productivity.[2].

The use of systemic insecticides is considered one of the main forms of control for this pest [3]. However, one of the major problems encountered in chemical control is the translocation of the active ingredient of these insecticides via sap to the fruits, which are mostly marketed "in natura" [4]. In this context another form of control which has been widely used is the use of attractive baits, produced with pieces of pseudostem split in half. These baits are based

28 on the attraction exerted by the volatile substances present in the banana pseudostem and
29 rhizome. [5].

30 Use of attractive baits may be enhanced after spraying with entomopathogenic fungus:
31 *Beauveria bassiana* (Bals.) Vuill, this way the fungus acts against insects which served as
32 parasite spreading agents to other parts of the banana plantations. The fungus *B. bassiana*
33 is one of the most effective fungi and studied in biological control. This fungus as well as
34 other entomopathogenic fungi penetrate the host via the integument, causing the death of
35 insects due to mycotoxin production, and due to vegetative growth promoting mechanical
36 blockage of the digestive tract and other physical damage due to mycelial growth. [6].

37 Several studies have been performed to detect *C. sordidus* olfactory preference to different
38 banana varieties. However, information on the olfactory preference of these insects for
39 varieties treated with entomopathogenic fungi is rare or almost nil.

40 Thus, the objective of this study was to investigate the *C. sordidus* olfactory responses for
41 different varieties and the possible olfactory interference after *B. bassiana* fungus application
42 in the culture.

43

44 **2. MATERIAL AND METHODS**

45

46 The research was carried out at the Phytosanitary Clinic in the Agriculture Sector of the
47 Center of Human, Social and Agrarian Sciences of the Federal University of Paraíba, located
48 in Bananeiras - PB. At the same site, *C. sordidus* specimens were obtained. The capture
49 took place by means of tile baits, made from pieces of banana pseudostems. The insects
50 were kept in the laboratory in plastic containers, measuring approximately 10 cm high by 80
51 cm wide, with 1 cm radius perforations on the sides, containing as a food source and shelter
52 fresh pseudostem pieces, changed every 05 days The four banana genotypes analyzed
53 were: Pacovan, Nanica, Maçã and Prata (these are kinds of banana), from which two parts
54 of the plant (pseudostem and rhizome) were used. Later the most preferred variety was
55 inoculated with the fungus: *B. bassiana* for possible verification of the preference after
56 inoculation by the banana tree coleobroca. The strain of the entomopathogenic fungus *B.*
57 *bassiana* was isolated from a mummified specimen of *C. sordidus*, found in the Rural
58 community of Roma, Bananeiras district.

59 For evaluation of *C. sordidus* olfactory response to banana genotypes, the rhizome and
60 pseudostem of each genotype were reduced in dimensions of approximately 2 cm, arranged
61 in a multiple arena similar to that described by Botelho et al. [7] (Figure 1). Which features a
62 central arena and four side arenas as options to choose from. The bioassays were
63 performed during the night, a time associated with greater activity of the banana tree. [8].


64 The experimental design used was a completely randomized design. The bioassays were
65 composed of four treatments (banana species) containing 11 repetitions and distributed in
66 two stages; where the first one evaluated the attractiveness to the banana genotypes and
67 the rhizome and pseudostem tissues. In this evaluation the tissues and each vegetable were
68 grouped in isolation on the olfactometer, where two arenas were filled with fresh tissue of
69 one genotype and the other two by cotton wicks soaked with distilled water as a control.
70 Subsequently, four *C. sordidus* adults were placed in the central arena and remained for 40
71 minutes exposed to the volatiles released by the tissues of the analyzed plants.

72 In the second stage to verify the attractiveness of *C. sordidus* to tissues contaminated with
73 the fungus: *B. bassiana*, the arenas were filled with fresh tissue (pseudostem) of the Nanica
74 banana plantation, this time using only, detected in the first stage of the bioassays as the

75 genotype and the tissue with the highest preference for the insect. For the contamination of
 76 banana tissues, the fungus was diluted in water and the tissues dipped in this solution for
 77 conidia adherence.

78
 79 The parameters evaluated were preference, non-preference and individuals with no
 80 response to odors. For this, the insects found in three conditions were quantified: on fresh
 81 rhizome and pseudostem tissues, found on the control and those that remained in the
 82 release arena. Insects not in these conditions were disregarded.

83 The results for banana olfactory response to volatile banana genotypes were analyzed using
 84 the non-parametric X² test (chi-square), which is composed by the formula: $X^2 = \sum(O-E)^2 / E$.
 85



86
 87 **Figure 1. Multiple choice arena for olfactometric analysis of *C. sordidus***
 88

89 **3. RESULTS**

90
 91 The results for *C. sordidus* olfactory preference (Table 1) showed a low insect attractive
 92 effect on maçã banana variety plant tissues, and the observed frequency was lower than
 93 expected in both tissues analyzed.

94
 95
 96 **Table 1. Attractiveness of *C. sordidus* to maçã banana variety plant tissues offered in a**
 97 **separated way**

Maçã banana plantation						
Observed Frequency (Expected) (%)						
Insect	Choice option	Preference	No preference	No answer	X ²	Value de (P)
<i>Cosmopolites sordidus</i>	Pseudocaulis	36,36 (40)	18,18 (40)	45,45 (20)	44,6	(0,001)
	Rizoma	18,18 (40)	9,09 (40)	72,72 (20)	174,8	(0,001)

98
 99 For the prata banana plantation the results indicate a high influence of pseudostem on *C.*
 100 *Sordidus*, when they showed an olfactory preference of 81.81%, while the expected was
 101 around 40%. However, the rhizome of this same variety was not attractive to the insect, and
 102 the expected frequency was higher than the observed frequency (Table 2).

103 **Table 2. Attractiveness of *C. sordidus* to prata banana variety plant tissues offered in a**
 104 **separated way**

Prata banana Plantation						
Observed Frequency (Expected) (%)						
Insect	Choice option	Preference	No preference	No answer	X ²	Value of (P)
<i>Cosmopolites sordidus</i>	Pseudocaulis	81,81 (40)	9,09 (40)	9,09 (20)	73,53	(0,001)
	Rizoma	9,09 (40)	18,18 (40)	72,72 (20)	174,8	(0,001)

105

106 Both tissues of Nanica banana plantation were attractive to *C. Sordidus*, obtaining
 107 preference values higher than the observed frequency. The pseudostem showed a
 108 preference of 81.81% and the rhizome 54.54%, the expected frequency for the tissues was
 109 40% (Table 3).

110 **Table 3. Attractiveness of *C. sordidus* to plant tissues of the Nanica banana variety**
 111 **offered in isolation.**

112

Nanica banana Plantation						
Observed Frequency (Expected)(%)						
Insect	Choice option	Preference	No Preference	No answer	X ²	Value of (P)
<i>Cosmopolites sordidus</i>	Pseudocaulis	81,81 (40)	9,09 (40)	9,09 (20)	73,53	(0,001)
	Rizoma	54,54 (40)	0,09 (40)	36,36 (20)	58,48	(0,001)

113

114 PTo cultivate Pacovan, the results indicate that the Insects presented a higher Preference
 115 for the pseudostem, which obtained 45.45% in the observed frequency against 40% of the
 116 expected frequency (Table 4).

117 **Table 4. *C. sordidus* attractiveness to plant tissues of the Pacovã variety offered alone**

Pacovan Plantation						
Observed Frequency (Expected)(%)						
Insect	Choice option	Preference	No Preference	No answer	X ²	Value of (P)
<i>Cosmopolites sordidus</i>	Pseudocaulis	45,45 (40)	9,09 (40)	45,45 (20)	57,01	(0,001)
	Rizoma	36,36 (40)	0,09 (40)	54,54 (20)	99,80	(0,001)

118

119 As it was observed in Table 5, *C. sordidus* showed Preference by pseudostem tissues when
 120 they were inoculated by the fungus *B. bassiana*.

121 **Table 5. Attractiveness of *C. sordidus* in tissues of inoculated Nanica Banana**
 122 **plantation and without application of the fungus *Beauveria bassiana*.**

Insect	Observed Frequency (Expected)(%)				Value of (P)
	Pseudostem with application of <i>B. bassiana</i>	X ²	Pseudostem without application	X ²	
<i>Cosmopolites sordidus</i>	60,0 (50)	2,0	40,0 (50)	2,0	(0,001)

123

124 4. DISCUSSION

125

126 Among the various managements used for the control of agricultural pests, varietal
 127 resistance has been researched and pointed as a viable alternative in the control of *C.*
 128 *sordidus* [9, 10]. We evidenced in this study the olfactory influence exerted by different plant
 129 tissues and genotypes, analyzed on *C. sordidus*. The attractiveness of Insect by the plant is
 130 produced due to the secondary volatile compounds released by the plant as semiochemicals
 131 present in the rhizome and pseudostem. (11, 12].

132 Facundo et al [13] states that the various banana genotypes have different volatile
 133 compounds that vary according to variety. According to Oliveira et al., [14], the volatiles
 134 found in banana genotypes, whether varieties or hybrids, have the same attractiveness for
 135 *C. sordidus*. However, there is variation in the composition and concentration of volatiles that
 136 may vary among banana genotypes, reflecting the responses of *C. sordidus* adults to each
 137 genetic material [13, 15].

138 We corroborate in this study the high attractiveness of the volatile compounds of Nanica
 139 Banana plantation for both plant tissues analyzed. The Nanica Banana Plantation is
 140 considered one of the most susceptible to attack by this insect, with losses of up to 80% of
 141 production, these losses are also affected in the Prata banana plantation, with a smaller
 142 proportion around 30% [16].

143 Infestations of this pest in banana plants can be reduced by planting less attractive
 144 plantations, which resulted in lower insects demand for the plant and consequently less
 145 oviposition and emergence of new individuals [1]. Another efficient type of control widely
 146 used by producers is the use of entomopathogenic fungi traps. The use of these fungi for the
 147 biological control of banana broth as with other agricultural pests is a viable alternative that
 148 reduces environmental impacts caused by the use of chemicals such as insecticides [16].
 149 The use of alternative pest controls also reduces the use of chemicals, reducing the amount
 150 of toxic waste to humans in banana consumption and at the time of application [17].

151

152 5. CONCLUSION

153

154 Given the conditions under which the study was conducted, it can be concluded that the
 155 banana plantations and plant tissues analyzed have an olfactory influence under
 156 *Cosmopolites sordidus*. The Nanica banana plantation is the most susceptible to Insect
 157 attack. The application of the fungus *Beauveria bassiana* in banana baits does not affect the
 158 power exerted by the nanica banan plantation under *C. sordidus*.

159

160

161 **COMPETING INTERESTS**

162

163 Authors have declared that no competing interests exist.

164

165

166 **REFERENCES**

167

168 1. Gold CS, Pena JE, Karamura EB. Biology and integrated pest management for the
169 banana weevil *Cosmopolites sordidus* (Germar) (Coleoptera: Curculionidae).
170 Integrated Pest Management Reviews. 2001;6:79-155. Springer Nature.
171 <http://dx.doi.org/10.1023/a:1023330900707>.

172 2. Dassou AG, Carval D, Dépigny S, Fansi G, Tixier P. Ant abundance and
173 *Cosmopolites sordidus* damage in plantain fields as affected by intercropping.
174 Biological Control. 2015; 81:51-57. Elsevier BV.
175 <http://dx.doi.org/10.1016/j.biocontrol.2014.11.008>.

176 3. Almeida AMB, Batista FA, Tavares FM, Leite LG. Seleção de isolados de *Beauveria*
177 *bassiana* para o controle de *Cosmopolites sordidus* (Germar, 1824) (Coleoptera:
178 Curculionidae). Arquivos Instituto Biológico. 2009; 76: 489-493.

179 4. Moreira FJC, Araújo AB, Silva VF, Luna NS, Araújo OP, Braga RDS. Controle de
180 *Cosmopolites sordidus* (Coleoptera: Curculionidae) com os fungos
181 entomopatogênicos *Beauveria bassiana* e *Metarhizium anisopliae* em
182 banana. Revista Verde de Agroecologia e Desenvolvimento Sustentável.
183 2017;12:366-373. Grupo Verde de Agroecologia e Abelhas.
184 <http://dx.doi.org/10.18378/rvads.v12i3.4538>.

185 5. Mendonça FAC, Vilela EF, Eiras ÁIE, Sant'Ana AEG. Resposta de *Cosmopolites*
186 *sordidus* (Germar) (Coleoptera, Curculionidae) aos voláteis da planta hospedeira e
187 de adultos coespecíficos em olfatômetro. *Revista Brasileira de Zoologia*.
188 1999;16:123-128. <https://dx.doi.org/10.1590/S0101-81751999000600014>.

189 6. ALVES SB. Controle microbiano de insetos. São Paulo: Manole, 1986.

190 7. Botelho PSM, Silveira Neto S, Barbin D, Boreges CG. Teste de atração de *Musca*
191 *domestica* L. com luzes de diferentes comprimentos de onda. O Solo, 65:42-45.
192 1973.

193

194 8. Gold CS, Pinese B, Peña JE. Pests of banana. In: Peña JE, Sharp JL & Wysoki M
195 (Eds.) Tropical fruit pests and pollinators: biology, economic importance, natural
196 enemies and control. Wallingford, CABI Publishing. 2002; 3-56.

197 9. Kiggundu A, Gold CS, Labuschagne MT, Vuylsteke D, Louw S. Components of
198 resistance to banana weevil (*Cosmopolites sordidus*) in *Musa* germplasm in
199 Uganda. Entomologia Experimentalis et Applicata. 2007;122:27-35.

200 10. Sadik K, Nyine M & Pillay M. A screening method for banana weevil (*Cosmopolites*
201 *sordidus* Germar) resistance using reference genotypes. African Journal of
202 Biotechnology. 2010 9:4725-4730.

- 203 11. Budenberg WJ, Ndiege IO, Karago, FW, Hansson BS. Behavioral and
204 electrophysiological responses of the banana weevil *Cosmopolites sordidus* to host
205 plant volatiles. *Journal of Chemical Ecology*. 1993;19:267-277.
- 206 12. Cerda H, López A, Sanoja O, Sánchez, JAFFÉ K. Atracción olfativa de
207 *Cosmopolites sordidus* Germar (1824) (Coleoptera: Curculionidae) estimulado por
208 volatiles originados en musaceas de distintas edades y variedades genómicas.
209 *Agronomia Tropical*. 1996;46:413-429.
- 210 13. Facundo HVV, Garruti DS, Dias CTS, Cordenunsi BR, Lajolo FM. Influence of
211 different banana cultivars on volatile compounds during ripening in cold storage.
212 *Food Research International*. 2012;49:626-633.
213 <https://doi.org/10.1016/j.foodres.2012.08.013>.
- 214 14. Oliveira FT, Neves PMOJ, Bortolotto OC, Ventura,U. Respostas olfativas do
215 moleque-da-bananeira (Coleoptera: Curculionidae) para diferentes genótipos de
216 bananeira. *Revista Ceres*. (2018); 65(4): 329-337. <https://dx.doi.org/10.1590/0034-737x201865040005>.
- 218 15. Pino JA, Flebles Y. Odour-active compounds in banana fruit cv. Giant Cavendish.
219 *Food Chemistry*. 2013;141:795-801.
- 220 16. Fancelli M, Dias A B, Jesus SC, Delalibera I, Nascimento AS, Silva S. Controle
221 Biológico de *Cosmopolites sordidus* (Germ.) (Coleoptera: Curculionidae) pelo Fungo
222 *Beauveria bassiana* (Bals.) Vuill. *Embrapa Mandioca e Fruticultura*. 2004; 3.
- 223 17. Rossetti NCS. Controle biológico e comportamental do moleque-da-bananeira.
224 Casa do Produtor Rural – ESALQ/USP. 2013; accessed 09 october 2019.
225 Available:<http://www.esalq.usp.br/cprural/boapratica/mostra/85/control-e-biologico-e-comportamental-do-moleque-da-bananeira>.
226
227