

1 **Abundance and morphometric study of some Lizards (Agama lizard, Skinks and Wall**
2 **gecko) in the University Community in Nigeria: Obafemi Awolowo University as a case**
3 **study**

4
5
6 **Abstract**

7 This study investigated the abundance and some morphometric parameters of lizards (*A.*
8 *agama*, *H. brookii*, *T. affinis* and *T. quinquetaeniata*) in the University campus, Ile-Ife. The
9 specimens were collected in five different locations in the University campus. The samples
10 were collected with sweep nets in all the locations. A total of 624 species of lizards (324 of *A.*
11 *agama*, 185 of *H. brookii*, 60 of *T. affinis* and 55 of *T. quinquetaeniata*) were caught in all the
12 locations and external body measurement was carried out with the help of calliper. *A. agama*
13 was found to be abundant in all the locations followed by *H. brookii*. The number of Skinks
14 (*T. affinis* and *T. quinquetaeniata*) in the residential area was low but high in University parks
15 and garden and markets. The morphometric parameters showed that *A. agama* was different
16 in all the parameters measured compared to the rest specimens and it was expected since the
17 specimens were not of the same family except the Skinks (*T. affinis* and *T. quinquetaeniata*).
18 There were similarity in values of some of the measured body parts of *H. brookii*, *T. affinis*
19 and *T. quinquetaeniata*. The Principal Components Analysis (PCA) showed that all the
20 specimens differ in Tail length, Tail width, and Trunk length. In conclusion, the abundance of
21 the Lizards in the University call for concern because their waste can contaminate food and
22 cause infection since they are hosts to a number of parasites.

23
24 **Keywords:** Infection, Parasite, Trunk length, Head length

25
26 **1.0 Introduction**

27 The parasite causing infectious disease which can be commonly found in some lizards living
28 around us (1) are of great fear to human existence. Parasites are found in lizards which are

29 found everywhere in many of the tropical climates of the world. Although, most lizard
30 species are harmless to humans unless provoked (2), but they can cause diseases if they lay
31 eggs or urinate into human food.

32 In some parts of the world, lizards and other reptiles such as snakes, crocodiles are used for
33 food (3). In Nigeria, the clouded-monitor lizard serves as a source of meat for people who
34 likes games. Clouded-monitor lizard help in insect control in some agricultural areas since
35 they feed on them. In Africa, the lizards commonly found are Agama lizard, Skinks, Gecko,
36 Chameleons, Monitor lizard, Alligator lizard (4).

37 *Agama agama* was reported to serve as transport and reservoir host to several protozoan and
38 helminth parasites (5). Parasites from most reptiles can be transferred into a human by
39 carefree attitude. Humans can be infected with *Raillietiella* species, by contaminating their
40 hands and food with the eggs, faeces or saliva of the reptile (6). Gecko (*Hemadactylus sp*) are
41 house lizards commonly found in human dwelling in tropical countries. They are nocturnal
42 lizards which feed mainly on insects and termites (7). Oluwafemi *et al.* (8) reported the
43 presence of *Raileteilla frenatus* and *Paraphayngodon sp* in Wall gecko (*Hemidactylus*
44 *frenatus*) species caught in Ile-Ife. These parasites can cause diseases for human if ingested
45 into the human system.

46 Skinks of the genus *Mabuya* is found around the world and it contains about 90 species which
47 are distributed around Sub-sahara Africa (9). Like all vertebrates, skinks are susceptible to
48 parasitism in their niche. *Spirura sp* a nematode was found in skink (*Mabuya*
49 *quinquetaeniata*) collected in Ile-Ife and Ibadan by Oluwafemi *et al.* (8). In Nigeria, there
50 have been various studies conducted on the parasites of lizards and other reptiles (10, 11 and
51 12). It is important to study the population of these lizards in our locality to determine the
52 best method to control them and reduce the risk of contamination of our foods.

53 Obafemi Awolowo University, Ile-Ife, Osun State where the research was conducted was one
54 of the University with a large number of students in Nigeria. Similar research was conducted

55 on the bait preference of Black rat (*Rattus rattus*) when the students' hostel was infected by
56 black rat (13). This study was conducted in the University (Obafemi Awolowo University) to
57 determine the most prevalence lizards (*A. agama*, Skinks and Wall gecko) in various part of
58 the University. This is aimed at controlling the lizards in other to reduce the risk of infection
59 among the University populaces.

60

61 **3.0 Materials and methods**

62 **3.1 Study Location**

63 This research work was carried out in Obafemi Awolowo University student's hostels,
64 academic area and staff quarters, Ile Ife. The university is located between Latitudes 7°26'N
65 and 7°32'N and between Longitudes 4°31'E and 4°35'E. The landmass is 5506 hectares with
66 an altitude of 300 m above sea level.

67

68 **3.2 Materials**

69 **3.2.1 Materials used in the research field**

70 Sweep net: For trapping the specimens, Chloroform: To anesthetize the specimens, Killing
71 jar: Where the specimens are kept immediately after collection, Cotton wool: Placed inside
72 the killing jar for a smooth landing for the anesthetized specimens. Measuring ruler, Thread,
73 Rubber gloves, Harvard trip balance, Camera

74

75 **3.2.2 Method of collection**

76 Agama lizards (*Agama agama*) and Skinks (*Trachylepis affinis*; *Trachylepis quinquetaeniata*)
77 were caught by using a sweep net to cover them, and they were put inside killing jar
78 containing chloroform. Wall geckos (*Hemidactylus brookii*) were caught in the night using a
79 broom. The broom was used to brush them into the sweep net after which they were
80 transferred into the killing jar containing chloroform. The chloroform was used to anesthetize
81 the specimens until they are taken into the laboratory.

82

83 **3.3 Identification of specimens**

84 The Agama lizard (*Agama agama*) was identified by the descriptions given by Harris (14).

85 The lizard was said to have a white underside, brown back limbs and a tail with a light stripe
86 down the middle. Breeding males of this subspecies have brilliant orange heads, and an
87 indigo blue or black body and legs. Their tail is bluish-white at the base and has an orange
88 middle area and black tail tip. The non-breeding male is paler in colour and might not have
89 the orange on the head. Females are brown and have olive green colour on their backs with
90 some barring marks. Wall gecko (*Hemidactylus brookii*) was identified with some
91 specification given by Bauer et al. (15). The snout of the specimen is longer than the distance
92 between the eye and the ear opening, nearly twice the diameter of the orbit; the forehead is
93 concave.

94 Skinks (*Trachylepis affinis*) has a yellow or white venter, not bright blue-green. Skink
95 (*Trachylepis quinquetaeniata*) males have the side of the head yellow and the throat black.
96 Juveniles and females have five white longitudinal stripes. Identification of skink species was
97 done using identification guides by Branch (16) and Spawls *et al.*, (17).


98

99 **3.4 Data collection**

100 The captured specimens (that have been anesthetized) were taken into the laboratory for sex
101 identification and measurement of morphometric parameters. Specimens without sexual
102 dimorphism (Agama Lizard) the sex was determined, they were tagged male or female before
103 being weighed and measured. In the laboratory, the sex of the geckos was determined using
104 the presence of pores and hemi penal bulges at the base of the tail to identify the males while
105 the absence of those aforementioned identified females. The sex of the skinks was determined
106 by dissecting them and checking for the presence of ovaries. Those with ovaries are females
107 while those without ovaries are males.

108 The following measurements were taken in cm:

- 109 1. Snout-vent length (SVL): from the tip of snout to anterior end of the cloaca.
- 110 2. Head height (HH): height of the head.
- 111 3. Head length (HL): ventral measurement from the tip of the lower jaw to immediately
- 112 posterior to the jaw.
- 113 4. Head width (HW): the widest portion of the head anterior to the ear.
- 114 5. Tail length (TL): from the anterior end of the cloaca to the tip of the tail
- 115 6. Tail width (TW): measured at the base of the tail from one side to another
- 116 7. Trunk length (TRL): from where the forelimb originates to where the hind limb
- 117 originates.
- 118 8. Mouth opening (MO): from snout to posterior border of the last supralabial scale
- 119 The weight of all specimens was measured in grams using the Harvard trip weighing balance.



120

121 **Figure 1:** Diagram showing some of the measured morphometric parameters on the Lizards.

122 Sources: (a & b) Modified from Kaliontzopoulou *et al.* (18), and (c) Modified from Uyeda *et*

123 *al.* (19)

124

125 3.5 Statistical Analysis

126 One-way analysis of variance (ANOVA) was used to determine the significant difference

127 between the means, while the significant mean was separated at $p \leq 0.05$ using Least

128 Significant Difference (LSD) test from System Analysis Software (SAS Institute, 20).

129 Principal Component Analysis (PCA) was carried out with IBM SPSS 20th version.

130 **4.0 Results**

131 The abundance of four types of lizards (*Agama agama*, *Hemidactylus brookii*, *Trachylepis*
 132 *affinis* and *Trachylepis quinquetaeniata*) studied at five different locations (University
 133 Hostels, University staff quarters, University Library, University Parks and garden and
 134 markets) are shown in Table 1. Among the four lizards studied, *A. agama* had the highest
 135 population in all locations (57.53, 48.02, 53.85, 54.33, and 49.30%), followed by *H. brookii*
 136 (30.97, 36.72, 30.77, 19.69, and 28.17%). Skinks (*T. affinis* and *T. quinquetaeniata*) have the
 137 least percentage from all the locations.

138 **Table 1:** The morphometric parameters of male of the four lizard species (*Agama agama*,
 139 *Hemidactylus brookii*, *Trachylepis quinquetaeniata*, and *Trachylepis affinis*) caught on
 140 Obafemi Awolowo University, Ile-Ife.
 141

Locations	Specimen name	Male	Female	Total	Percentage total (%)
University Hostels	Agama lizards (<i>A. agama</i>)	25	40	65	57.53
	Wall geckos (<i>H. brookii</i>)	20	15	35	30.97
	Skinks (<i>T. affinis</i>)	3	5	08	7.08
	Skinks (<i>T. quinquetaeniata</i>)	2	3	05	4.42
	Total	50	63	113	100
University Staff Quarters	Agama lizards (<i>A. agama</i>)	30	55	85	48.02
	Wall geckos (<i>H. brookii</i>)	25	40	65	36.72
	Skinks (<i>T. affinis</i>)	7	9	16	9.04
	Skinks (<i>T. quinquetaeniata</i>)	5	6	11	6.22
	Total	67	110	177	100
University Library	Agama lizards (<i>A. agama</i>)	15	20	35	53.85
	Wall geckos (<i>H. brookii</i>)	12	8	20	30.77
	Skinks (<i>T. affinis</i>)	2	4	06	9.23
	Skinks (<i>T. quinquetaeniata</i>)	1	3	04	6.15
	Total	30	35	65	100
University Parks & Garden	Agama lizards (<i>A. agama</i>)	27	42	69	54.33
	Wall geckos (<i>H. brookii</i>)	14	11	25	19.69
	Skinks (<i>T. affinis</i>)	7	8	15	11.81
	Skinks (<i>T. quinquetaeniata</i>)	7	11	18	14.17
	Total	55	72	127	100
University Markets	Agama lizards (<i>A. agama</i>)	28	42	70	49.30
	Wall geckos (<i>H. brookii</i>)	18	22	40	28.17
	Skinks (<i>T. affinis</i>)	7	8	15	10.56

Skinks (<i>T. quinquetaeniata</i>)	8	9	17	11.97
Total	61	81	142	100

142

143 In all the locations, the highest number of Agama lizard was collected in the University
 144 Hostels (57.53%) (Figure 2) followed by University parks and garden (54.33%), followed by
 145 University Library, followed by University markets (49.30%). The least percentage of Agama
 146 lizard was collected in University staff quarters (48.02%).

147

148 The highest value of *H. brookii* was collected in University staff quarters (36.72%), followed
 149 by University Hostels (30.97%) and University Library (30.77%). The least was collected in
 150 University parks and garden (19.69%). Skinks (*T. affinis* and *T. quinquetaeniata*) abundance
 151 was low in University Hostels (7.08 & 4.42%), University staff quarters (9.04 & 6.22%) and
 152 University Library (9.23 & 6.15%). The highest percentage of Skins (*T. affinis* and *T.*
 153 *quinquetaeniata*) were from University parks & garden (11.81 & 14.17%) and University
 154 market (10.56 & 11.97%).

155 The ratio of male to female of all the lizards did not follow a specific pattern in that female
 156 ratio was higher than male in *A. agama* in all the locations while the male ratio was higher
 157 than female in *H. brookii* collected in all the locations. The ratio of female Skins (*T. affinis*
 158 and *T. quinquetaeniata*) collected in all the locations were higher than the male in this study.

159

160 Table 2 showed the morphometric parameters measured on the male specimens collected in
 161 all the locations. *A. agama* had the highest average body weight (68.38 g) followed by
 162 *Trachylepic quinquetaeniata* (23.55 g). The bodyweight of *Trachylepic affinis* (11.68 g) was
 163 higher than that of *Hemidactylus brookii* (8.99 g). The Height of the head (HH) of all the
 164 male specimens followed the same pattern with the bodyweight with *A. agama* having the
 165 highest average value (1.10 cm) followed by *T. quinquetaeniata* (0.90 cm). *T. affinis* and *H.*
 166 *brookii* have 0.75 & 0.55 cm respectively.

167

168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204




Figure 2: Percentage of four different lizards (*Agama Lizard (Agama agama)*, *Wall gecko (Hemidactylus brookii)*, *Skinks (Trachylepis quinquetaeniata & Trachylepis affinis)*) in various collections site on the University

- Keys**
- University Hostels
 - University Parks and garden
 - University Library
 - University Staff Quarters
 - University Market

205 **Table 2:** The morphometric parameters of male of the four lizard species (*A. agama*, *H. brookii*, *T. affinis* and *T. quinquetaeniata*) caught on
 206 Obafemi Awolowo University, Ile-Ife.
 207

Species	Statistics	BW (g)	HH (cm)	HL (cm)	HW (cm)	TL (cm)	TW (cm)	TRL (cm)	MO (cm)	SVL (cm)
AA	Mean	68.38 ^d ±1.2	1.10 ^d ±0.1	3.82 ^c ±0.1	3.45 ^b ±0.1	16.85 ^c ±0.5	0.86 ^a ±0.1	5.09 ^c ±0.4	1.99 ^c ±0.1	12.87 ^d ±0.2
	Range	50.4-80.8	0.7-1.5	2.8-4.5	2.5-4.0	11.6-20.9	0.4-1.6	3.0-8.0	1.3-2.6	11.2-14.6
HB	Mean	8.99 ^a ±0.3	0.55 ^a ±0.1	1.86 ^a ±0.1	1.16 ^a ±0.1	5.50 ^a ±0.3	0.65 ^a ±0.1	2.69 ^a ±0.1	1.04 ^b ±0.1	5.87 ^a ±0.1
	Range	7.4-12.4	0.4-0.8	1.6-2.4	0.6-1.5	4.0-7.0	0.4-0.9	1.7-3.1	0.6-1.7	5.4-6.4
TA	Mean	10.68 ^b ±0.4	0.75 ^b ±0.12	1.93 ^a ±0.2	1.15 ^a ±0.2	11.18 ^b ±0.4	0.71 ^a ±0.1	4.52 ^b ±0.3	0.89 ^a ±0.1	7.15 ^b ±0.5
	Range	10.1-14.0	0.5-1.5	1.1-2.6	0.6-2.0	9.3-12.9	0.6-1.1	3.5-5.4	0.5-1.3	5.4-9.4
TQ	Mean	15.55 ^c ±0.7	0.90 ^c ±0.1	2.15 ^b ±0.5	1.65 ^a ±0.1	13.00 ^b ±0.6	0.90 ^a ±0.1	4.10 ^b ±0.4	1.35 ^b ±0.1	9.20 ^c ±0.3
	Range	20.8-26.3	0.8-1.0	2.0-2.3	1.6-1.7	12.4-13.6	0.8-1.0	3.7-4.5	1.3-1.4	8.9-9.5

208 *Means within column with different Superscript are significantly different ($P \leq 0.05$) from each other

209 Footnotes

210 AA is *Agama agama* (Agama Lizard), HB is *Hemidactylus brookii* (Wall gecko), TA is *Trachylepis affinis* (Skinks) and TQ is *Trachylepis quinquetaeniata* (Skinks)

211 BW is the Body weight, HH is the Height of the head, HL is the Head length, HW is the Head width, TAL is the Tail length, TAW is the Tail width, TRL is the Trunk length,

212 MO is the Mouth opening and SVL is the Snout-vent length

213

214 There was no significant difference ($p > 0.05$) in Head length average value among three male
215 specimens (*H. brookii*, *T. affinis* and *T. quinquetaeniata*) measured when compared to *A.*
216 *agama* (3.45 cm). The Head width average value of male specimens collected in all the
217 locations showed that there was no significant difference ($p > 0.05$) between *H. brookii* and *T.*
218 *affinis*. There was a significant difference ($p < 0.05$) between the average Head width of *A.*
219 *agama* (3.45 cm) and *T. quinquetaeniata* (1.65 cm).

220 The Tail length (TL) average value of *A. agama* (16.85 cm) was the highest and *H. brookii*
221 (5.50 cm) have the least value of TL. There was no significant difference ($p > 0.05$) between
222 the TL of *T. affinis* and *T. quinquetaeniata*. The Tail width (TW) of all the specimens (*A.*
223 *agama*, *H. brookii*, *T. affinis* and *T. quinquetaeniata*) showed no significant difference at $p >$
224 0.05 , while the average value of TW for *T. quinquetaeniata* was the highest (0.90 cm) among
225 all the male specimens. *A. agama* has average TAW value of 0.86 cm followed by *T.*
226 *quinquetaeniata* (0.71 cm) and *T. affinis* (0.65 cm).

227 The average Trunk length (TRL) of *T. affinis* (4.52 cm) and *T. quinquetaeniata* (4.10 cm)
228 have no significant difference ($p > 0.05$), while average TRK of *A. agama* (5.09 cm) and *H.*
229 *brookii* (2.69 cm) showed a significant difference at $p < 0.05$. *H. brookii* have the list average
230 TRL value (2.69 cm) and *A. agama* have the highest value of TRL (5.09 cm). The average
231 value of Mouth opening (MO) of all the specimens was significantly differenced at $p < 0.05$.
232 *A. agama* has the highest value of MO (1.99 cm) and *T. affinis* have the least value of MO
233 (0.89 cm). *H. brookii* and *T. quinquetaeniata* have MO average values of 1.04 and 1.35 cm
234 respectively. The Snout-vent length (SVL) average of all the specimens showed a significant
235 difference at $p > 0.05$. *A. agama* has the highest value of 12.87 cm, followed by *T.*
236 *quinquetaeniata* (9.20 cm). *T. affinis* have an average value of 7.15 cm for SVL, while *H.*
237 *brookii* have an average value of 5.87 cm for SVL.

238

239 The values of morphometric parameters measured from female specimens (*A. agama*, *H.*
240 *brookii*, *T. affinis* and *T. quinquetaeniata*) were shown in Table 3. The average Body weight
241 (BW) of *A. agama* was the highest (35.23 g) followed by Skinks (*T. affinis* (11.67 g) and *T.*
242 *quinquetaeniata* (15.40 g)) and *H. brookii* (6.23 g). The of Height of head (HH) of all the
243 specimens collected in different locations showed a significant difference ($p < 0.05$) with *T.*
244 *quinquetaeniata* having the highest value (0.70 cm) and *A. agama* having the least (0.34 cm).
245 There was no significant difference ($p > 0.05$) among the average value of Head length (HL)
246 among three lizards (*H. brookii*, *T. affinis* and *T. quinquetaeniata*). There was a significant
247 difference ($p < 0.05$) in HL value between *A. agama* and the rest of the specimen. The
248 average value of the Head width (HW) of *A. agama* was the highest (2.15 cm) followed by *H.*
249 *brookii* (1.55 cm), followed by *T. affinis* (1.14 cm) and *T. quinquetaeniata* (1.06 cm).
250
251 The least value of Tail length (TL) was recorded from *H. brookii* (5.23 cm), while the highest
252 average TL was recorded for *A. agama* (14.15 cm). The average value of the Tail length (TL)
253 of the specimens showed a significant difference ($p < 0.05$) between *A. agama* (14.15 cm) and
254 *H. brookii* (5.23 cm). There was no significant difference ($p > 0.05$) in TL between the two
255 species of Skinks (*T. affinis* and *T. quinquetaeniata*). Tail width (TAW) followed the same
256 pattern as TL. There was a significant difference ($p < 0.05$) between TW of *A. agama* and *H.*
257 *brookii*, while there was no significant difference ($p > 0.05$) in TW between *T. affinis* and *T.*
258 *quinquetaeniata*.

259 **Table 3:** The morphometric parameters of female of the four lizard species (*A. agama*, *H. brookii*, *T. affinis* and *T. quinquetaeniata*) caught on
 260 Obafemi Awolowo University, Ile-Ife.
 261

Species	Statistics	BW (g)	HH (cm)	HL (cm)	HW (cm)	TL (cm)	TW (cm)	TRL (cm)	MO (cm)	SVL (cm)
AA	Mean	35.23 ^d ±1.6	0.34 ^a ±0.1	2.74 ^b ±0.2	2.15 ^c ±0.1	14.15 ^c ±0.3	0.39 ^a ±0.1	4.05 ^b ±0.2	1.51 ^c ±0.1	10.28 ^c ±0.2
	Range	44.7-52.8	0.1-1.3	1.9-4.5	1.5-3.4	11.3-16.7	0.2-1.0	2.2-6.0	1.2-2.2	8.6-13.6
HB	Mean	6.23 ^a ±0.2	0.48 ^b ±0.1	1.64 ^a ±0.1	1.55 ^b ±0.4	5.23 ^a ±0.2	0.53 ^b ±0.1	2.69 ^a ±0.1	0.88 ^a ±0.1	5.35 ^a ±0.14
	Range	4.2-7.5	0.3-0.6	1.4-2.2	0.8-1.0	3.3-6.4	0.3-0.9	2.0-4.1	0.7-1.2	3-6.2
TA	Mean	11.67 ^b ±0.3	0.58 ^c ±0.1	1.68 ^a ±0.2	1.14 ^b ±0.1	10.62 ^b ±0.6	0.69 ^c ±0.1	4.35 ^b ±0.2	0.91 ^b ±0.1	6.98 ^b ±0.3
	Range	9.6-13.4	0.3-0.9	1.0-2.6	0.4-1.9	5.0-13.0	0.4-0.9	3.1-5.8	0.3-1.4	5.0-8.9
TQ	Mean	25.40 ^c ±0.8	0.70 ^d ±0.1	1.67 ^a ±0.1	1.06 ^a ±0.1	11.3 ^b ±0.4	0.63 ^c ±0.1	4.14 ^b ±0.1	1.03 ^b ±0.1	7.30 ^b ±0.2
	Range	13.2-20.0	0.5-0.9	1.4-1.9	0.8-1.3	9.1-12.2	0.4-0.9	3.8-4.4	0.7-1.4	6.5-8.0

262 *Means within column with different Superscript are significantly different ($P \leq 0.05$) from each other

263 Footnotes

264 AA is *Agama agama* (Agama Lizard), HB is *Hemidactylus brookii* (Wall gecko), TA is *Trachylepis affinis* (Skinks) and TQ is *Trachylepis quinquetaeniata* (Skinks)

265 BW is the Body weight, HH is the Height of the head, HL is the Head length, HW is the Head width, TAL is the Tail length, TAW is the Tail width, TRL is the Trunk length,

266 MO is the Mouth opening and SVL is the Snout-vent length

267

268

269

270 The Trunk length (TRL) average value of three specimens (*A. agama*, *T. affinis*, and *T.*
271 *quinquetaeniata*) have similar values (4.05, 4.35, and 4.14 cm) without significant difference
272 ($p > 0.05$), but differed from *H. brookii* (2.69 cm) which have the least value. The Mouth
273 opening (MO) average value of *A. agama* was the highest (1.51 cm) and *H. brookii* (0.88 cm)
274 have the least MO value. There was no significant difference ($p > 0.05$) in MO value of *T.*
275 *affinis* and *T. quinquetaeniata*. Snout-vent length (SVL) of *A. agama* was the highest (10.28
276 cm) followed by *T. quinquetaeniata* (7.30 cm). The SVL value of *H. brookii* was the least
277 (5.35 cm) and there was no significant difference ($p > 0.05$) in SVL of *T. affinis* and *T.*
278 *quinquetaeniata*.


279

280 Figure 3 showed the Principal Component Analysis (PCA) scatter plot of the morphometric
281 relationship among the lizards (*A. agama*, *H. brookii*, *T. affinis* and *T. quinquetaeniata*)
282 captured in various location in the University. From the figure, the morphometric parameters
283 measured on *A. agama* were different from other specimens. Although, it can be said that
284 there was a slight similarity between *A. agama* and *T. affinis*. The remaining specimens (*H.*
285 *brookii*, *T. affinis* and *T. quinquetaeniata*) have morphometric similarities with *T. affinis*
286 having morphometric parameters common to the remaining two specimens (*H. brookii* and *T.*
287 *quinquetaeniata*).

288

289

290



292 **Figure 3:** Principal Component Analysis (PCA) Scatter Plot showing Morphometric
 293 relationship in lizard species captured in O.A.U.

294 Keys

295 Red colour is Agama Lizard (*Agama agama*)


296 Blue colour is Wall gecko (*Hemidactylus brookii*)

297 Pink colour is Skinks (*Trachylepis affinis*)

298 Green colour is Skinks (*Trachylepis quinquetaeniata*)

299

300 The Principal Component Analysis (PCA) loading showed some of the morphometric
 301 parameters that were different in the specimens (Figure 4). Tail length, Tail width and Trunk
 302 length were differed from one specimen to another with cut off of 0.4.



303

304 **Figure 4:** PCA loading Plot showing the significant morphometric parameters within the four
 305 species of lizard with a cut off of 0.4.

306

307 4.1 Discussions

308 The increase in the population of *A. agama* compared to other specimens (*H. brookii*, *T.*
309 *affinis* and *T. quinquetaeniata*) in all the location in the University can be attributed to their
310 ability to feed on different varieties of insects and plant materials (21). *A. agama* is regarded
311 as frugivorous and seed dispersion as well as insectivorous animals. The abundance of *H.*
312 *brookii* residence (University Hostel, University staff quarters and University Library) may
313 be attributed to easy accessibility to food since they are commonly found in the house.
314 Tkaczenko et al. (22) reported that Hemidactylus species are highly adapted to living close to
315 people since they feed on insects that are attracted to artificial light sources which are
316 commonly found where people live. The report of Tkaczenko et al. (22) was supported by
317 Weterings (23). He reported that Hemidactylus spp also forages in garbage bins and on tables
318 where they feed on leftover foods.

319 In case of Skinks (*T. affinis* and *T. quinquetaeniata*), the decrease in their population in most
320 of the locations (University Hostel, University staff quarter and University Library) may be
321 attributed to the absence of grasses and shrubs. The presence of grasses and shrubs in the
322 University market and University parks and garden may be responsible for the increases in
323 the amount of Skinks specimens collected in both locations. Kadry et al. (24) report on five-
324 lined Skinks showed that Skinks are ground-dwelling and associated with grasses and shrubs
325 for protection against predators. They also reported that grasses and shrubs provided suitable
326 temperature for Skins behaviour.

327 The ratio of the female specimen in this study was higher than that of the male specimen in
328 all the locations in the University. The ratio of the female specimen in this study may be due
329 to the search for food and early sexual maturity. Female lizards (e.g *A. agama*) reach sexual
330 maturity between fourteen and eighteen months, while it takes two years for male *A. agama*
331 to sexual maturity (25). Also, the reduction amount of male specimen in this study may be
332 attributed to competition and territorial behaviour which may lead to death or accessibility to

333 the predator (26). This result was similar to our previous findings on the Black rat (*Rattus*
334 *rattus*) in Obafemi Awolowo University (13).

335 Variation in the measured morphometric parameters of the specimens may be due to different
336 factors. Some of the factor to be considered are sex, age, habitats, an abundance of food and
337 predator (27). The report of phenotypic analysis of sexual size dimorphism of lizards showed
338 that female small lizard's species tend to be smaller than males, while males of large species
339 have higher body weight than female (28). In this study, the bodyweight of *A. agama* male
340 was higher than the female, while the bodyweight of female Skinks (*T. affinis* and *T.*
341 *quinquetaeniata*) was higher than male. Maturity in respect to the age of the species may also
342 be responsible for variation in measured morphometric parameters. Although the age of the
343 specimens was not considered in this study, the report of Jena et al. (29) showed that young
344 lizards morphometric parameters are lower when compare with the older ones.

345 The habitats of lizards may be important when considering the differences in their
346 morphometric parameters. Lizards being commonly found in a warm climate but leave in
347 cold climate may develop some adaptive measure to survive the cold condition (30).
348 Kaliontzopontou et al. (31) reported that morphology and habitat is a central theme in
349 evolutionary biology because it reflects the way organisms adapt to their environment. The
350 reports of Kohlsdorf et al. (32) showed that there is always a relationship between
351 morphology and habitat which affect the body size and other parts of lizards.

352 The abundance of food is a major factor of life that determines how an individual's species
353 allocate energy among competing demands and maturation (33). Food abundance in lizards
354 can have direct long-lasting effects on external morphometric of lizards (34). The abundance
355 of food may also attract the predator such as snakes to the habitat where lizards dwell.
356 Lizards serve as food for snakes and other predators like Hawks (35).

357 The difference in the morphometric parameters of *A. agama* may be responsible for the
358 differences seen in the scatter plot (Figure 3). *A. agama* can be considered bigger lizard

359 when compared with Wall gecko (*H. brookii*) and Skinks (*T. affinis* and *T. quinquetaeniata*)
360 used for this study. It was shown from the morphometric parameters that *A. agama* have
361 higher values in all the parameters measured than the rest species. Among the rest species (*H.*
362 *brookii*, *T. affinis* and *T. quinquetaeniata*), there was interaction in all the parameters
363 measured and the interaction may be attributed to similarity in average values of the
364 measured parameters. Although the Principal Component Analysis (PCA) loading plot
365 (Figure 4) showed that Sex, Tail length (TL), Tail width (TW) and Trunk length of all the
366 lizards' species differs. The difference in the above mentioned parameters may be due to the
367 difference in species of lizard studied. The tail of male *A. agama* was reported used in
368 fighting when defending their territory (36) while *H. brookii* do not use tail in the fighting.
369 This may be responsible for an increase in TL, TW, and TRL in *A. agama* than the rest
370 species.

371 The abundance of *A. agama* and *H. brookii* in University Hostels, University staff quarters
372 and Library call for great concern since they are parasite carrier which can cause a great
373 health concern when defecated or die in the food or urinate inside the food. It is important for
374 the University authority to look for the best way to reduce the percentage of these lizards in
375 the University.

376

377 **Conclusion**

378 *A. agama* and *H. brookii* are the most abundant species of lizards in all the collection
379 locations and the measured morphometric parameters show a significant difference in all the
380 lizards measured. It is important that University authority looks into ways to reduce the
381 percentage of these lizards in University residence in order to prevent diseases.

382

383

384 **Conflict of interest**

385 Authors have declared that no competing interests exist.

386

387 **References**

- 388 1. Prokop, P., & Fedor, P. (2013). The effects of parasites on human behaviour: An
389 evolutionary perspective. *Problems of Psychology in the 21st Century*, 5(5), 46-64.
- 390 2. Bridges, V., Koprak, C., & Johnson, R. (2001). The reptile and amphibian communities
391 in the United States. *USDA: APHIS: VS Centers for Epidemiology and Animal Health*,
392 *Fort Collins*.
- 393 3. Hoffman, L. C., & Cawthorn, D. M. (2012). What is the role and contribution of meat
394 from wildlife in providing high quality protein for consumption?. *Animal Frontiers*, 2(4),
395 40-53.
- 396 4. Leaché, A. D., Rödel, M. O., Linkem, C. W., Diaz, R. E., Hillers, A., & Fujita, M. K.
397 (2006). Biodiversity in a forest island: reptiles and amphibians of the West African Togo
398 Hills. *Amphibian and Reptile Conservation*, 4(1), 22-45.
- 399 5. Sowemimo, O. A., & Oluwafemi, T. A. (2017). A Survey of Helminth Parasites of the
400 Lizard, *Agama agama* in Ile-Ife and Ibadan Southwest Nigeria. *J. Bacteriol. Parasitol*, 8,
401 1-6.
- 402 6. Nash, H. (2005). Pentastomes: Respiratory Parasites of Reptiles. *Veterinary Services*
403 *Department, Drs. Foster & Smith, city, country*.
- 404 7. Rocha, C. F. D., & Anjos, L. A. (2007). Feeding ecology of a nocturnal invasive alien
405 lizard species, *Hemidactylus mabouia* Moreau de Jonnès, 1818 (Gekkonidae), living in
406 an outcrop rocky area in southeastern Brazil. *Brazilian Journal of Biology*, 67(3), 485-
407 491.
- 408 8. Oluwafemi, T., Sowemimo, O., & Bamidele, A. (2017). Parasitic infections of two
409 species of lizard (*Hemidactylus frenatus* and *Mabuya quinquetaeniata*) from two cities in
410 Southwest Nigeria. *Annual Research & Review in Biology*, 18(2), 1-13.
- 411 9. Castiglia, R., Annesi, F., Bezerra, A. M., Garcia, A., & Flores-Villela, O. (2010).
412 Cytotaxonomy and DNA taxonomy of lizards (Squamata, Sauria) from a tropical dry

- 413 forest in the Chamela-Cuixmala Biosphere Reserve on the coast of Jalisco,
414 Mexico. *Zootaxa*, 2508(1), 1-29.
- 415 10. Adeoye, G. O., & Ogunbanwo, O. O. (2007). Helminth parasites of the African lizard
416 *Agama agama* (Squamata: Agamidae), in Lagos, Nigeria. *Revista de biología*
417 *tropical*, 55(2), 417-425.
- 418 11. Nwadike, C. C., & Ilozumba, P. C. O. (2015). Prevalence of helminth parasites in
419 rainbow lizard, *Agama agama* L.(Squamata: Agamidae) in Nsugbe, Anambra State,
420 Nigeria. *Zoologist (The)*, 13, 22-27.
- 421 12. Omonona, A. O., Adedokun, O. A., & Adekoya-Gafaar, S. A. (2011). Parasitological
422 studies on *Agama agama* lizard (Squamata: Agamidae) in Ibadan. *Adv. Environ. Biol*, 5(5), 803-807.
- 423 13. Bamidele, A. O., & Kowobari, E. D. (2019). Morphology and Bait Preference of Black
424 Rat (*Rattus rattus*) in the University Community in Nigeria: Obafemi Awolowo
425 University as a Case Study. *Asian Journal of Research in Zoology*, 1-12.
- 426 14. Harris, V.A. (1964): The life of the rainbow lizard. Hutchinson, London. pp. 1-174.
- 427 15. Bauer, A. M., Pauwels, O. S., & Chanhom, L. (2002). A new species of cave-dwelling
428 *Cyrtodactylus* (Squamata: Gekkonidae) from Thailand. *Tropical Natural History*, 2(2),
429 19-29.
- 430 16. Bauer, A. M. (2003). On the identity of *Lacerta punctata* Linnaeus 1758, the type species
431 of the genus *Euprepis* Wagler 1830, and the generic assignment of Afro-Malagasy
432 skinks. *African Journal of Herpetology*, 52(1), 1-7.
- 433 17. Spawls, S., Howell, K., Drewes, R., & Ashe, J. (2004). *A field guide to the reptiles of*
434 *East Africa: Kenya, Tanzania, Uganda, Rwanda and Burundi*. London: A & C Black.
- 435 18. Kaliontzopoulou, A., Carretero, M. A., & Llorente, G. A. (2007). Multivariate and
436 geometric morphometrics in the analysis of sexual dimorphism variation in *Podarcis*
437 lizards. *Journal of Morphology*, 268(2), 152-165.

- 438 19. Uyeda, L. T., Iskandar, E., Kyes, R. C., & Wirsing, A. J. (2015). Encounter rates,
439 agonistic interactions, and social hierarchy among garbage-feeding water monitor lizards
440 (*Varanus salvator bivittatus*) on Tinjil Island, Indonesia. *Herpetological Conservation
441 and Biology*, 10(2), 753-764.
- 442 20. SAS Institute, (1997). System analysis software, version 9.1. SAS Institute Inc, Carry
443 NC
- 444 21. Herrera, C. M. (2002). Seed dispersal by vertebrates. *Plant–animal interactions: an
445 evolutionary approach*, 185-208.
- 446 22. Tkaczenko, G. K., Fischer, A. C., & Weterings, R. (2014). Prey preference of the
447 Common House Geckos *Hemidactylus frenatus* and *Hemidactylus
448 platyurus*. *Herpetology Notes*, 7, 483-488.
- 449 23. Weterings, R. (2017). Observations of an adaptive feeding strategy in flat-tailed house
450 geckos (*Hemidactylus platyurus*) living in buildings. *Herpetology Notes*, 10, 133-135
- 451 24. Kadry, M. A., Mohamed, H. R., & Hosney, M. (2017). Ecological and biological studies
452 on five-lined skink, *Trachylepis* (= *Mabuya*) *quinquetaeniata* inhabiting two different
453 habitats in Egypt. *Cell Mol Biol (Noisy le Grand)*, 63(11).
- 454 25. Adams, E. S. (2001). Approaches to the study of territory size and shape. *Annual Review
455 of Ecology and Systematics*, 32(1), 277-303.
- 456 26. Miaud, C., Andreone, F., Ribéron, A., De Michelis, S., Clima, V., Castanet, J., ... &
457 Guyétant, R. (2001). Variations in age, size at maturity and gestation duration among
458 two neighbouring populations of the alpine salamander (*Salamandra lanzai*). *Journal of
459 Zoology*, 254(2), 251-260.
- 460 27. Kidawa, D., & Kowalczyk, R. (2011). The effects of sex, age, season and habitat on diet
461 of the red fox *Vulpes vulpes* in northeastern Poland. *Acta theriologica*, 56(3), 209-218.

- 462 28. Colwell, R. K. (2000). Rensch's rule crosses the line: convergent allometry of sexual size
463 dimorphism in hummingbirds and flower mites. *The American Naturalist*, 156(5), 495-
464 510.
- 465 29. Jena, B. S., Nayak, S. B., & Patnaik, B. K. (2002). Age-related effect of aluminium on
466 the catalase activities of the brains of two species of poikilothermic
467 vertebrates. *Gerontology*, 48(1), 34-38.
- 468 30. Harding, J. H., & Mifsud, D. A. (2017). *Amphibians and reptiles of the Great Lakes*
469 *region*. University of Michigan Press.
- 470 31. Kaliontzopoulou, A., Carretero, M. A., & Llorente, G. A. (2010). Intraspecific
471 ecomorphological variation: linear and geometric morphometrics reveal habitat-related
472 patterns within *Podarcis bocagei* wall lizards. *Journal of Evolutionary Biology*, 23(6),
473 1234-1244.
- 474 32. Kohlsdorf, T., Garland Jr, T., & Navas, C. A. (2001). Limb and tail lengths in relation to
475 substrate usage in *Tropidurus* lizards. *Journal of Morphology*, 248(2), 151-164.
- 476 33. Mugabo, M., Marquis, O., Perret, S., & Le Galliard, J. F. (2010). Immediate and delayed
477 life history effects caused by food deprivation early in life in a short-lived lizard. *Journal*
478 *of evolutionary biology*, 23(9), 1886-1898.
- 479 34. Taborsky, B. (2005). The influence of juvenile and adult environments on life-history
480 trajectories. *Proceedings of the Royal Society B: Biological Sciences*, 273(1587), 741-
481 750.
- 482 35. Cortés-Gomez, A. M., Ruiz-Agudelo, C. A., Valencia-Aguilar, A., & Ladle, R. J. (2015).
483 Ecological functions of neotropical amphibians and reptiles: a review. *Universitas*
484 *Scientiarum*, 20(2), 229-245.
- 485 36. Cooper Jr, W. E. (2003). Social behavior and antipredatory defense in lizards. *Lizard*
486 *social behavior*. Johns Hopkins University Press, Baltimore, 107-141.
- 487