

ANNUAL EFFECTIVE DOSE AND LIFETIME CANCER RISKS DUE TO NATURAL RADIOACTIVITY IN GROUND WATER OF TAI RIVERS STATE, NIGERIA

Abstract

The presence of radionuclide in water poses a number of health hazards, especially when the radionuclide is deposited in the human body through drinking water. The aim of this study was to evaluate natural radioactivity and its associated health risk in hand dug well water of Tai Local Government Area of Rivers State, Nigeria by means of gamma spectroscopy techniques and radiation models. The well water was collected from five selected coastal communities of Tai and chemically treated by adding nitric acid and then pre-concentrated further by evaporating to certain levels. The residue were transferred to small cylindrical containers were kept sealed in order to ensure secular equilibrium between ^{238}U , ^{232}Th and their progenies for 28 days and counted with sodium iodide activated with thallium detector. The specific activity concentration of ^{40}K , ^{226}Ra and ^{228}Ra (^{232}Th) obtained were used to estimate the annual effective doses for different age categories by taking into consideration the ingested dose conversion factors as well as their yearly average water consumption. The annual effective dose due to ingestion of the sampled water were above the recommended values by WHO, IAEA and UNSCEAR. The paper presents the overview of the techniques used and the summary of the findings. The result of this study gives the radiological baseline data for effective monitoring of the study area.

Keywords: Tai, Radionuclide, Spectroscopy, lifetime cancer risk and hereditary effects.

1. Introduction

There are numerous liquids found in nature, water being one of them has a vast usage to animals, plants, humans and humans' activities, operations, productions and other services. According to Stabans and Kinsars [1] water is a global solvent on our planet, whose main sources include rivers, springs, wells, boreholes, seas and other fresh water bodies. Environmental radiation is either natural or artificial, however results to production of radionuclide, energies, particles and waves. Saghatchi [2] reported that radioactive materials exist everywhere in human environment and the largest portion of human exposure comes from natural sources. Stabans E.I and Kinsars [1] noted that, many radioactive compounds or radionuclides are released into the environment and hence into drinking water supply due to human activities. He added that radionuclide can also enter the food chain if the contaminated water is used for drinking or irrigation purpose.

Ajayi and Awolabi [3], declared that the presence of radionuclide in water poses a number of health hazards, especially when the radionuclide is deposited in the human body through drinking water.

Ononugbo, Avwiri and Egbeya [4] stated that, many people are increasingly concerned about a variety of contaminants in drinking water, especially those which affect human health. Ononugbo *et al.*, [4] stated that the presence of radionuclide in drinking water can be attributed to a variety of sources, including the improper disposal of household waste products, ground storage tanks, discharge from oil and gas industry and increased use of pesticides.

Radionuclide in drinking water causes human internal exposure, caused by the decay of radionuclide taken into the body by ingestion and inhalation indirectly when they are incorporated as part of the human food chain [4, 5]. Several naturally occurring alpha and beta emitting radionuclide such as ^{238}U , ^{226}Ra , ^{216}pb , ^{228}Th and others are frequently dissolved in ground water supplies (well water inclusive) and their concentrations vary over an extremely wide range, mainly depending upon the amount of radioelement in bedrock and soil with which the water comes in contact [6].

Many part of the world get their water from ground water or deep wells and the quality of water we consume at every in time determines our state of health. Most materials and all environments are exposed to ionizing radiation and few of the materials exposed to this ionizing radiation are used as building materials for our accommodations which during processing add up to the amounts of radioactive elements in the environment [7]. Completely removing radionuclides from drinking water especially well water before consumption will probably reduce the cases of terminal diseases such as cancers, benign tumors and even cataracts by few percentage but mostly radon enters the air in a home through exposure to soil and rocks, uranium are eliminated via urine and radium is stored in the bone while others are stored in the blood [8].

In Tai Local Government Area, oil exploration, drilling and local refining of crude oil has resulted to oil leakage on river waters, farm lands and seas. Moreover, oil drilling companies including shell, Elf and Chevron do flare gases into the open air, thereby increasing soot concentration in the atmosphere, acid rain and more production of radionuclide like uranium, radon gas, thorium, and potassium. In most industrial areas and host communities in Rivers State (Tai Inclusive), availability of clean and safe drinking water has been a critical issue to tackle, due to oil reservoirs, crude oil co-exist with underground water called formation water [4]. Produced water contains some level of naturally occurring radionuclide [9]. The aim of this study is to evaluate activity concentration of radionuclide in hand dug well water from some selected communities in Tai Local Government Area in order to quantify the health implication of ingesting such water.

2. Materials and Methods

2.1 Study Area

The study was carried out in five selected coastal Communities Tai Local Government Area of Rivers State, Nigeria these communities are Nonwa community, Borobara community, Kira community, Korokoro community and Kpите community. It covers an area of 159 km^2 and lies between latitude $4^{\circ}43'0''\text{N}$ and longitude $7^{\circ}18'0''\text{E}$. the major communities within this region (seat of the monarch of Tail) include: Nonwa, Sime, Kira, Borobara, Kpите, Korokoro, Koroma, Bunu, Ban-ogoi, Horo, Ueken. Generally, Tail Local Government Area has two board sections. The Tua-Tua Kingdom and the

Nonwa kingdom and is situated in the South East Area of River State. It is bounded by Eleme L.G.A, Gokana L.G.A, Khana L.G.A and Oyigbo L.G.A.




Fig. 1: Map of the Study Area

Sampling was conducted from Nonwa, Borobara, Kira, Kpite and Korokoro community respectively. A total of 22 water samples were collected, and four each from each community and 2 sample was collected from outside the communities which serve as control. Plastic bottles of 1 liter each were used to collect water samples. The containers were raised twice with the water sample in order to minimize contamination from the original content of the container[10]. The water sample were then acidified with 2 drops of diluted HCL in each of the water samples to minimize the precipitation of the radionuclide present in the water sample and also prevent the absorption of the radionuclide on the walls of the containers. The water samples were tightly covered and it was stored for four weeks to reach secular equilibrium.

2.3 Gamma-ray Activity Measurement

Activity measurements of radionuclides in surface water were performed at the University of Agriculture, Abeokuta with a gamma-ray spectrometry system with a thallium activated 3" x 3" sodium iodide on a Na(Tl) detector connected to ORTEC 456 amplifier. The detector in a 100mm thick lead shield, was connected to a computer program called SAMPO 90 window that matched gamma energies to a library of possible isotopes. Since the accuracy of the quantitative measurement is dependent on the calibration of the spectrometry system and adequate energy; background measurement and efficient calibration of the system was made using Cs-137 and Co-60 standard sources from IAEA, Vienna [11].

The analysis was performed using a Canberra S 100 computer analyzer. Standard of natural origin were prepared in the same manner as the samples, these standards are uranyl nitrate ($\text{UO}_2(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$) 502.18 mol/g, potassium chloride (KCl) 74.55 mol/g and thorium nitrate ($\text{Th}(\text{NO}_3)_4 \cdot 5\text{H}_2\text{O}$) 570.13 mol/g. One gram of each of the standard was taken and dissolved into a 200 ml distilled water to form a standard solution. It is subtle that 1g of uranyl nitrate contains 0.474g of uranium which has activity of 0.0294 Bq l^{-1} , also 1g of potassium chloride contains 0.534g of potassium which has activity of 0.706 Bq l^{-1} and 1g of thorium nitrate contains 0.859 g of thorium with activity of 0.0175 Bq l^{-1} .

Spectrum were accumulated for background for a period of 2900s at volts to produce strong peaks at gamma emitting energies of 1460 Kev for ^{40}K ; 609 Kev of ^{232}Bi and 911 Kev of ^{228}Ac , which were used to estimate the concentration of ^{238}U and ^{232}Th respectively. The detector was calibrated with cesium-137 and cobalt-60 sources and the energy resolution is 39.5 and 22.2% respectively. The configuration and geometry was maintained throughout the analysis. The individual radionuclide concentration was calculated using relative method as in the Equation below:

$$\frac{\text{Activity of } U1}{\text{Activity of } S1} = \frac{\Sigma U1 - \Sigma b}{\Sigma S1 - \Sigma b} \quad 1$$

Where $U1$ = the unknown sample activity concentration in the unit of Bq l^{-1} , $S1$ = activity of the standard source, $\Sigma U1$ = sum under the peak of $U1$ in cps $\Sigma S1$ = the sum under the peak $S1$ in cps.

3. Radiological Risk Assessment

The annual effective dose from ingestion of radionuclide in water samples was estimated on the basis of the mean activity concentration of the radionuclides. This was done for different age brackets. In this work the intake rates and dose conversion factors for the radionuclides based on the International commission on radiological protection [12] publication are used: 0.5 L/day and 1.0 l/day for infants (age $\leq 1\text{yr}$) and Children (age 1-12 years) respectively. And 2 L/day for teenagers ($\leq 17\text{yr}$) and adult ($> 17 \text{ yrs}$) as presented in Table 1.

The annual effective dose from ingestion of ground water was computed by the following equation [13].

$$H_{\text{ing}}(w) = \sum DCF_{\text{ing}}(i) \times A_{\text{spi}} \times I \quad (2)$$

DCFing is dose conversion coefficient of a particular radionuclide ith in Sv/Bq for a particular age category, As_{pi} is the specific activity concentrations of radionuclide ith in the water samples in Bq/l and I is radionuclide intake in litres per year for each age category.

In addition to the estimated annual effective dose, the cancer and hereditary risk due to low dose without any threshold dose known as stochastic effect were estimated using the ICRP cancer risk model [14]. Radiation risk to population result from exposure to low dose radiation are normally known as chronic risk of somatic or hereditary damage of human tissues, thus much emphasis is always placed on the reduction of these radiological risks to natural radiation.

The nominal lifetime risk coefficient of fatal cancer recommended in the 2007 recommendations of the ICRP for members of the public is $5.5 \times 10^{-2} \text{ Sv}^{-1}$. For hereditary effects, the detriment adjusted nominal risk coefficient for the whole population as stated in ICRP [14] for stochastic effects after exposure at low dose rates is estimated at $0.2 \times 10^{-2} \text{ Sv}^{-1}$.

The risk to population was then estimated using the recommended risk coefficient in ICRP report and assumed 70 years lifetime of continuous exposure of population to low level radiation. According to the ICRP methodology:

$$\text{Cancer Risk} = \text{Total annual Effective Dose (Sv)} \times \text{cancer risk factor} \quad (3)$$

$$\text{Hereditary Effects} = \text{Total annual Effective Dose (Sv)} \times \text{hereditary effect factor} \quad (4)$$

Table 1: Committed Effective Dose Conversion Factor (Sv/Bq) for members of the Public (ICRP, 2012)

S/N	Radioisotope	Infant $\leq 1\text{yr}$	Children (1-12yr)	Teenage (12-17)	Adult $\geq 17\text{yr}$
1	^{226}Ra	4.7 E-06	6.2 E-07	1.5 E-06	2.8 E-07
2	^{232}Th	3.0E-05	3.4 E-06	5.3 E-06	6.2 E-07
3	^{40}K	6.2 E-08	2.1 E-08	7.6 E-09	6.2 E-09
	H ₂ O Intake	0.5 L	1.0 L	1.5 L	2.0 L

4. Results and Discussion

The specific activity concentrations of ^{40}K , ^{226}Ra , and ^{228}Ra (^{232}Th) identified in hand dug well water samples from the selected coastal communities in Tai Local Government area of Rivers State as well as their radium equivalent (Raeq) are presented in Table2. From Table 2, the activity concentration of ^{40}K varies distribution of the radionuclide identified from 5.68 ± 4.59 to $88.99 \pm 3.01 \text{ Bq l}^{-1}$ with an average value of 25.90 Bq l^{-1} . The specific activity concentration of ^{226}Ra ranged from 4.51 ± 2.9 to $44.73 \pm 3.64 \text{ Bq l}^{-1}$ with average value of 19.21 Bq l^{-1} . The activity concentration of ^{228}Ra (^{232}Th) varied between BDL to $221.81 \pm 2.61 \text{ Bq l}^{-1}$ with an average value of 18.50 Bq l^{-1} .

This variation in activity concentration of ^{40}K , ^{226}Ra , and ^{228}Ra (^{232}Th) observed in the samples indicate that the origins of this water are not the same and they come from different depths and pass

through different geological layers. Also this irregular distribution of the radionuclides identified could be due to industrial activities in these communities and their content in solid aquifers in the area of study. These variations in activity concentration of ^{40}K , ^{226}Ra , and ^{228}Ra (^{232}Th) strongly depend on the physical and chemical properties of each water sample.

Activity concentration for ^{40}K , ^{226}Ra , and ^{232}Th were high in all the samples except in Borobara 2 where the activity concentration of ^{232}Th were below the detectible limit. This could be due to agrochemical which include nutrients of plants, fertilizers, and discharge of oil and gas products into the environments. The increase in the activity concentration across all communities could be a result of effluent production from the oil and gas drilling companies which might have concentrated potassium and radium at those locations. The highest activity concentration of ^{40}K ($88.99\pm 3.01 \text{ Bq l}^{-1}$) was recorded at Kira 2. This could be due to massive use of NPK fertilizer in the area to improve the crop yield which might get into the water bodies and ground water. Also the highest activity concentration of ^{228}Ra (^{232}Th) was recorded in Borobara 1 ($24.81\pm 2.61 \text{ Bq l}^{-1}$), ^{40}K , ^{232}Th and ^{226}Ra was detected in all the well water and this could be because of the frequent and constant discharge of oil and gas into ground water and other maritime activities which has concentrated the activity concentration of these radionuclides.

The results obtained shows that the activity concentration of ^{226}Ra , ^{232}Th and ^{40}K in sampled hand dug well water were higher than safe recommended value set by the environment protection agency and World Health Organization as shown in Figures 2 to 4. The range of thorium (^{232}Th) in natural water is set at $1.0\text{-}10.0 \text{ mBq l}^{-1}$ and uranium (^{226}Ra) range from $1.0\text{-}10 \text{ Bq l}^{-1}$ but the range for ^{40}K was not specified by Environmental Protection Agency and WHO, [15] recommended the maximum allowable committed effective dose equivalent of 4 mrem/y for annual intake for adult.

The average activity concentration of ^{40}K (25.90 Bq l^{-1}) in the present study is comparatively higher than the average value obtained in Cameron (0.107 Bq l^{-1}) by Ndontchueng et al., [13]. It is also higher than the average values obtained in Italy (Milano) (0.05 Bq l^{-1}) by Roscuni [16].

Table 2: Specific Activity Concentrations of Radionuclide in Various Sampling Locations and its Radium Equivalent.

S/N	Sampling Code	Activity Concentration in Bq l^{-1}			Ra_{eq} (Bq l^{-1})
		^{40}K	^{226}Ra	^{228}Ra (^{232}Th)	
1	KPITE1	16.42 ± 3.13	37.36 ± 1.83	23.43 ± 1.74	72.13
2	KPITE2	14.36 ± 2.56	9.50 ± 2.32	16.11 ± 3.18	33.64
3	KPITE3	5.68 ± 2.54	36.54 ± 1.45	23.84 ± 1.74	71.07
4	KPITE4	13.58 ± 4.52	11.96 ± 4.05	24.25 ± 2.46	47.68
5	KIRA 1	12.00 ± 2.06	4.59 ± 2.54	22.62 ± 2.67	37.86

6	KIRA 2	88.99±3.01	7.78±4.85	25.06±3.27	50.47
7	KIRA 3	10.79±4.72	10.65±3.89	27.77±3.58	51.19
8	KIRA 4	11.99±4.60	7.50±3.00	20.23±2.92	37.35
9	NONWA 1	26.99±4.50	4.59±3.12	27.73±1.92	46.32
10	NONWA 2	5.68±4.59	19.17±2.34	23.98±2.67	53.90
11	NONWA 3	14.58±4.21	31.63±2.67	27.37±1.55	71.89
12	NONWA 4	43.57±3.03	28.35±2.74	22.89±3.36	64.44
13	BOROBARA 1	30.15±4.79	11.31±2.24	221.81±2.61	45.35
14	BOROBARA 2	45.93±4.81	35.72±2.46	BDL	39.26
15	BOROBARA 3	32.52±4.41	28.27±2.74	23.98±3.18	65.07
16	BOROBARA 4	13.31±4.52	28.35±2.67	25.87±2.24	66.37
17	KOROKORO 1	38.73±5.25	44.73±3.64	23.30±2.61	50.14
18	KOROKORO 2	40.65±3.15	4.51±2.9	26.82±3.31	45.99
19	KOROKORO 3	34.72±3.67	15.77±2.67	27.91±4.40	58.35
20	KOROKORO 4	27.25±1.22	8.68±3.12	24.61±2.30	45.97
21	CONTROL 1	7.04±4.07	6.14±1.83	5.33±1.55	83.88
22	CONTROL 2	4.89±4.71	1.34±3.42	6.28±1.92	59.61
	Mean	25.90	19.21	18.50	54.43
	UNSCAR, 2000	10.0	10.0	1.0	-




Fig.2: Comparison of Activity concentration of ^{40}K with UNSCEAR, (2000) Recommendation




Fig.3: Comparison of Activity concentration of ^{226}Ra with UNSCEAR, (2000) Recommendation




Fig.4: Comparison of Activity Concentration of ²³²Th with UNSCEAR Recommendation

In the same way, the average concentration of ²²⁸Ra (²³²Th) 18.50 Bq⁻¹ is comparatively higher than the value obtained in Cameroon (0.036 Bq⁻¹) by Nduentgen *et al.*, [13]. The values obtained in this work were thus compared favorably with the reported average values published by other authors. The activity concentration of ⁴⁰K, ²²⁶Ra and ²²⁸Ra (²³²Th) obtained in this study were also compared with the guideline activity concentration values of the radionuclide in drinking water recommended by the World Health (WHO) and other data obtained from IAEA. This comparison showed that our results were found to be higher than the recommended safe value.

The annual effective dose due to ingestion of the monitored ground water was estimated for different age groups including infants, children, teenage and adults considering only the ingestion of ²²⁶Ra and ²²⁸Ra (²³²Th) as shown in Table 3. Potassium (⁴⁰K) values were not considered during the calculation of the radiation dose because the absorption of the essential potassium element is under homeostatic control and takes place mainly from ingested food. Thus, the potassium contribution to the dose from ingestion in water, with its relatively low dose conversion factor will be much less than that of many other radionuclides.

The estimated effective dose for different age groups were ranged from 0.00007 to 0.318 Svy⁻¹ for infants, between 0.00005 and 0.078 Svy⁻¹ for children, varied from 0.00004 to 0.238 Svy⁻¹ for teenagers and varied from 0.00003 to 0.051 Svy⁻¹ for adults with average values of 0.115, 0.027, 0.071 and 0.013 Svy⁻¹ respectively. It can be seen that radiation dose received by infants are relatively higher than that received for children, teenagers and adults. This result is an agreement with the work done in Cameroon by Ndontchueng *et al.*, [13]. Comparing with the WHO, IAEA and UNSCEAR [15, 11, 17] recommended reference levels of the effective dose for infants, children, teenagers and adults corresponding to one year consumption of drinking water are 0.26, 0.2 and 0.1 mSvy⁻¹, respectively. The doses obtained in this study are higher than the recommended reference level and from radiation protection point of view, life-long consumption of these investigated hand dug well waters may cause significant radiological health risk.

Table 3: Annual Effective doses for different Ages and Estimated Cancer risks and Hereditary Effects on Adult member of the Public

S/N	location	Total Effective Dose (Svy ⁻¹)				Cancer Risk to Adult		Hereditary Effect in Adult	
		E _T Infant	E _T Child	E _T Teenage	E _T Adult	FCR x10 ⁻³	LFCR x10 ⁻³	SHE x10 ⁻⁵	ELHE x10 ⁻³
1	KPITE1	0.111	0.026	0.065	0.011	0.62	43.48	2.26	1.58
2	KPITE2	0.318	0.078	0.238	0.051	2.83	198.11	10.29	7.20
3	KPITE3	0.142	0.033	0.088	0.016	0.90	62.86	3.27	2.29
4	KPITE4	0.121	0.028	0.071	0.013	0.71	49.61	2.58	1.80
5	KIRA 1	0.185	0.043	0.109	0.019	1.06	73.97	3.84	2.69
6	KIRA 2	0.115	0.027	0.066	0.011	0.62	43.29	2.25	1.57
7	KIRA 3	0.088	0.020	0.046	0.007	0.40	27.94	1.45	1.02
8	KIRA 4	0.083	0.020	0.056	0.011	0.63	44.04	2.29	1.60
9	NONWA 1	0.104	0.025	0.068	0.013	0.73	51.17	2.66	1.86
10	NONWA 2	0.159	0.036	0.089	0.015	0.83	58.04	3.02	2.11
11	NONWA 3	0.138	0.032	0.076	0.013	0.71	49.39	2.57	1.80
12	NONWA 4	0.00007	0.00005	0.00004	0.00003	0.02	0.11	0.006	0.004
13	BOROBAR1	0.021	0.006	0.020	0.005	0.28	19.91	1.03	0.72
14	BOROBA 2	0.097	0.023	0.062	0.012	0.66	45.98	2.88	1.67
15	BOROBA 3	0.065	0.015	0.040	0.008	0.42	29.49	1.53	1.07
16	BOROBA 4	0.083	0.020	0.057	0.012	0.65	45.78	2.38	1.66
17	KOROKOR1	0.140	0.032	0.079	0.013	0.73	50.87	2.64	1.85
18	KOROKO 2	0.083	0.020	0.056	0.011	0.63	44.04	2.29	1.60
19	KOROKO 3	0.141	0.033	0.080	0.014	0.74	51.98	2.70	1.89
20	KOROKO 4	0.144	0.033	0.081	0.014	0.76	53.39	2.77	1.94
21	CONTROL 1	0.196	0.047	0.127	0.025	1.35	94.42	4.90	3.43
22	CONTROL 2	0.146	0.034	0.083	0.014	0.76	53.52	2.78	1.95
Mean		0.11507	0.0271	0.07063	0.01343	0.74	0.05172	2.69	1.88

E_T = Total Annual Effective, FCR = fatality cancer risk to adult per year, LFCR = lifetime fatality cancer risk to adult, SHE = severe hereditary effect in adult per year, ELHE = Estimated lifetime hereditary effect in adult.

In order to evaluate the radiation risk due to ingestion of the selected radionuclides, the ICRP methodology was adopted in this study and the results shown in Table 3. The results of the cancer and non-cancer risk components were evaluated from the estimated annual effective dose of the sampled water. The result of the evaluated fatal cancer risk to adult per year in each sample ranged from 0.02×10^{-3} (NONWA 4) to 2.83×10^{-3} (KPITE 2) with an associated lifetime fatality cancer risk of 0.11×10^{-3} (NONWA 4) to 198.11×10^{-3} (KPITE2). The evaluated lifetime hereditary effect to adult per year varied from 0.006×10^{-5} (NONWA 4) to 10.29×10^{-5} (KPITE 2) with an associated lifetime hereditary effect in adult of 0.004×10^{-3} (NONWA 4) to 7.20×10^{-3} (KPITE 2).

This means that in terms of the lifetime fatality cancer risk to adult, approximately 2 out of 1000 may suffer from some form of cancer fatality and for the lifetime hereditary effect, approximately 7 out of 1000 suffer some hereditary effects. The negligible cancer fatality risk value recommended by USEPA is in the range of 1.0×10^{-6} to 1.0×10^{-4} (ie 1 person out of 1 million or 10,000 suffering from some form of cancer fatality is considered trivial). Comparing the estimated results of the lifetime fatality cancer risk in this study with the acceptable risk factor, it can be inferred that all the estimated results of the lifetime fatality risk in adult citizen of selected communities of Tai Local Government Area of Rivers State, Nigerian population due to ingestion of radionuclide in hand dug well water are relatively higher than the range of acceptable risk values recommended by USEPA [18].

5. Conclusion

The annual effective dose and lifetime cancer risk due to natural radioactivity in ground water of Tai Local Government Area of Rivers State, Nigeria has been estimated using Gamma spectroscopy techniques and some radiation models. The activity profiles of the radionuclides have clearly shown high activity concentrations across the sampled hand dug well waters.

The estimated annual effective dose and lifetime fatality risk in adult member of population of the selected communities of Tai, Rivers State due to ingestion of radionuclide in the sampled water are higher than the range of acceptable safe values and acceptable risk values recommended by USEPA as contained in Ndontcheung *et al.*,(2013).

Based on the findings of this work, we can conclude that hand dug well waters are not suitable for human consumption especially infants due to their shallow depths and susceptible to high level of pollution and therefore presents some significant radiological risk related to its life-long consumption. The result of this work serves as baseline radioactivity data for future studies in the study area.

References

- [1] Stabans E.I and Kinsars (2014). Radioactivity in the groundwater of a high background radiation area. *Journal of environmental Radioactivity*, 137.
- [2] Saghati F. and Salouti M. (2012). natural radioactivity in drinking water of drilled wells in Zanjan, NorthWestern Iran. *J. Biomed Phys* 2(1):27-32.
- [3] Ajayi, O.S. and Owolabi (2008). Determination of natural radioactivity in drinking water in private dug wells in Akure, SouthWestern , Nigeria. *Radiation Protection Dosimetry* 128(4): 477-484.
- [4] Ononugbo C. P, Avwiri G. O and J. M. Egieya (2013). Evaluation of natural radionuclide content in surface and groundwater and excess life time cancer risk due to gamma radioactivity. *Academic Research International*, 5(3):79-84
- [5] Uosif L.M. Issa, S.A.M., Issa M.A.M. and Abd El-Salam (2012). Natural radionuclide concentration in granite rocks in Aswan and Central Southern Eastern desert, Egypt and their radiological implications. *Radiation protection dosimetry*, 150(4):488-495.
- [6] Agbalagba, E.O., Gregory O. Avwiri, Yehuwadah E. Chad-Umoren (2013). Gross alpha and beta activity concentration and estimation of adults and infants dose intake in surface and ground water of Ten oil fields environment in Western Niger Delta of Nigeria.
- [7] Nwankwoals H.O and Walter I.O (2012) Assessment of Groundwater quality in Shallow Coastal Aquifer of Okirika Island, Eastern Niger Delta Nigeria Ife . *Journal of Science*, 14(2): 21-29
- [8] Ugbede, F.O (2018): Measurement of background ionizing Radiation Exposure levels in selected farms in communities of Ishielu L.G.A, Ebonyi State, Nigeria. *Journal of Applied Science Environmental Management* 22(9) 1427-1483.
- [9] Ajayi O.S. and Adesida G. (2009). Radioactivity in some sachet drinking water samples produced in Nigeria. *Iran J. Radiat Res.* 7(3):151-158.
- [10] Onoja, R.A. (2004). Survey of gross alpha and beta radioactivity in well water from Zaria area. M.sc thesis. Unpublished.
- [11] IAEA (2010). Programmes and systems for source and Environmental radiation monitoring. IAEA Report Series No. 64. International Atomic Energy Agency Vienna.
- [12] ICRP (2012). Compendium of Dose coefficients based on ICRP publication 60. ICRP publication 119, Ann. ICRP 41(suppl.)
- [13] Ndontchueng M.M., Simo A., Nguelem, E.J.M. Beyala, J.F. and Kryeziu 92013). Preliminary study of Natural Radioactivity and Radiological Risk Assessment in some Mineral Bottled water Produced in Cameroon. *International Journal of Science and Technology* 3(12):372 – 377

- [14] ICRP (2007). The 2007 Recommendations of the International commission on radiological protection. Annals of ICRP. Publication 103.
- [15] WHO (2008). Guidelines for drinking water quality in cooperating First Addendum 1, Recommendations, 3rd edition Radiological Aspect Geneva: World Health organization
- [16] Rusconi R., Forte M. Badalamenti P., Bell-inzona S. Gallini R. Maltese S Romeo C. and Sgorbati G.(2003). The monitoring of tap water in Milano: planning, methods and results. Lombardia, Italy: internal Report ARPA.
- [17] United Nations Scientific Committee On the effects of Atomic Radiation (UNSCEAR 2000). Sources and effects of ionizing Radiation (Report to the General assembly) New York: United Nations.

UNDER PEER REVIEW