

1 **Influence of Kraal Manure, Chicken Manure and Inorganic Fertilizer on Growth, Yield**  
2 **and Post-harvest Quality of Pepper (*Capsicum annuum* L.) in a Sub-tropical**  
3 **Environment**

4 **ABSTRACT**

5 *The excessive unjustified use of some kinds of fertilizers has seen some farmers realizing poor*  
6 *quality fruit that does not appeal to the final consumer, thus negatively affecting the effort of*  
7 *alleviating poverty in the Kingdom of Eswatini. This experiment was carried out at the*  
8 *Horticulture Department Lath House, Faculty of Agriculture, Luyengo campus of the*  
9 *University of Swaziland to determine the growth, yield and shelf life of green pepper when*  
10 *fertilized with kraal manure, poultry manure and inorganic fertilizer. The experiment was*  
11 *conducted to find the optimum levels of fertilizers that promotes the growth of pepper and to*  
12 *find the effects of different fertilizers on yield and quality of pepper. The treatments were of*  
13 *kraal manure applied at 60tones/hectare, 2:3:2 (37) at 370 kg/ha and chicken manure at 40*  
14 *tonnes per hectare. The results showed that growing pepper using the three treatments*  
15 *significantly affected its growth rate, leaf number, fruit number and its (fruit) shelf life.*  
16 *Pepper grown using inorganic fertilizer had the highest leaf number as compared to chicken*  
17 *manure and the least was recorded in kraal manure. There were no significant ( $P>0.05$ )*  
18 *difference in the growth rate of the inorganic fertilizer and chicken manure treatments. There*  
19 *was also no significant ( $P>0.05$ ) difference of pepper grown with chicken manure and*  
20 *chemical fertilizer in the number of days it took the harvested pepper to 100% decay stored at*  
21 *room temperature at 21 days. Yet pepper grown with kraal manure was significantly different*  
22 *from the two as it showed 20% decay rate in the same number of days. Kraal manure at 60*  
23 *t/ha is recommended in the production of pepper with a longer shelf life.*

## 24 INTRODUCTION

25 Green pepper (*Capsicum annuum L.*) is amongst the most important vegetables grown in  
26 most countries with warm climate. Recently large scale production of pepper for export has  
27 been undertaken under irrigation in Southern Africa (Norman, 1992). Green pepper  
28 originated from Central and Tropical America. The fruit is berry like tomato but with large  
29 locules without the gel with its seeds tightly compressed to the central stalk (Norman, 1992).

30 Green pepper is a warm season crop and its growth and development is similar to that of  
31 tomato but requires relatively higher temperatures. Soils preferred are sandy soils or loamy  
32 soils with a lot of organic matter well drained soils and pH ranging from 5.5 - 6.8 is best for  
33 its successful production. The crop may be directly seeded or transplanted to the field. Soil  
34 moisture must be relatively uniform throughout the growing season for optimal production.  
35 The number and frequency of irrigation will depend on the type of soil, developmental stage  
36 of the plant, atmospheric temperature and humidity (Norman, 1992).

37 Vegetables such as green pepper are very important nutritionally and economically in  
38 Swaziland. Pepper is used as a vegetable, salad or to add flavour in stews. It is also used for  
39 medicinal purposes to cure fever and colds (Norman, 1992). The challenges observed though  
40 are those of proper nutrition for specific vegetables. It is important to ensure that there is  
41 enough nutrients to hasten the plant growth so that flowering fruit setting does not occur  
42 whilst the plant structure is still small to carry the load of fruits.

43

44

45 Farmers in Swaziland normally use both inorganic and organic fertilizers in vegetable  
46 production. The question is the amount and effects of the fertilizer of choice on yield and  
47 product quality.

48 Most local farmers apply kraal manure (Boma) on their plots, some use chicken manure and  
49 most use commercial (synthetic) fertilizers. Synthetic fertilizers commonly used by farmers  
50 in Swaziland include 2:3:2 (22), 2:3:2 (38) and straight fertilizers such as Lime Ammonium  
51 Nitrate (LAN) or Urea but the latter is rarely used. A general recommendation is to apply  
52 250 kg 15-15-15 NPK prior to planting (Norman, 1992), but it is wise to test the soil for  
53 nutrient status and apply fertilizers/soil ammendments as recommended.

54 Swazis enjoy their meals with vegetables. The challenges observed are that farmers that  
55 produce the vegetables use all types of fertilizers without exactly knowing the benefits and  
56 demerits of the various fertilizers. The amount applied is also another challenge that needs to  
57 be addressed. Product quality is also a point of concern, including nutritional status and shelf  
58 life of the final product. However, growth response of plants differs with different fertilizers  
59 due to their differences in nutrient and other element composition. Therefore, the  
60 determination of the best source of nutrition is one of the fundamentals for effective plant  
61 production to meet consumer satisfaction. Cost effective means of production are a necessity  
62 to effectively produce safe, healthy and adequate food thereby alleviating poverty and  
63 building a healthy nation.

64 **To find out the effects of different fertilizers on yield and quality of pepper....etc**

65

66 Since the use of fertilizers has been introduced in horticulture, different fertilizer sources  
67 have been used which have different compositional properties that causes differences in  
68 growth rate and yield of plants and quality of the final product. The excessive unjustified use

69 of some kinds of fertilizers has seen farmers realizing poor yields of poor quality fruit that  
70 does not appeal to the final consumer, thus negatively affecting the effort of alleviating  
71 poverty in the Kingdom of Eswatini.

72 **The objectives of this research paper are to ....**

### 73 **MATERIALS AND METHODS**

74 Experimental site: This experiment was conducted in the lath house of the Horticulture  
75 Department, University of Swaziland, Luyengo Campus during the summer from November  
76 2012 to February 2013. It is located at Luyengo between latitude 26°34 'S and 31°12' E at  
77 750m above sea level with an average of temperature 21°C and receives about 800mm of  
78 rainfall per annum. The soil at the experimental site is classified under Malkerns series, deep  
79 red loam. Ferrasolic or merely a ferralitic soil intergrades to fersialitic soils or typical utisols.  
80 The soil was mixed with sand to create a sandy loam (Brady and Weil, 2007).

81 **Plant Material:** Six week old green pepper seedlings of the cultivar 'Mayo' obtained from  
82 Vickery Seedlings, Malkerns. They were transplanted on the 2<sup>nd</sup> November 2012. The pepper  
83 seedlings were transplanted in pots in medium that was prepared in advance. The medium  
84 comprised of top-soil obtained from the campus farm, river sand and sawdust at the ratio  
85 1:1:1. The medium was steam sterilized to kill unwanted microorganism and soil borne  
86 diseases like bacterial wilt. The pots were arranged into four blocks with 4 treatment, 4 pots  
87 per row and replicated 4 times. The plants were provided with optimal growing conditions,  
88 and all cultural practices according to need, for example irrigation, weeding, and pest and  
89 disease control.

90

91 **Nutrient Sources:** Three types of nutrition sources were used as treatments. 2:3:2 (37) was  
 92 used as a source of inorganic fertilizer. Two types of organic fertilizers were used, kraal  
 93 manure was sourced from the Luyengo animal dairy section while chicken manure was  
 94 sourced from the poultry section of the Luyengo campus.

95 The organic and inorganic soil ammendments were applied 2 weeks after transplanting  
 96 (WAT).

97 Samples were analysed for pH status at the Malkerns Research Station Soil Testing Unit. It  
 98 was found that adjustments were necessary at the rate of 1 tonne per hectare. The pH was  
 99 below the acceptable level as it was found to be 5.1. The pH had to be raised to acceptable  
 100 levels of 5.5 – 7.0.

101 Table 2: Nutrient composition per 1kg of poultry manure

Nutrient	Concentration per kg of manure
Iron	256mg
Cadmium	20mg
Zinc	72mg
Copper	96mg
Manganese	20mg
Phosphorus	180mg
Potassium	84mg
Magnesium	240mg

Calcium	1372mg
pH	6.92

102

103 Table 3: Nutrient composition per 1kg of cattle manure

Nutrient	Concentration per kg of manure
Iron	0.19mg
Cadmium	0.02mg
Zinc	0.11mg
Copper	0.12mg
Manganese	0.02mg
Phosphorus	0.18mg
Potassium	0.14mg
Magnesium	0.38mg
Calcium	2.04mg
pH	6.97

104

105

106

107


108 **Experimental Design:** The experiment was a 1x3 factorial experimental laid in a randomized  
109 complete block design (RCBD) with three types of fertilizers for nutrition. Plant pots were  
110 arranged in blocks and were laid down on the ground in the lath house with 80% light  
111 transmission.

112 **Data Collection and Analysis:** Data were taken on a fortnightly basis and recorded until  
113 harvest and through postharvest. Three plants were randomly selected at the beginning of the  
114 experiment in each of the replications and the following parameters were measured: plant  
115 height, number of leaves, leaf area index, time to flowering, number of flowers, number of  
116 fruits harvested per plant, fruit weight(g), percentage of nonmarketable fruits, shoot and root  
117 dry mass determined.

118 Data collected were analysed using MSTAT-C (Nissen, 1989). Analysis of variance  
119 (ANOVA) was done on plant height, number of leaves, leaf area, leaf area index, number of  
120 flowers, number of fruits produced, number of harvested fruits, fruit weight, percentage of  
121 non-marketable fruits, and shoot to root ratio. Mean separation was done using Duncan's  
122 New Multiple Range Test (DNMRT) where the F test showed significant differences (Gomez  
123 and Gomez, 1994).

124 **RESULTS**

125 **Plant height:** Plant height increased steadily in the different fertilizer treatments with plant  
 126 growth up to 9 WAT. Plant height was significantly ( $P<0.05$ ) affected by the different  
 127 fertilizers. The highest plant height of 46 cm was obtained in chicken manure treatment  
 128 whilst the lowest (8cm) was obtained in control 9 (WAT) (Figure 1). There was no significant  
 129 difference ( $P<0.05$ ) in plant height of pepper treated with 2:3:2(37) and chicken manure in  
 130 block 2 after 9 WAT, but both were significantly higher than plants treated with kraal manure




131 and control.

132 Figure 1: Effects of different fertilizers on plant height of green pepper. Bars are standard  
 133 error below and above the mean.

134



135 **Number of leaves:** The number of leaves per plant was significantly ( $P < 0.05$ ) affected by the  
 136 different fertilizers. Plants treated with synthetic fertilizer (2:3:2 (37) had the highest number  
 137 of leaves while those treated with kraal manure had the lowest number of leaves (22) (Figure  
 138 2). However there was no significant ( $P > 0.05$ ) difference in the number of leaves produced  
 139 by green pepper treated with kraal manure and chicken manure (15 and 17 respectively) at 6  
 140 WAT.



141

142 Figure 2: Effects of different fertilizers on number of leaves of green pepper. Bars are  
 143 standard error below and above the mean.

144

145

146


147

148

149

150

151 **Leaf area:** There were significant ( $P < 0.05$ ) differences in leaf area of green pepper plants for  
152 the various fertilizer treatments. Plants treated with synthetic fertilizer (2:3:2 (37) had the  
153 highest leaf area than in kraal manure but not significantly ( $P > 0.05$ ) higher from pepper  
154 treated with chicken manure at 9 WAT (Figure 3).




155

156 Figure 3: Effects of different fertilizers on leaf area of green pepper. Bars are standard error  
157 below and above the mean.

158

159 **Leaf area index:** The leaf area index (LAI) was significantly ( $P < 0.05$ ) affected by the  
160 different fertilizers. Plants treated with synthetic fertilizer (2:3:2 (37) had the highest LAI  
161 (13.6 cm) cm than those treated with kraal manure and chicken manure. Plants treated with  
162 kraal manure had the lowest LAI (6.7cm) (Figure 4).




163

164 Figure 4: Effects of different fertilizers on LAI of green pepper plants. Bars are standard error  
165 below and above the mean.

166

167 **Number of flowers:** There were significant ( $P < 0.05$ ) differences in number of flowers  
168 produced per plant treated with different fertilizers. Plants treated with synthetic fertilizers  
169 (2:3:2 (37) and chicken manure produced significantly ( $P < 0.05$ ) higher number of flowers  
170 (26) than plants treated with kraal manure (17) (Figure 5). A variation in the flowering dates  
171 among the treatments was observed. Plants treated with 2:3:2(37) flowered first at 35 days  
172 after transplanting (DAT), followed by plants treated with chicken manure (39 DAT) and  
173 green pepper treated with kraal manure flowered last at (42 DAT).



174

175 Figure 5: Total number of flowers on plants from different fertilizer treatments


176 Bars with the same letters are not significantly different ( $P > 0.05$ ) from one another. Mean  
177 separation by Duncan's New Multiple Range (DNMRT).

178

179

180

181 **Fruit number:** There were significant ( $P < 0.05$ ) differences in the number of fruits produced  
182 by green pepper plants treated with the different fertilizers. Plants treated with chicken  
183 manure produced the highest number of fruits (28) followed by pepper treated with 2:3:2 (37)  
184 at (25) while lowest number of fruits were obtained from fruits treated with kraal manure (16)  
185 (Figure 6). However the number of harvested fruits was significantly ( $P < 0.05$ ) affected by the  
186 different fertilizer treatments (Figure 7). Highest number of harvested fruits (9.8) was  
187 obtained in pepper treated with 2:3:2(37) and chicken manure and the lowest were obtained  
188 in pepper treated with kraal manure (3.5).



189

190 Figure 6: Total number of fruits produced by plants from different fertilizer treatments

191 Bars with the same letter are not significantly different ( $P > 0.05$ ) from one another. Mean  
192 separation by Duncan's New Multiple Range Test (DMRT).


193

194

195

196

197



198


199

200 Figure 7: Effects of different fertilizers on total number of harvested fruits

201 Bars with the same letter are not significantly different ( $P>0.05$ ) from one another. Mean

202 separation by Duncan's New Multiple Range Test (DNMRT).


203 **Change of colour:** The change of colour of fruits stored at room temperature was  
204 significantly ( $P < 0.05$ ) affected by the different fertilizer treatments (Figure 8). The highest  
205 change of fruit colour was observed on fruits from plants previously treated with chicken  
206 manure (3) and the lowest was obtained on fruits from plants previously treated kraal manure  
207 (2)



208

209 Figure 8: Effects of fertilizer treatments on colour change of fruits from green to red stored at  
210 room temperature. Bars with the same letters are not significantly different ( $P > 0.05$ ) from  
211 one another. Mean separation by Duncan's New Multiple Range Test (DN MRT).

212 **Fruit Shrinking:** The different fertilizer treatments significantly ( $P < 0.05$ ) affected the  
213 shrinkage of fruits during storage. Fruits harvested from chicken manure and synthetic  
214 fertilizer (2:3:2 (37) showed a significant difference in the number of fruit shrinking. Fruits  
215 treated with 2:3:2(37) started to shrink 3 days after harvest (DAH) while those treated with  
216 chicken manure started to shrink at 6 DAH (Figure 9).




217

218

219 Figure 9: Effects of the different fertilizers on the rate of shrinking of pepper harvested and  
220 stored at room temperature. Bars with the same letter are not significantly different ( $P > 0.05$ )  
221 from one another. Mean separation by Duncan's New Multiple Range Test (DNMRT).



222 **Fruit Decay:** The different fertilizer treatments significantly ( $P < 0.05$ ) affected the rate of  
 223 fruit deterioration when stored at room temperature. Fruits fertilized with 2:3:2 (37) and  
 224 chicken manure had a 100% fruit decay/ rot at 14 days after harvest (DAH), while fruits from  
 225 plants fertilized with kraal manure started to rot 21 DAH (Figure 10).



226  
 227 Figure 10: Percentage of decayed fruits at 21 days after harvest (DAH). Bars with the same  
 228 letters are not significantly different ( $P > 0.05$ ) from one another.

## 230 DISCUSSION

231 Different types of organic and inorganic fertilizers had varying effects on the growth and  
 232 yield of green pepper. The highest leaf number was recorded in plants grown from chemical  
 233 fertilizer applied at recommended rates of 370kg which was significantly ( $P < 0.05$ ) different  
 234 from chicken manure applied at 40 t/ha and the lowest was obtained from the kraal manure at  
 235 60t/ha.

236 These differences may be due to the fact that the growth medium of all the treatments were  
 237 added with the same amounts of sawdust that ensured the same water holding capacity in all  
 238 blocks and that the treatments of chemical fertilizers had nutrients readily available for

239 uptake by plants. The other two treatments had to undergo organic matter breakdown by  
240 micro-organisms before nutrients were released for plant uptake thus delaying the availability  
241 of nutrients (Jacobs *et al.*, 2003). As the number of weeks increased a steady increase was  
242 obtained from the organic fertilizer treatments. Most probably as the manure decomposed the  
243 nutrient availability was increased and that the water holding capacity increased in the  
244 manure treatments. Replacement of chemical fertilizer by organic manures has been reported  
245 to enhance soil biological activity, efficiency and the rate of microbial substrate use (Van  
246 Averbek and Yoganathan, (2003). Increased vegetable yield with the use of manure have  
247 been previously reported for okra (Ogunlela *et al.*, 2005).

248 The mean leaf area and yield between the three treatments showed some variations. Overall  
249 chemical fertilizer applied at 370kg/ha had the highest leaf area and yield. However yield of  
250 pepper fertilized with chemical fertilizer was not significantly ( $P>0.05$ ) different from that of  
251 chicken manure fertilizer plants. Kraal manure fertilized plants had the lowest mean leaf area  
252 and yield compared to the other treatments. These results showed that the release of nutrients  
253 for plant utilization was delayed. If the organic fertilizers were given enough time to  
254 decompose before planting the results would possibly have been different as reported by  
255 (Gandy *et al.*, 2002).

256 The rate of shrinking of harvested fruits fertilized with chemical manure was significantly  
257 ( $P<0.05$ ) different from those grown with chicken manure and the least affected was kraal  
258 manure fertilized plants. This trend was also evident in the total number of days it took the  
259 fruits to start rotting when stored at room temperature.

260 After 21 days a 100% rotting of stored fruits was recorded for fruits previously fertilized with  
261 chicken manure and chemical fertilizers. This may be due to the increased content of

262 elements than in kraal manure which recorded 20% at the same number of days (Ferguson  
263 and Ziegler, 2004)

## 264 **CONCLUSIONS AND RECOMMENDATIONS**

265 The results of this study showed that the highest growth rate and yield of green pepper was  
266 obtained in plants treated with chicken manure (60 t/ha) but the highest yield was obtained in  
267 the chemical fertilizer treatment followed by chicken manure which both had a 100% rot rate  
268 at 21 days of storage compared to the 20% of kraal manure in the same number of storage  
269 days.

270 The applications of kraal manure at 60 t/ha are recommended for farmers to obtain products  
271 with a higher shelf life. Consumers who do not have the necessary storage means will benefit  
272 by just storing their vegetables at room temperature. The results suggest that it is possible to  
273 produce higher vegetables yield with longer shelf life through organic farming than that of  
274 conventional farming with inorganic fertilizer. It is recommended that more research be  
275 conducted to establish the optimum period of applying organic fertilizer before planting and  
276 to validate the recommendation.

## 277 **REFERENCES**

- 278 Aiyelaagbe, I.O.O. and Fawusi, M.O.A. (1986). Growth and yield of pepper to mulching.  
279 *Biotronics* 15:25-29.
- 280 Bauer, S., Schulte, E., Their, H.P. (2005). Composition of the surface waxes from bell pepper  
281 and eggplant. *European Food Research and Technology*, 220 (1): 5-10.
- 282 Blay, E.T. and Aflakpui, G.K.S. (1985). Effects of stage of harvesting on the quality and  
283 shelf life of tomato (*Lycopersicon esculentum* Mill.) *Legon Agriculture Research*  
284 *Bulletin*. 1:1-12.

- 285 Borovsky, Y., and Paran, I. (2008). Chlorophyll breakdown during pepper fruit ripening in  
286 the chlorophyll retainer mutation is impaired at the homolog of the senescence  
287 inducible stay-green gene. *Theoretical and Applied Genetics*, .117(2): 235-240.
- 288 Brady, N and Weil.R. (2007). *Nature and properties of soils*, 13<sup>th</sup> edition, Prentice Hall,  
289 New York. USA.
- 290 Browder, J.O. (1990) Ecosystem Science for the Future *BioScience*, 40 (9):
- 291 Deborah, K. (2006). "The case against synthetic fertilizers," San Francisco Chronicle,  
292 1/14/06.<http://www.sfgate.com/cgi-bin/article.cgi?f=/c/a/2006/01/14/HOG71>  
293 GLP6A1.DTL Accessed 5/19/09.
- 294 Eghball, B. (2001). Composting manure and other organic residue. Cooperative Extension  
295 (*NebGuide*), *Institute of Agriculture and Natural Resources, University of Nebraska*,  
296 Lincoln, Nebraska, USA.
- 297 FAO. (2008). *Organic Materials as Fertilizers*. Longman/FAO. Rome, Italy
- 298 Ferguson, J. and Ziegler, M. (2004). Guidelines for purchase and application of poultry  
299 manure for organic crop production.<http://edis.ifas.ufl.edu/HS217>
- 300 Gomez, A.A., and Gomez, K.A. (1994). *Statistical procedures for agricultural research*.  
301 John Wiley and Sons, New York, New York, USA.

- 302 Grandy, A.S., Porter, and Enrich, N.S. (2002) Organic management and crop rotation  
303 effects on the recovery of organic matter and aggregation in potato cropping  
304 systems. *Soil Science Society Journal* 66: 1311-1319.
- 305 Haynes, C. (2008). The Organic Green Revolution.  
306 [Http://www.rodaleinstitute.org/files/ GreenRevUP.pdf](http://www.rodaleinstitute.org/files/GreenRevUP.pdf) retrievecontent/12/06/2008.
- 307 Hamilton, R. (2009 “Agriculture’s Sustainable Future: Breeding Better Crops”. *Scientific*  
308 *American Journal*, [http://www.:edis.future.ufl.edu/HS217.retrievecontent/09/24/2009](http://www.edis.future.ufl.edu/HS217.retrievecontent/09/24/2009).
- 309 Jacobs, R.D., Sloan, D. and Jacob J.(2003). Cage layer manure: *An important resource for*  
310 *land use* <http://edis.ifas.ufl.edu/ps005>.
- 311 Masarirambi, M.T., Dlamini, P., Wahome, P.K. and Oseni, T.O. (2012). Effects of chicken  
312 manure on growth, yield and quality of lettuce (*Lactuca sativa* L.) ‘Tiana’ under a  
313 lath house in a semi-arid sub-tropical environment. *American-Euasian Journal*  
314 *Agriculture and Environment* 12 (3): 399-406.
- 315 Ministry of Agriculture, Fisheries and Food (1976). *Organic manures*. London. UK.
- 316 Muse, J.K. (1993). *Inventory and Evaluation of Paper mill By-products for Land Application*.  
317 .Unpublished. M.Sc. Thesis, Auburn University, USA, pp. 9-13.
- 318 Nissen, O. (1989). *MSTAT-Ca micro computer programme design, management and*  
319 *analysis of agronomic research projects*. Michigan State University, East Lansing,  
320 Michigan, USA.

- 321 Norman, J.C. (1977b). Effects of age of transplants on hot pepper, (*Capsicum sinense*)  
322 *Acta Horticulturae*. 53: 43-48.
- 323 Norman, J.C. (1992) Tropical Vegetable Production, Auther H. Stockwell L.T.D, Elms Court  
324 Ifracombe, Devon, U.K.
- 325 Ogunlela, V.B., Masarirambi, M.T. and Makuza, S.M. (2005). Effects of cattle manure  
326 application on pod yield and yield indices of okra (*Abelmoschus esculentus* L.  
327 Moench) in semi-arid and subtropical environment. *Journal of Food, Agriculture*  
328 *and Environment* 3:5-15.
- 329 Ossom, E.M. (2005). Effects of weed control methods on weed infestation, soil temperature  
330 and yield in Swaziland. *UNISWA Research Journal of Agriculture, Science and*  
331 *Technology* 8: 5-15.
- 332 Pilon, L., Oetterer, M., and Gallo, M.H.F. (2006). *Shelf life of minimally processed carrot*  
333 *and green pepper*. Vol. 26. No.1, pp. 150- 158.
- 334 Pimentel, D., Hepperly, P., Hanson, J., Douds, D. and Seidel, R. (2005). Environmental,  
335 energetic and economic comparisons of organic and conventional farming systems.  
336 *Bioscience*. 55(7): 573-582.
- 337 Van Averbek, A. and Yoganathan, S. (2003). *Using kraal manure as a fertilizer*.  
338 Department of Agriculture. Pretoria. South Africa  
339