

## Original Research Article

# STRUCTURE OF MACROZOOBENTHOS (GASTROPODS) COMMUNITY IN MANGROVE FOREST ECOTOURISM PANDANSARI KABUPATEN BREBES, CENTRAL JAVA

### ABSTRACT

**Aims:** This research aims to analyze the abundance and diversity of macrozoobenthos in the Gastropod Class in the Ecotourism Waters of Pandansari Mangrove Forest, Brebes Regency, Central Java. The research was conducted in the Ecotourism Waters of Pandansari Mangrove Forest, Brebes Regency, Central Java

**Study design:** The research was conducted survey

**Place and Duration of Study:** This research was carried out for 3 months between December 2018 until February 2019 in Ecotourism Forest Areas in Bakau Pandansari, Brebes Regency, Central Java

**Methodology:** The method used in this research is survey methods using primary data in the form of physical chemistry of aquatic data, gastropod abundance, diversity index, equatibility index. Data collection techniques using purposive sampling in determining 3 stations and 3 repetitions. Data analysis in this research uses descriptive analysis.

**Results:** Research results found 9 species of Gastropods that settled in the mangrove ecosystem, namely *Casidulla aurisfelis*, *Cerithidea* sp, *Cerithidea alata*, *Cerithidea cingulata*, *Cerithidea obtusa*, *Polinices* sp, *Puperita* sp, *Telescopium telescopium*, and *Turiculla nellial-suprius*. Gastropod abundance ranges from 67 - 166 ind / m<sup>2</sup>. Gastropod diversity index in the medium category  $2 < H' \leq 3$ . Uniformity values ranged from 0.79 to 0.81 with a high category. The relationship between water quality and gastropods has an R<sup>2</sup> value of 97%, water quality can affect gastropod life and as much as 3% is influenced by other factors.

**Conclusion:** Based on the results of research that has been done, it can be concluded that in general environmental parameters in the Pandansari mangroves are still viable in supporting the survival of the gastropods contained in them.

**Comment [J1]:** Scientific names in italic

**Keywords:** [*Gastropods*, *Abundance*, *Diversity Index*, *Equatibility Index*]

**Comment [J2]:** Replace keywords that are already in the title

### 1. INTRODUCTION

Brebes is one of the districts in Central Java reside in the north coast of Java (Pantura). The North Coast region of Brebes has a coastal area that stretches around 54 kilometers with mangrove forests covering an area of approximately 852 hectares from Brebes, Losari Sub-district. Brebes mangrove

forests are used for timber, some land is used for shrimp ponds and recently for salt production. So that on 2007 it was estimated that the remaining mangrove forest in Brebes was 275 hectares [1]. Mangrove area having damaged due to abrasion and excessive logging. Starting from 2000 to 2008, abrasion Brebes reached 640.45 hectares with a coastline of 27.043 km in the period of 8 years or 29.60 m per year[1]. One of the regions with the highest abrasion level is located in Kaliwlingi Village, Brebes Subdistrict, reaching 385.98 hectares. The activities of mangrove rehabilitation in the coastal area of Brebes Regency were coordinated and carried out by the Department of Agriculture, Forestry and Soil Conservation of the Regency of Brebes which was realized since 2004[2]. Efforts to rehabilitate mangrove forests from 2008 to 2017 from one stakeholder have planted as many as 3,500,000 mangrove stems[3]. The most commonly planted species is *Rhizophora* sp because it is relatively resistant to estuary environmental fluctuations-

Comment [J3]: Italic

Mangrove forests are dominated by several species of distinctive trees or shrubs that have the ability to grow in salty waters[4]. Mangrove forest, creating a healthy climate atmosphere aquatic biota, and has contribute to balance the biological cycle in the waters. Mangrove has a very important function for feeding ground, as the venue for biological processes such as spawning ground, nursery grounds some aquatic organisms one macrozoobenthos[5]. Macrozoobenthos is a group of invertebrate animals that live in sticking to the substrate at the bottom of the body of the water. Macrozoobenthos are generally relatively sedentary or can move with slow movements [6]. Macrozoobenthos in mangrove forests mostly live on hard substrates to mud. Generally macrozoobenthos in the Indonesian mangrove region are macrozoobenthos from the Gastropod, Bivalvia, Crustacea, and Polychaeta[7].

Gastropod is a class that has a very wide spread, because it has a fairly high adaptability [4]. Gastropods are part of the mangrove ecosystem and have a very complex role because their presence in aquatic ecosystems is unique. Unique because in addition to functioning as a food ingredient for other organisms, it can also be used as an indicator of the quality of a waters[8]. In the mangrove ecosystem , the existence of the macrozoobent as very closely related to water quality and the type of sediment that exists because in addition to water quality, sediment is a living medium that is directly in contact with macrozoobenthos [8]. Therefore the research entitled "Community Structure Macrozoobenthos (Gastropod) On Bodies Ecotourism Mangrove Forest Pandansari Brebes, Central Java" needs to be done. This research observes and analyzes species diversity and Gastropod abundance which can be used as bioindicators of the environment and is associated with abiotic factors in mangrove ecosystems for the sustainability of mangrove forest conservation.

## 2. MATERIAL AND METHODS

### 2.1 Time and Place of Research

This research was carried out on December 2018 to March 2019 at Ecotourism Pandansari Mangrove Forest located in Kaliwlingi Village, Brebes Regency, Central Java.

Comment [J4]: Add geographic coordinates of the study area

### 2.2 Materials and Tools

The tools used in this research are Global Posting System (GPS), quadrant transect, refractometer, DO meter, pH meter. poles, shovel, plastic bag. While the material used is, Gastropods and sediment.

### 2.3 Research Methods

The method used in this research is survey, determination of sampling stations using purposive sampling method. Gastropod sampling was carried out using the quadrant transect method. 1m x 1m quadrat size.

Station I is a muddy high density mangrove with a coordinate point S 6°47'8,70"-: E 109°2'16,62". Station II is a mangrove forest with a medium density of mud and gravel with a coordinate point of S 6°47 ' 13"-: E 109°2'22 ". Station III is a low density mangrove forest with muddy substrates with a coordinate point of S 6°47'9"-: E 109°2'20,23" Figure 1. Sampling was carried out three times with a sampling interval every month according to the fast life cycle of macrozoobenthos.




Figure 1. The map of study station located at Ecotourism Pandansari Mangrove Forest Brebes Central Java

## 2.4 Research Parameters

Data analysis was conducted using the Benthos density equation as shown below:

$$D_i = n_i/A$$

Whereas:

$D_i$  = Makrozoobenthos abundance (ind/m<sup>2</sup>)

$n_i$  = Number of makrozoobenthos that had been found

$A$  = Quadrat area (m<sup>2</sup>)

The next stage was to analyze the community structure as shown below:-

Diversity ( $H'$ )

$$H' = -\sum p_i \log_2 p_i$$

$$p_i = n_i/N$$

Whereas :

$H'$  = Indeks Diversity Shannon Wiener

$P_i$  = Proportion of species  $i$  ( $n_i$ ) to total number ( $N$ )

Equatibility ( $E'$ )

$$E = H'/H_{max}$$

## 2.5 Data Analysis

Statistical data analyzed for relation water quality between gastropods abundance using multiple regression and [correlation](#).

## 3. RESULTS AND DISCUSSION

Water quality is one of the main environmental factors in the fish farming system. Water quality is a physical and chemical factor that can affect the maintenance media environment and indirectly affect

**Comment [J5]:** To detail in the methods the physical and chemical parameters of water that were evaluated in the study.

the metabolic processes of test fish. Poor water quality can cause fish to become susceptible to disease [14,3].

### 3.1 Temperature

This research found 9 species of Gastropods which divided into 6 Genus. We found that total abundance of Gastropods in station 1 abundance of 166 ind/m<sup>2</sup>, station 2 abundance 144 ind/m<sup>2</sup>, and station 3 abundance 67 ind/m<sup>2</sup> (Table 1).

**Table 1. Abundance of Gastropods in study stations (ind/m<sup>2</sup>)**

Species	Station 1	Station 2	Station 3
<i>Casidulla aurisfelis</i>	63	55	26
<i>Cerithidea</i> sp	34	32	12
<i>Cerithidea alata</i>	3	7	4
<i>Cerithidea cingulata</i>	9	8	6
<i>Cerithidea obtusa</i>	18	15	0
<i>Polinices</i> sp	22	15	10
<i>Puperita</i> sp	9	6	5
<i>Telescopium telescopium</i>	6	5	2
<i>Turiculla nellial-suprius</i>	2	1	2
Total	166	144	67

**Comment [J6]:** Add author and year of taxon according to scientific nomenclature norms

The higher abundance of Gastropods was obtained in study station 1(166 ind/m<sup>2</sup>) which was dominated by *Cerithidea* sp-. These results are in agreement with Supratman [9] who reported that mangrove forest is the most preferred habitat for *Cerithidea* sp. Moreover, this species are thought to have a wide habitat and are able to live in various types of habitat both in the mangrove ecosystem, muddy to sandy beaches, and species are able to live in unstable conditions such as in mining areas. Overall Station 1 have a high abundance of mangroves, so that the production of mangrove waste at Station 1 is more than other stations. According to Nybakken [4] mangrove leaf decay will become an important organic material for organisms. In addition, the type of substrate station1 is mud-, this type is the most suitable for gastropods life. The abundance of Gastropods at station 3 is lower than all station because the location of the station is closer to the sea, the location of the substrate deeper makes the sampling technique difficult, besides that the substrate at the station is dominated by mud and waterlogged soil so that distribution of gastropods are expanded.

**Comment [J7]:** ?

### 3.2 Strucur Community

Diversity is identical to the stability of an ecosystem, if the diversity of an ecosystem is high then the condition of that ecosystem tends to be stable. Ecosystem environments that have diversity disorders tend to be moderate, and ecosystem environments that are polluted with diversity tend to be low. Diversity-Diversity index of Gastropods in the mangrove ecosystem of Pandansari Brebes can be seen in Figure 2. Gastropod diversity index from observations in each station included in the medium category  $2 < H' \leq 3$ . This condition shows that the distribution of the number individuals of each species is moderate. -This productivity is quite high, the condition of the ecosystem is balanced, and the ecological pressure is moderate. The highest diversity index was found at station 2, with a diversity index value of 2.555. The high diversity value of station 2 because the stability of the community and the distribution of the number of gastropods at station 2 was relatively evenly distributed. The lowest gastropods diversity index are at station 3 with a value of 2.506. The low diversity index value is due to the small number of gastropod species found compared to other observation stations. High and low levels of diversity are also influenced by sediment conditions and the quality of waters in the mangrove environment.

These results are in agreement with Suwodo [10] who reported diversity index will decrease with decreasing water conditions or quality. Species diversity is declared high if the community is

composed by many species and with the same or nearly the same species abundance, conversely if the community is composed of a few species and if only a few species are dominant then the diversity of species is low [11].

The equitability-equitability index is used to determine the evenness of the type in a community, the equitability index can also be used as an indicator of the presence of symptoms of species dominance in a community [12]. Uniformity values from the three stations ranged from 0.79 to 0.81 with a high category. Based on the criteria of uniformity index values it can be said that all species found at each station are almost evenly distributed. At each station the species were found almost the same, allegedly the equitable availability of food sources for Gastropods and in the area enabled Gastropods to defend themselves and reproduce well. Base of substrate is also an influencing factor because of the type of substrate that has mud texture and the content of organic material which is spread evenly in the mangrove ecosystem [5].




Figure 2. Graphic of Diversity Index and Equitability Index Gastropods

### 3.3 Relation of Water Quality with Gastropods

Result of regression test between abundance and water quality in Pandansari Mangrove Ecotourism in Brebes Regency, Central Java obtained  $y = 14.04 + 2.31 \text{ pH} - 0.22 \text{ DO} + 0.44 \text{ Salinity} + 0.09 \text{ temperature} - 14.60 \text{ c-organic} + 0.80 \text{ clay}$  with an R2 value of 0.97, which means as much as 97% of water quality can affect gastropod life and as much as 3% is influenced by other factors.

Tabel 2.

	Gastropods
pH	-0,3215973
DO	0,20078189
Salinity	-0,0641675
Temperature	-0,368975
c-organic	-0,9752935
clay	-0,4009145

The correlation coefficient is divided into several group such as 0.00 - 0.199 very low relation, 0.20 - 0.399 low relation, 0.40 - 0.599 moderate, 0.60 - 0.799 strong relation, and 0.80 - 1,000 very strong

**Comment [J8]:** Standardize text font throughout the manuscript, Times parts, Arial parts

**Comment [J9]:** Add x and y axis information; delete inner lines

**Comment [J10]:** Where and When? Inform.

**Comment [J11]:** Relativize the result. Of course, it is not just physical and chemical factors that define the occurrence and abundance of any organism in any environment.

Correlation Test

**Comment [J12]:** Add information about What, Where and When

relation [13]. Gastropods correlation test results on water quality namely pH 0.32 has a low relation, DO 0.20 has a very low relation, salinity 0.06 has a very low relation, temperature 0.36 has a low relation, c-organic 0.97 has the relation is very strong, and 0.40 clay has a moderate relation [Table 2](#). The c-organic correlation test has a very strong relation to influence the life of the gastropods, this is accordance with the existing criteria.

From this result, it was found that gastropod abundance was influenced by the conditions of the surrounding waters. Based on the correlation value, it was found that the abundance correlated most with organic carbon. The higher c-organic content, the gastropod abundance will tend to increase, this is in accordance with the statement Denisenko [14] high macrozoobenthos abundance is influenced by the high c-organic value at each station. The high c-organic in a waters generally will result in an increase in population [15]. In general, the results of environmental parameters in the Pandansari mangrove waters are still feasible in supporting the survival of the gastropods in them. Intensive community activities in the surrounding area will have an impact on the survival of gastropods in these waters.

#### 4. CONCLUSION

Based on the results of research that has been done, it can be concluded that in general environmental parameters in the Pandansari mangroves are still viable in supporting the survival of the gastropods contained in them.

**Comment [J13]:** Review the conclusion. It is not the environmental factors evaluated that define the occurrence and abundance of species. Actually, the evaluated environmental factors were correlated, but there are unevaluated factors that are also determinant.

#### REFERENCES

- [1] Mackay, P. 2012. Brebes Mangrove Restoration and Reforestation for Climate Change Adaptation Project. Central Java Green Belt Mangrove Corridor Program, Brebes Regency, Indonesia.
- [2] Suryono., Supriharyono., B, Hendrarto., O, K, Radjasa. 2015. Mapping of Mangrove Ecosystem Degradation and Coastal Abrasion Based on Geographic Information System in Brebes Regency, Central Java. *Oceatek*. 9(01).
- [3] Burhanudin, M. 2017. Kaliwlingi Proves Coastal Green Belt is Not an Impossible. <http://kehati.or.id/kaliwlingi-buktikan-sabuk-hijau-pesisir-bukan-hal-mustahil/>. Diakses pada tanggal 10 November 2018
- [4] Nybakken. 1992. *Marine Biology, An Ecological Approach*. PT. Gramedia. Jakarta.
- [5] Pramudji. 2000. Mangrove Forests in Indonesia: The Role of the Problem and its Processing. *Oseana*. 25(1) : 13-20.
- [6] Odum, E. P. 1993. *Ecological Basics*. Third edition. Gadjah Mada University press. Yogyakarta.
- [7] Arief, A. M. P. 2003. *Mangrove Forest Functions and Benefits*. Canisius Publisher. Yogyakarta.
- [8] Simanungkalit, V., dan Afrizal, T. 2013. Distribution of Macrozoobenthos In The Mangrove Forest Of Marine Protected Areas (MPA) Pariaman Of West Sumatra.
- [9] Supratman, O., Arthur, M. F., dan Jemi, F. 2018. Abundance and Gastropod Diversity in the Intertidal Zone on Eastern Bangka Island. *Enggano Journal*. 3(1) : 10-21.

**Comment [J14]:** Only 1/3 of the references are from the last five years. Update.

- [10] Suwondo, E. F., F. Sumanti. 2005. Gastropod Community Structure in Mangrove Ecosystems on Sipora Island, Mentawai Islands District, West Sumatra. *Biogenesis*. 2(1) : 25-29.
- [11] Janestia, R. N., Ali, S., dan Syahrul, P. 2017. Makrozoobentos Community Structure in the Mangrove Ecosystem Keudee Lung Waters, Pidie Jaya Regency. *Scientific Journal of Marine and Fisheries Unsyiah*. 2(3) : 406-414q.
- [12] Romdhani, A. M., Sukarno., Eko, S. 2016. The Biodiversity Of Gastropods Identified In The Mangrove Forest Of Baban Village, Gapura Districts Sumenep Regency As The Resource Of Learning Biology. *Jurnal Pendidikan Biologi Indonesia*. 2 (2) : 161-167.
- [13] Sugiyono. 2007. *Qualitative Quantitative Research Methods and R&D*. Bandung: Alfabeta.
- [14] Denisenko. S. G., N. V. Denisenko., K. K. Lehtonen., A. B. Andersin., A. O. Laine. 2003. Macrozoobenthos of the Pechora Sea (SE Barents Sea): community structure and spatial distribution in relation to environmental conditions. *Marine Ecology Progress Series*. Vol. 258: 109–123.
- [15] Asriani, W.O., Emiyarti. E. Ishak. 2013. Study of environmental quality around nickel loading ports (ni) and their relationship to the macrozoobenthos community structure in the waters of Motui Village, Konawe Utara Regency. *Jurnal Mina Laut Indonesia* 3(12) : 22 – 35