

# **ROCK PHOSPHATE, ZEOLITE AND QUAIL MANURE TO ENHANCE POTASSIUM UPTAKE AND YIELD OF SOYBEAN ON ALFISOLS**

## **ABSTRACT**

Soybean seeds are the source of vegetable protein-based that most consumed in Indonesia, but apparently, the production is unable to compensate the rate of increase in community needs. The aims of this study ~~was~~ ~~were~~ to investigate the effect of Rock Phosphate (RP), zeolite and quail manure to enhance potassium (K) uptake and yield of soybean in Alfisols. A field experiment was conducted in June-October with a single factor Randomized Complete Block Design (RCBD) consisting of 9 treatments and 3 blocks (replicates). The dosage of RP, zeolite and quail manure used ~~were~~ ~~was~~ 2.5 tons/ha and 5 tons/ha. The results showed that zeolite 5 tons/ha + quail manure 2.5 tons/ha tended to increase K uptake. The combination of RP 2.5 tons/ha + quail manure 5 tons/ha significantly ~~increase~~ ~~increases~~ the number of filled pods and seed weight per plot. RP 5 tons/ha + zeolite 5 tons/ha + quail manure 5 tons/ha also affects the number of filled pods and seed weight per plot. The treatment zeolite 2.5 tons/ha + quail manure 2.5 tons/ha also affects the number of filled pods.

*Keywords: Soybean, Alfisols, Rock Phosphate, Organic Manure, Zeolite*

## **1. INTRODUCTION**

Soybean seeds are the source of vegetable protein-based that most consumed in Indonesia. Soybean seed needs continue to increase along with increasing population growth, per capita income, and public awareness of food nutrition. National soybean production fell 7.06% from 963.18 tons in 2015 to 887.54 tons in 2016 [1]. The average soybeans consumption per year in Indonesia is 2.2 million tons, but ironically, 67.99% of soybean needs are obtained from imports [2]. Therefore there is a need to increase soybean production in order to meet the needs and overcome import dependence.

Land ~~expansion~~ ~~expansion~~ by utilizing the potential of suboptimal lands such as Alfisol need to increase soybean production. Alfisol is a potential soil for agriculture, but there are still obstacles that need to be considered in its management [3]. Alfisols have acidic pH due to high Al and Fe reduction and low cation exchange capacity (CEC). In addition, Alfisol has low levels of organic matter and low macro nutrient content such as potassium [4].

Potassium (K) is one of the essential nutrients needed in large quantities for plant growth. The K content in the soil is quite high, but plants can only absorb 1-2% of the total K soil [5]. Potassium plays a various role in

physiological processes in plants such as photosynthesis, protein synthesis, and enzyme activator [6]. The K element also helps increasing plant resistance of pests, diseases, and drought stress. Adequate availability and uptake of K not only increases plant growth and yield, but also the quality of crop yields [7].

Application of rock phosphate (RP), zeolite and quail manure is expected to increase the availability and uptake of K and other nutrients, so that soybean production increases. RP contains a group of phosphorus oxides [Ca<sub>3</sub>(PO<sub>4</sub>)<sub>3</sub>F<sub>2</sub>]. The Ca<sup>+</sup> in rock phosphate has a base effect so that the soil pH increases. Zeolite is an aluminosilicate crystal mineral that has the ability as an ion absorber and exchanger and pH buffer [8]. Zeolite ~~have~~ ~~has~~ high CEC and naturally contain K. Zeolite is negatively charged that can absorb K ions and prevent K from washing.

Quail manure has the potential to be used as a source of organic material, because its availability is abundant and not yet widely used. Adding organic manure as a source of organic material can increase total N, available P, Ca, and K, and increase land productivity [9]. Organic manure in the study [10] was able to improve soil fertility and increase nutrient uptake, plant height, plant dry weight, number of pods per plant, and dry weight of soybean. This study was conducted to determine the effect of

RP, zeolite, and quail manure on K uptake and soybean seed yield on Alfisol.

## 2. MATERIALS AND METHODS

### 2.1 Study Area

The study was carried out on Alfisol land in Soko, Sukosari Village, Jumantono Subdistrict, Karanganyar, Surakarta, Indonesia in July-October 2018. The research location was at an altitude of 180 meters above sea level with coordinates 7°37'41"S and 110°57'7"E. Laboratory analysis was carried out at the Laboratory of Chemical and Soil Fertility, Faculty of Agriculture, Universitas Sebelas Maret, Surakarta. The materials used in the study were soybean DEGA 1, rock phosphate, zeolite, and quail manure. The tools used include spectrophotometers, flame photometers, analytical scales, and ovens.

### 2.2 Experimental Design

This study used a single factor Complete Randomized Block Design (RAKL) consisting of 9 treatments with 3 block (replicates), so there were 27 experimental units. Treatments include: P0: Control, P1: RP 2,5 tons/ha + quail manure 5 tons/ha, P2: zeolite 2,5 tons/ha + quail manure 5 tons/ha, P3: RP 2,5 tons/ha + zeolite 2,5 tons/ha + quail manure 5 tons/ha, P4: RP 2,5 tons/ha + quail manure 2,5 tons/ha, P5: zeolite 2,5 tons/ha + quail manure 2,5 tons/ha, P6: RP 5 tons/ha + quail manure 2,5 tons/ha, P7: zeolite 5 tons/ha + quail manure 2,5 tons/ha, P8: RP 5 tons/ha + zeolite 5 tons/ha + quail manure 5 tons/ha.

### 2.3 Crop Management

The experimental field was divided into 3 equal blocks (replicates) with an inter-block spacing of 40 cm. Each block divided into 9 equal plots (1.7 m x 3 m) with interplot spacing of 20 cm, giving a total of 27 lots. Laboratory analysis was conducted to determine the characteristic of RP, zeolite, quail manure, initial soil, and post-harvest soil. Soil samples from each plot were taken before the application of treatment and planting. Application of RP, zeolite, and manure according to predetermined treatment. Soybean seed were planted directly into planting hole with a plant spacing of 25 cm x 25 cm, so there were 84 planting holes. Plant maintenance includes replanting, watering as well as control of weed, pests and diseases. At the flowering stage (4 weeks after planting) K tissue and K uptake were measured.

Soybean harvesting was carried out at 70 days after planting. Post-harvest soil samples were collected from each plot to measure soil pH, organic matter, CEC and available K. Soybean growth and yield variables were observed i.e. plant height, dry matter, tissue K, K uptake, number of filled pods and seed weight per plot.

### 2.4 Statistical Analysis

All the collected data were subjected to analysis of variance (ANOVA) using SPSS statistics package. Significant ANOVA results were tested further using the Duncan Multiple Range Test (DMRT) at 5%.

## 3. RESULT AND DISCUSSION

### 3.1 Initial Soil Condition

Alfisols on the experimental field had low soil fertility with 1.65% organic C (low), 2.85% organic matter (medium) and pH 4.9 (acid). Rao and Reddy [11] state that soil conditions with pH <6 will inhibit soybean growth because soil pH affects solubility and nutrient availability [12]. Soil with low pH will have a relatively low CEC, because more H<sup>+</sup> ions are attached to colloids and cause soil cations to be pushed into the soil solution. This condition, as reported by Sudadi et al. [13] in their study that Alfisols had low CEC (5.7 me/100g) and very low available K (0.33 me/100g). Application of RP, zeolite and quail manure is needed to improve the soil condition in providing and increasing nutrient uptake to support optimal plant growth and yield.

### 3.2 Characteristic of RP, zeolite dan quail manure

Rock phosphate, zeolite and quail manure had a pH of 7.9-8.3 (moderately alkaline) and expected to improve the soil conditions of the experiment field which has low pH. Quail manure has a N content of 1.32%, P 4.54%, K 1.51%, and C-org 17.56% with C / N 13.31 [14]. RP is slow release phosphate source, and manure contains organic acids that can increase the solubility of RP [15]. Zeolite with CEC 128.6 me / 100g meet the Indonesian National Standard of zeolite quality requirements. Chmielewska and Lesný [16] state that zeolite have pores that contain cations and channels that are broad in their structure, so that CEC is important parameter to determining the quality of zeolite.

Tabel 1. Effect of rock phosphate, zeolite and quail manure on soil pH, organic matter, cation exchange capacity and available K

Treatment	pH	Organic matter (%)	Cation Exchange Capacity (me/100g)	Available K (me/100g)
P0	6.4a	2.86a	41.79a	0.87a
P1	6.8b	3.06a	47.23a	1.00a
P2	6.5a	2.67a	48.07a	0.78a
P3	6.8b	3.58a	45.06a	0.82a
P4	6.7b	2.59a	50.18a	0.62a
P5	6.3a	3.20a	45.04a	1.09a
P6	6.9b	3.37a	43.96a	1.10a
P7	6.4a	3.24a	45.36a	1.36a
P8	6.8b	3.17a	40.26a	0.86a

The numbers followed by the same letters show no significant difference in the DMRT 5%

### 3.3 Soil pH

Soil pH is a characteristic that describes the amount of H<sup>+</sup> ion concentration in the soil. Application of RP, zeolite and quail manure significantly increase the soil pH (p=0.00). Combination of RP 2.5 tons/ha + quail manure 5 tons/ha, RP 2.5 tons/ha + zeolite 2.5 tons/ha + quail manure 5 tons/ha, RP 2.5 tons/ha + quail manure 2.5 tons/ha, RP 5 tons/ha + quail manure 2.5 tons/ha and RP+zeolite+manure each 5 tons/ha gave better results than the control treatment, while other treatments showed the same results with the control treatment.

The pH value in various treatments were compared with the initial soil pH (4.9) has increased. The increase in pH occurs due to the RP, zeolite and quail manure given to the soil have a higher pH of 7.9-8.3 (moderately alkaline). Quail manure releases organic compounds in the form of organic acids that are able to bind H<sup>+</sup> ions so that the soil pH increases. Appropriate soil pH will increase nutrient availability, growth and yield of plants [12].

Treatment with the addition of RP gives an increase in pH higher than treatment without RP. RP is slow release and has a Ca content that has an alkaline effect. Acidic initial soil conditions tend to accelerate the solubility of RP [17], which has implications for increasing pH.

### 3.4 Organic Matter

Statistical analysis showed that RP, zeolite and quail manure did not affect soil organic matter. Combination of Rp 2.5 tons/ha + quail manure 2.5 tons/ha has the lowest organic matter content (2.59%). Treatment of RP 2.5 tons/ha + quail manure 5 tons/ha +

zeolite 2.5 tons/ha tends to increase organic matter better than other treatments with organic matter content 3.58%.

RP+zeolite+quail manure each 5 tons/ha only increases the organic matter 11.07%, while the combination of RP 2.5 tons/ha + zeolite 2.5 tons/ha + quail manure 5 tons/ha tends to increase soil organic matter 25.41% compared to treatment control. Zeolite is able to influence soil aggregate stability [18]. Soils with stable aggregates have the ability to infiltrate water and high nutrient retention. Zhang et al. [19] stated that manure also increases soil aggregation which physically protects soil organic matter from degradation by microorganisms and exoenzymes.

### 3.5 Cation Exchange Capacity (CEC)

Cation Exchange Capacity (CEC) is the ability of the soil to bind and exchange cations. Analysis results showed that the application of RP, zeolite and quail manure did not affect the soil CEC. Table 1. shows that overall soil CEC tends to increase compared to initial soil condition.

Cation Exchange Capacity (CEC) is the ability of the soil to bind and exchange cations on its surface. Soils with high CEC are able to absorb and provide nutrients better than low CEC. Figure 1 shows a combination of Rp. 2.5 tons/ha + quail manure 2.5 tons/ha tends to increase CEC 20.08% compared to control treatment. RP is a rock with high content of phosphate (P). Yuniarti et al. [20] reported that the administration of P and organic matter to the soil contributed to adding negative charges to the soil causing CEC to increase.

### 3.6 Available K

Potassium (K) is a macro essential nutrient that highly mobile and is need in large

Formatted: Highlight

Formatted: Highlight

Comment [RD1]: repeated

quantities for plant growth. K availability in the soil determines the amount of K that can be absorbed by plants [21]. Application of RP, zeolite and quail manure did not effect the K availability in the soil (Table 1.). Combination of zeolite 5 tons/ha + quail manure 2.5 tons/ha tended to increase K available 56.18% compared to the control treatment. Manure as a source of organic matter contains humus which can increase the release of K elements and determine the dynamic balance status of

potassium in the soil [22]. Organic matter can increase the ability to hold water so that soil moisture is maintained. Soil under humid conditions will affect the release mechanism of K which is absorbed to become available to plants. Zeolite can increase K availability due to the capability of zeolite to absorb K<sup>+</sup> from soil solution and prevent K<sup>+</sup> from leaching. Zeolite also contain basic cations such as K and Ca, as well as macro elements that can be added to the soil as a plant nutrient [23].

Table 2. Effect of rock phosphate, zeolite and quail manure on K uptake, plant height and dry matter

Treatment	K Uptake (mg/plant)	Plant Height (cm)	Dry Weight (gram)
P0	155.98a	21.33a	4.66a
P1	419.80a	27.95b	12.76a
P2	370.92a	25.97b	8.25a
P3	379.53a	26.97b	9.17a
P4	385.08a	25.87b	9.37a
P5	503.55a	25.41ab	10.36a
P6	505.67a	25.26ab	12.04a
P7	509.83a	25.13ab	11.40a
P8	375.85a	25.07ab	10.22a

The numbers followed by the same letters show no significant difference in the DMRT 5%

### 3.7 K Uptake

Potassium (K) absorbed by plants in the form of exchangeable K and K<sup>+</sup> ions in the soil solution. RP, zeolite, and quail manure application tended to increase K uptake compared to the control treatment which had a K uptake of 155.98 mg/plant. The highest K uptake was found in the treatment of zeolite 5 tons/ha + quail manure 2.5 tons which was 509.83 mg/plant. Zeolite naturally contains K and increases K availability in the soil and prevents K from leaching [24]. As reported by Rabai et al. [25] that zeolite increases the concentration and uptake of K as well as the nutrient use efficiency of K elements by plants.

Treatment with quail manure 2.5 tons/ha tended to give higher K uptake than dose 5 tons/ha. Quail manure contain organic material that capable to realese K nutrients into the soil solution. However, excessive application of manure can result in leaching up soil cations such as K, Ca and Mg which cause decreased of CEC and soil pH [26], those condition can inhibit nutrient absorption by plant roots

### 3.8 Plant Height

Plant height is one of the plant growth indicator and its irreversible. Table 2. showed

that RP, zeolite and quail manure were significantly increase the plant height ( $p=0,006$ ). The RP 2.5 tons/ha + quail manure 5 tons/ha; zeolite 2.5 tons/ha + quail manure 5 tons/ha; RP 2.5 tons/ha + zeolite 2.5 tons/ ha + quail manure 5 tons/ha; and RP 2.5 tons/ha + quail manure 2.5 tons/ ha significantly increases plant height. Other treatments have similar effect with control treatment.

Plant height in each treatment has increased every week, and application of RP, zeolite and quail manure was able to increase plant height 21.29-31% compared to control treatment. RP contains P elements which play a significant role in the formation of root nodules in legume plants. Zeolite can increase the water holding capacity due to its physical properties, so it can improve soil structure and increase macro and micro pores in the soil [27]. Manure as organic material supply plant nutrients that stimulate root growth [28]. A proper root system will promote nutrient absorption and optimal plant growth.


### 3.9 Dry Matter

Dry weight determined the amount of assimilates that translocated to plant organs as the result of metabolic processes in the form of photosynthesis, nutrient absorption


and water. The amount of assimilate produced during growth is related to the rate of photosynthesis [29]. The control treatment had the lowest dry weight of 4.66gram. Combination of RP 2.5 tons/ha + quail manure 5 tons/ha tended to give the highest result compared to other treatments. Wang et al. [30] stated that potassium plays an important role in water retention in plant tissues, as well as cell turgor during the opening and closing of stomata. Potassium in the mechanism of opening and closing of stomata related to

plant physiology processes, especially CO<sub>2</sub> fixation which will produce assimilates to meet plant needs for proper growth.

Dry weight was positively correlated with K uptake ( $r = 0.808$ ;  $p = 0,000$ ). Accumulation of photosynthate during plant growth related to the rate of photosynthesis, and potassium plays an important role in these process. Wang et al. [30] stated that adequate of K availability and K uptake will increase the total accumulation of plant dry weight.



The numbers followed by the same letters show no significant difference in the DMRT 0.05  
Figure 1. Effect of rock phosphate, zeolite and quail manure on number of filled pods



The numbers followed by the same letters show no significant difference in the DMRT 0.05  
Figure 2. Effect of rock phosphate, zeolite and quail manure on seed weight per plot

### 3.10 Number of Seed per Pods

Number of seed per pods was significantly increase due to the application of RP, zeolite and quail manure. The treatments with significant effect were RP 2.5 tons/ ha + quail manure 5 tons/ha; RP 5 tons/ha + quail manure 2.5 tons/ha; and RP 5 tons/ha + zeolite 5 tons/ha + quail manure 5 tons/ha. The increase in the number of pods of these treatments were 80.41-115.46% compared to the control treatment. Other treatments showed similar result, except the zeolite 2.5 tons/ha + quail manure 5 tons/ha which showed the same effect as the control treatment.

RP as a source of phosphate (P) was increases the soil pH (table 1.) and its also effect the availability of P in the soil. Legumes use P for the formation of pods and seed filling during the generative phase. The absorbed P is translocated from the roots and leaves to the seed [31]. The P and K elements are needed in hight amount for proper plant growth and development. Fageria and Oliveira [32] found that the combination of both P and K gave a large increase in plant production compared to the effects of each individual element. These interactions indicate a synergy between P and K [33].

The number of filled pods correlated with dry r weight ( $r = 0.835$ ;  $p = 0,000$ ). Sabilu et al.

[34] stated that photosynthate in the generative phase widely used for the process of pods formation and seed filling.

### 3.11 Seed Weight per Plot

The combination of RP 2.5 tons/ha + quail manure 5 tons/ha and RP 5 tons/ha + zeolite 5 tons/ha + quail manure 5 tons/ha significantly increases the seed weight per plot. Other treatments showed the similar effect with the control treatment. Seed weight per plot correlates with the number of filled pods ( $r = 0.921$ ,  $p = 0,000$ ). Manure does not only contain macro and micro nutrients needed by plants, but also produces derivative products in the form of humic acid [35]. Mindari et al. [36] mentioned that humic acid can increase K uptake so that it can trigger ion transport and assimilation of carbohydrates and affected the increase of plant yield.

## 4. CONCLUSION

Rock phosphate, zeolite and quail manure is needed to increase K uptake and soybean yield in Alfisol. RP, zeolite and quail manure tend to increase potassium uptake. Potassium plays an important role in photosynthesis and translocation of photosynthate to support plant growth and production. The combination of RP 2.5 tons/ha and quail manure 5 tons/ha gave the highest yield on soybean yield variables compared to other treatments, and significantly increased the number of filled pods and seed weight per plot.

## REFERENCES

1. BPS [Biro Pusat Statistik]. 2016. Data Produksi Kedelai Nasional. Badan pusat statistik republik indonesia. Jakarta.
2. KEMENTAN [Kementerian Pertanian]. 2016. Outlook komoditas pertanian tanaman pangan: kedelai. Kementerian Pertanian Republik Indonesia. Jakarta
3. Prasetyo A, Listyorini E, Utomo WH. 2014. Hubungan sifat fisik tanah, perakaran dan hasil ubi kayu tahun kedua pada Alfisol Jatikerto akibat pemberian pupuk organik dan anorganik. J Tanah dan Sumberdaya Lahan 1(1): 27-37.
4. Chinnadurai C, Gopalaswamy G, Balachandar D. 2014. Impact of long-term organic and inorganic nutrient managements on the biological properties and eubacterial community diversity of the Indian semi-arid Alfisol. J Archives of Agronomy and Soil Science, 60(4): 531-548.
5. Munawar A. 2011. Kesuburan tanah dan nutrisi tanaman. IPB Press: Bogor
6. Amtmann A, Rubio F. 2012. Potassium in Plants. In: eLS. John Wiley & Sons Ltd, Chichester. <http://www.els.net>. doi:[10.1002/9780470015902.a0023737](https://doi.org/10.1002/9780470015902.a0023737)
7. Zörb C, Senbayram M, Peiter E. 2014. Potassium in agriculture: status and perspectives. J Plant Physiol. 171(9): 656-69. doi: [10.1016/j.jplph.2013.08.008](https://doi.org/10.1016/j.jplph.2013.08.008).
8. Sabilu Y. 2016. Aplikasi zeolit meningkatkan hasil tanaman pada tanah ultisol. J Biowallacea 3(2): 396-407.
9. Akter F, Rahman M, Alam A. 2019. Soil chemical properties as influenced by long term manuring and nitrogen fertilization in Bangladesh. Asian Journal of Soil Science and Plant Nutrition. 4(4): 1-9. doi:[10.9734/AJSSPN/2019/v4i430049](https://doi.org/10.9734/AJSSPN/2019/v4i430049)
10. Sudarsono WA, Melati M, Aziz SA. 2013. Pertumbuhan, serapan hara dan hasil kedelai organik melalui aplikasi pupuk kandang sapi. J Agron. Indonesia 41 (3) : 202 – 208
11. Rao AS, Reddy KS. 2010. Nutrient management in soybean. In: Singh G (ed) The soybean: botany, production and uses. London: CABI.
12. Zhao J, Dong Y, Xie X, et al. 2011. Effect of annual variation in soil pH on available soil nutrients in pear orchards. Acta Ecologica Sinica, 31(4): 212–216. doi:[10.1016/j.chnaes.2011.04.001](https://doi.org/10.1016/j.chnaes.2011.04.001)
13. Sudadi, Sumarno, Handi W, 2012. Pengaruh pupuk organik berbasis azolla, fosfat alam dan abu sekam terhadap hasil padi dan sifat kimia tanah alfisol. Sains Tanah 11(2): 77-84. doi: [10.15608/stjssa.v11i2.223](https://doi.org/10.15608/stjssa.v11i2.223)
14. Alvernia P, Minardi S, Suntoro. 2017. Zeolite and organic fertilizer application to the improvement of available P and soybean (*Glycine max* L) seed yield in Alfisols. J of Soil Science and Agroclimatology 14 (2): 84-90.
15. Ditta A, Muhammad J, Imtiaz M et al. 2018. Application of rock phosphate enriched composts increases nodulation, growth and yield of chickpea. International Journal of Recycling of Organic Waste in Agriculture 7 :33–40. doi.org/[10.1007/s40093-017-0187-1](https://doi.org/10.1007/s40093-017-0187-1)
16. Chmielewska E, Lesný J. 2012. Selective ion exchange onto Slovakian natural zeolite in aqueous solutions. Journal of Radioanalytical and Nuclear Chemistry, 293(2): 535–543.
17. Wahba MM, Bahna FL, Amal MA. 2018. Improving the availability of phosphorus from rock phosphate in calcareous soils by

- natural materials. *Bioscience Research* 15(3):1796-1804.
18. Aminiyan MM, Safari AA, Sheklabadi SM. 2015. Aggregation stability and organic carbon fraction in a soil amended with some plant residues, nanozeolite, and natural zeolite. *Int J Recycl Org Waste Agricult* (2015) 4:11–22. doi: [10.1007/s40093-014-0080-0](https://doi.org/10.1007/s40093-014-0080-0)
  19. Zhang H, Ding W, Yu H, et al. 2015. Linking organic carbon accumulation to microbial community dynamics in a sandy loam soil: result of 20 years compost and inorganic fertilizers repeated application experiment *Biol Fertil Soils* 51(2): 137. doi.org/[10.1007/s00374-014-0957-0](https://doi.org/10.1007/s00374-014-0957-0)
  20. Yuniarti A, Arifin M, Sofyan ET, et al. 2018. The effect of Sinabung volcanic ash and rock phosphate nanoparticle on CEC (cation exchange capacity) base saturation exchange (K, Na, Ca, Mg) and base saturation at Andisol soils Ciater, West Java. AIP Conference Proceedings. doi.org/[10.1063/1.5021196](https://doi.org/10.1063/1.5021196)
  21. Káš M, Mühlbachová G, Kusá H, et al. 2016. Soil phosphorus and potassium availability in long-term field experiments with organic and mineral fertilization. *J. Plant Soil Environ.* 62(12): 558–565. doi: [10.17221/534/2016-PSE](https://doi.org/10.17221/534/2016-PSE)
  22. He W dan Chen F. 2013. Evaluating status change of soil potassium from path model. *PLoS ONE* 8(10):1-15. doi: [10.1371/journal.pone.0076712](https://doi.org/10.1371/journal.pone.0076712)
  23. Abdillah A, Syamsiyah J, Riyanto D et al. 2011. Pengaruh pupuk zeolit dan kalium terhadap ketersediaan dan serapan K di lahan berpasir pantai Kulonprogo, Yogyakarta. *Bonorowo Wetlands* 1 (1): 1-7.
  24. Moraetisa D, Papagiannidou S, Pratikakis A. 2015. Effect of zeolite application on potassium release in sandy soils amended with municipal compost. *Desalination and Water Treatment* 57(28):1-12. doi: [10.1080/19443994.2015.1065440](https://doi.org/10.1080/19443994.2015.1065440)
  25. Rabai KA, Ahmed OH, Kasim S. 2013. Use of formulated nitrogen, phosphorus, and potassium compound fertilizer using clinoptilolite zeolite in maize (*Zea mays* L.) cultivation. *Emir. J. Food Agric.* 25 (9): 713-722. doi: [10.9755/ejfa.v25i9.14541](https://doi.org/10.9755/ejfa.v25i9.14541)
  26. Han SH, Ji YN, Hwang J, Kim SB, Park BB. 2016. The effects of organic manure and chemical fertilizer on the growth and nutrient concentrations of yellow poplar (*Liriodendron tulipifera* Lin.) in a nursery system. *J Forest Science and Technology.* 12(3): 137-143. doi: [10.1080/21580103.2015.1135827](https://doi.org/10.1080/21580103.2015.1135827)
  27. Juarsah I. 2016. Pemanfaatan zeolit dan dolomit sebagai pembenah untuk meningkatkan efisiensi pemupukan pada lahan sawah. *J Agro* 3(1):10-19
  28. Ditta A, Arshad M, Zahir ZA et al. 2015. Comparative efficacy of rock phosphate enriched organic fertilizer vs. Mineral phosphatic fertilizer for nodulation, growth and yield of lentil. *Inter J Agric Bio.* 17 (3): 589-595. doi: [10.17957/IJAB/17.3.14.954](https://doi.org/10.17957/IJAB/17.3.14.954).
  29. Wahono E, Izzati M, Parman S. 2018. Interaksi antara tingkat ketersediaan air dan varietas terhadap kandungan prolin serta pertumbuhan tanaman kedelai (*Glycine max* L. Merr). *Buletin Anatomi dan Fisiologi* 3(1): 11-19
  30. Wang M, Zheng Q, Shen Q, et al. 2013. The critical role of potassium in plant stress response. *Int. J. Mol. Sci.* 14: 7370-7390; doi:10.3390/ijms14047370
  31. Taliman NA, Dong Q, Echigo K, et al. 2019. Effect of phosphorus fertilization on the growth, photosynthesis, nitrogen fixation, mineral accumulation, seed yield, and seed quality of a soybean low-phytate line. *Plants* 2019, 8:1-13. doi: [10.3390/plants8050119](https://doi.org/10.3390/plants8050119)
  32. Fageria NK, Oliveira JP. 2014. Nitrogen, phosphorus and potassium interactions in upland rice. *Journal of Plant Nutrition* 37 (10):1586–600
  33. Rietra RPJJ, Heinen M, Dimkpa CO, et al. 2017. Effects of nutrient antagonism and synergism on yield and fertilizer use efficiency. *Communications in Soil Science and Plant Analysis*, 48(16): 1895-1920, doi: [10.1080/00103624.2017.1407429](https://doi.org/10.1080/00103624.2017.1407429)
  34. Sabilu Y, Damhuri, Imran. 2015. Kadar N, P, dan K kedelai (*Glycine max* (L) Merrill) yang diaplikasi *Azotobacter* sp., mikoriza dan pupuk organik. *Biowallacea*, 2 (1): 153-161
  35. Murmu K, Swain DK, Ghosh BC. 2013. Comparative assessment of conventional and organic nutrient management on crop growth and yield and soil fertility in tomato-sweet corn production system. *Aust J Crop Sci.* 7(11):1617-1626.
  36. Mindari W, Aini N, Kusuma Z, et al. 2014. Effects of humic acid-based buffer + cation on chemical characteristics of saline soils and maize growth. *Journal Of Degraded And Mining Lands Management.* 2(1): 259-268. doi: [10.15243/jdmlm.2014.021.259](https://doi.org/10.15243/jdmlm.2014.021.259)