

Economic assessment of Napier grass production using different fertiliser combinations under smallholder farming conditions in the Central Highlands of Kenya

ABSTRACT

Aims: The objective of the study was to evaluate the cost effectiveness of using different fertiliser combinations to improve Napier grass Production within the smallholder farms.

Study design: The experimental design was a Randomised Complete Block Design (RCBD) with three replicates per treatment. The treatments were: Di-Ammonium Phosphate (DAP), rabbit (*Oryctolagus cuniculus*) manure; rabbit manure plus rabbit urine, DAP plus Calcium Ammonium Nitrate (CAN), DAP plus rabbit urine, Control and Conventional method.

Place and Duration of Study: The study was done in Embu County, Kenya from March 2015 to January, 2016.

Methodology: The economic analysis to determine the most cost-effective fertiliser was done using gross margins and cost-benefit ratios approach. The economic analysis to determine the most cost-effective fertiliser was done using gross margins and cost-benefit ratios approach.

Results: Rabbit manure plus urine had the highest cost of production averages at US\$.154 8.13 per year at $p < 0.05$ while the conventional method was US\$ 494.59 at $p < 0.05$. The study revealed that the most cost-effective fertiliser in Embu County was DAP plus rabbit urine treatment under "Tumbukiza" pits.

Conclusion: The projections are that by the end of the second cropping year, the treatment top-dressed with either rabbit urine or CAN would be having higher gross margins since the initial cost would have been recovered. Farmers in Embu County are encouraged to integrate the use of both organic and inorganic fertilisers to achieve high production in a cost-effective way.

Keywords: Cost, Fodder, inputs, profitability

1. INTRODUCTION

The dairy industry is an integral sub-sector of livestock production in Kenya, which supports the key players within the entire value chain [1]. Total annual milk production in Kenya is has been approximated at 3.43 billion litres, of which more than 80% is from the smallholder farms [2]. Currently, the milk production per cow per day is averaged at 6 Kgs, which is lay way below the expected 15 Kgs [3]. Dairy production performance in most smallholdings is are below optimal due to some factors associated with dairy production systems. These factors comprise of low quality feeds, poor feeding, a declining genetic base, animal diseases, poor access to credit facilities, effects of climate change and diminishing land [4, 5, 6].

To realise milk from a lactating cow, the animal genetic base and environment are critical. The environment consists of housing and Feeding of which feeding stands at 70% of the production cost. Studies have been done on improving milk production, but the yields have

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

30 | remained low with the milk unit cost being comparatively high, which **makes made** it
31 | unaffordable to most consumers [7, 8]. A research done in Embu County [3] showed that the
32 | average cost of producing a litre of milk was US\$ 0.374. Further studies indicated that the
33 | highest percentage of the cost of producing milk **is** from fodder constituting 55-70% [5, 9].

Formatted: Highlight

Formatted: Highlight

34 |
35 | Napier grass **is has been?**the most popular perennial fodder used within the smallholder
36 | crop-livestock farming systems in Kenya, where over 80% of the national milk is produced
37 | [10]. The reason for these is because of its advantageous traits such as vigorous growth,
38 | high biomass productivity, deep root system for drought tolerance, a wide range of soil
39 | conditions, high photosynthetic and its water-use efficiency [11]. Napier grass acts as a
40 | windbreak in crop fields and stabilises the soil by holding particles together in this manner,
41 | preventing soil erosion [12]. Milk production in smallholdings could be increased by reducing
42 | the cost of production, especially for fodder. There **is are** limited empirical data on the
43 | economic assessment of Napier production to achieve high production. Hence we evaluated
44 | the Economic assessment of using different fertiliser combinations to improve Napier grass
45 | Production within smallholder farming conditions.

Formatted: Highlight


Formatted: Highlight

46 | 47 | **2.0 MATERIAL AND METHODS / EXPERIMENTAL DETAILS / METHODOLOGY** 48 | **NO NEED TO MENTION ALL!**

Formatted: Highlight

49 | 50 | **2.1 study site**

51 | The study was done in Kirigi (0°24'14.71" S, 37°32'10.6" E), Embu County, Eastern Kenya.
52 | Kirigi is located in Agro-Ecological Zone (AEZ) UM1 (Upper midland zone 1), a coffee-tea
53 | zone and lies at an altitude of 1650 m above sea level. The average temperature is 18.7°C,
54 | and the precipitation pattern is bimodal with an annual average rainfall of 1677 mm [13]. The
55 | daily rainfall pattern and amounts experienced during the study period is shown in Figure 1.



56 |
57 | Figure 1: Daily rainfall during the study period.

58 | 59 | **2.2 Experimental design**

60 | The field trial **was were** laid in a randomised complete block design replicated thrice. The test
61 | crop was Napier grass, Kakamega 1 variety. The treatments were: Di-Ammonium Phosphate
62 | (DAP), rabbit manure, rabbit manure plus urine, DAP plus Calcium Ammonium Nitrate (CAN),
63 | DAP plus rabbit urine, conventional method and Control (no fertiliser input). The treatments
64 | were assigned randomly within the three replicates, and the blocking was done based on

Formatted: Highlight

slope and soil homogeneity as the major influencing factors. The fertiliser application rate was based on N nutrient at 45kg of N ha⁻¹ from the assorted sources: DAP, CAN, rabbit manure, and rabbit urine. The plot size measured 3m by 2.1 m consisting of five "Tumbukiza" pits measuring 0.9m length by 0.6m width by 0.6m depth. On the other hand, the conventional method pits measured 0.2 m length by 0.15 m width by 0.2 m depth ([how much fertilizer and what type of fertilizer are applied under conventional?](#)). Five cuttings of Napier grass were planted in each "Tumbukiza" pit while one cutting was planted in the conventional method pit. [How many napier cutting were planted in the other treatments?](#)
[How were the fertilizer applied, interval of application](#)

2.3 Data collection

The economic analysis to determine the most cost-effective fertiliser was done using gross margins and cost-benefit ratios approach. The gross margin (GM) was calculated by subtracting total variable cost (TVC) from total revenue (TR) of Napier production per hectare (equation 1).

$$GM = TR - TVC \quad \text{Equation 1}$$

Where: GM is gross margin (US\$/ha), TR is total revenue or the total value of output from the Napier Production (US\$/ha). It is the product of average output per hectare multiplied by the market price, and TVC is total variable cost or the costs that are specific in producing Napier (US\$/ha). TVC varies according to output and is incurred on variable inputs. This includes the cost of inputs like canes, fertiliser, and hired/family labour per treatment.

Formatted: Highlight

Formatted: Highlight

2.4 Data analysis

Data were subjected to analysis of variance (ANOVA) using SAS 9.2. Mean separation was done using Tukey's Kramer Honest significant difference (HSD) at $P = 0.05$. Differences between means were considered significant if P values were less than 0.05. Data were analysed using SAS edition 9.2.

$$Y_{ijk} = \mu + B_i + T_j + E_{ijk} \quad \text{Equation 2}$$

Where: Y_{ijk} is the dependent variable, μ is the mean, B_i is the effect due to i^{th} replication, T_j is the effect due to j^{th} treatment and E_{ijk} is the residual effect.

3.0 RESULTS AND DISCUSSION

3.1 cost of production

During the study, it was observed that all means were significantly different from the control in the 1st crop while DAP and rabbit manure were not significantly different from the control in the 2nd, 3rd and 4th crops. The highest costs incurred were observed in the 1st crop while during the other crops the costs were almost constant. The conventional method had the lowest cost of production while rabbit manure plus urine had the highest cost.

109 **Table 1: Analysis of the cost of production using different fertiliser combinations on**
 110 **Napier grass in Embu County**

Treatment	Production costs (US\$) <u>per plot or ha/?</u>			
	1 st crop	2 nd crop	3 rd crop	4 th crop
DAP	786.47 ^d	92.74 ^c	92.74 ^c	92.74 ^c
Rabbit manure	1178.92 ^b	92.74 ^c	92.74 ^c	92.74 ^c
Rabbit manure+Urine	1201.99 ^a	115.81 ^b	115.81 ^b	115.81 ^b
DAP+CAN	817.31 ^c	123.86 ^a	123.58 ^a	123.58 ^a
DAP+Rabbit urine	809.54 ^b	115.81 ^b	115.81 ^b	115.81 ^b
Control	717.17 ^e	92.74 ^c	92.74 ^c	92.74 ^c
Conventional method	259.33 ^f	78.42 ^d	78.42 ^d	78.42 ^d
P	<.0001	<.0001	<.0001	<.0001

111 Means in the same column followed by the same letter are not significantly different at
 112 $P < 0.05$
 113

114 The study showed **s** that the highest cost was incurred during the 1st crop since planting
 115 material, fertilisers and more labour were used due to the land preparation. In the 2nd, 3rd and
 116 4th crop, more cost was incurred where topdressing was done since there was the cost of
 117 fertiliser and extra labour for the fertiliser application. On the other hand, the conventional
 118 method was cheaper to establish since it used less labour to establish. The study found that
 119 the labour cost was the highest with estimated at 52% of the production cost. This result is in
 120 agreement with [5] who found that labour cost forms a large proportion in the dairy
 121 smallholder farms. Despite the fact that Rabbit manure plus urine had the highest cost of
 122 production, its gross margins were higher compared to the conventional method, which had
 123 the lowest gross margins.

Formatted: Highlight

124 **3.2 Cost-benefit analysis**

125 The study found that during the 1st crop, all the Gross margins were negative with the
 126 conventional method having the lowest gross margin though, in the 2nd, 3rd and 4th crop
 127 positive gross margins were realised. All the treatments means were significantly different
 128 from the control throughout all crops apart from the conventional method, which was not
 129 significantly different from control apart from the 1st crop. The study on the economic
 130 evaluation of the most cost-effective fertiliser in Embu County revealed that DAP plus rabbit
 131 urine treatment under "Tumbukiza" pits was leading, followed closely by rabbit manure plus
 132 urine.
 133

134 **Table 2: Assessment of the cost-effectiveness of using different fertiliser**
 135 **combinations on Napier grass in Embu County**

Treatment	Gross Margins (US\$) <u>to produce for how much?/land size</u> <u>or kg of Napier?</u>			
	1 st crop	2 nd crop	3 rd crop	4 th crop
DAP	-382.68 ^b	129.77 ^{bc}	224.93 ^{cdde}	4663.97 ^a
Rabbit manure	-948.01 ^e	280.48 ^a	377.77 ^{ab}	508.60 ^a
Rabbit manure+Urine	-793.43 ^d	314.92 ^a	441.00 ^b	654.00 ^a
DAP+CAN	-585.80 ^d	205.03 ^b	252.37 ^{bcd}	613.93 ^{ab}
DAP+Rabbit urine	-445.67 ^b	312.97 ^a	662.00 ^a	803.31 ^a
Control	-624.43 ^c	1.26 ^d	34.64 ^{de}	34.96 ^b
Conventional method	-177.15 ^a	9.39 ^d	72.50 ^{de}	22.90 ^b
LSD	118.84	82.19	211.70	355.01
P	<.0001	<.0001	0.001	0.007

136 Means in the same column followed by the same letter are not significantly different at
137 $P < 0.05$

138

139 The study on the economic evaluation of the most cost-effective fertiliser in Embu County
140 revealed that DAP and Rabbit urine combinations were leading, followed closely by Rabbit
141 manure and rabbit urine combinations all under “*Tumbukiza*” plots. Both treatments realised
142 high yields in all the harvests. The reason why the first was leading compared to the latter
143 was that the first had less labour and time for fertiliser application, unlike the manure that had
144 more time and labour. The control and Conventional method had low gross margins in all the
145 harvests due to their low yields and high cost involved in their establishment. Gross margins
146 from treatments with “*Tumbukiza*” plots had high gross Margin apart from the control despite
147 their high cost of establishment particularly digging the holes compared to the conventional
148 method. The results differed with a study was done by [14] who found the gross margins for
149 the “*Tumbukiza*”, and Conventional method was similar.

150

151 4. CONCLUSION

152

153 The study revealed that the most cost-effective fertiliser in Embu County was DAP plus rabbit
154 urine under “*Tumbukiza*” pit treatment since it performed better compared to the others. The
155 reason as to why the treatment was doing well **is was** because it used less labour and time
156 for fertiliser application, unlike where manure was used since there were more time and
157 labour involved. The projections are that by the end of the second cropping year, the
158 treatment top-dressed with either rabbit urine or CAN would be having higher gross margins
159 since the initial cost would have been recovered. Farmers in Embu County are encouraged to
160 integrate the use of both organic and inorganic fertilisers to achieve high production in a cost-
161 effective way.

162

163

164 COMPETING INTERESTS

165

166 We have no conflicts of interest to disclose.

167

168

169 REFERENCES

170

171

1. Muriuki HG. Dairy Development in Kenya. FAO Report. 2011. Pp 3.

172

173

174

2. Kenya Dairy Board. Status and Outlook of the dairy industry in Kenya. Proceedings of the
14th Africa dairy conference and exhibition, 22nd to 24th August 2018. 2018. Nairobi,
Kenya.

175

176

177

3. Mugambi DK., Maina MW, Kairu S, Gitunu, AMM. Assessment of performance of
smallholder dairy farms in Kenya: an econometric approach. Journal of Applied
Biosciences. 2015.85:7891– 7899.

178

179

180

181

4. Jayne TF, Milu M. Land constraint in Kenya's densely populated rural areas:
implementation for food policy and institutions reforms. Proceedings of the 86th annual
conference of agricultural economics society on 16th to 18th April, 2012. 2012. University
OF Warwick, United Kingdom.

Formatted: Highlight

- 182 5. Kibiego MB, Lagat JK, Bebe BO. Competitive of Smallholder Milk Production Systems in
183 Uasin Gishu of Kenya. Journal of Economics and Sustainable Development. 2015.6 (10):
184 39-45.
- 185 6. Mutavi SK, Kanui TI, Njarui DM, Musimba NR, Amwata DA. Innovativeness and
186 Adaptations: The way forward for Small scale Peri-Urban Dairy Farmers in semi-arid
187 regions of South Eastern Kenya. International Journal of scientific research and
188 innovative technology. 2016. 3(5): 1-14.
- 189 7. Omiti J, Wanyoike F, Staal S, Delgado C, Njoroge L. Will small-scale dairy producers in
190 Kenya disappear due to economies of scale in production? Contributed paper prepared
191 for presentation at the International Association of Agricultural Economists Conference.
192 2006. Gold Coast, Australia.
- 193 8. Staal SJ, Alejandro NP, Jabber M. Dairy development for the resource poor Part 2:
194 Kenya and Ethiopia dairy development case studies pro-poor livestock policy initiative.
195 2008.
- 196 9. Hussain M, Ghafoor A, Saboor A. Factors affecting milk production in buffaloes: A case
197 study. Pakistan Vet. J. 2010.30(2):115-117
- 198 10. Njarui DMG, Gatheru M, Wambua JM, Nguluu SN, Mwangi DM, Keya GA. Feeding
199 management for dairy cattle in smallholder farming systems of semi-arid tropical Kenya.
200 Livestock Research for Rural Development. 2011.23(111).
- 201 11. Tessema Z. Effect of plant density on morphological characteristics, yield and chemical
202 composition of Napier grass (*Pennisetum purpureum* (L.) Schumach). East African
203 Journal of Sciences. 2008. 2:55–61.
- 204 12. Khan Z, Pickett P, Midega C, Jimmy PJ. Climate-smart push-pull: A conservation
205 agriculture technology for food security and environmental sustainability in Africa. First
206 Africa Congress on Conservation Agriculture Lusaka, Zambia, 18th to 21st March 2014.
207 2014.
- 208 13. Jaetzold R, Schmidt H, Hornetz B, Shisanya C. Farm management hard book; Natural
209 conditions and farm management information 2. Ministry of Agriculture & GTZ, Nairobi,
210 Kenya. 2006. Pp 23.
- 211 14. Nyambati EM, Lusweti CM, Muyekho FN, Mureithi JG. Up-scaling Napier grass
212 (*Pennisetum purpureum* Schum) production using “*Tumbukiza*” method in smallholder
213 farming systems on north-western Kenya. Journal of Agricultural Extension and Rural
214 Development. 2011.3:1-7.