

1 **Consequence of ~~Ground Water~~ Groundwater Irrigation on Physico-Chemical**  
2 **Properties of Soils of Kanholibara Village in Nagpur District, Maharashtra.**

3  
4 **ABSTRACT**

5 A field experiment entitled “Effect of Ground-water irrigation on various soil properties of  
6 Kanholi-Bara in Nagpur District Was conducted during Kharif season of 2015-2016 at Kanholi-  
7 Bara of Hingana tahsil in Nagpur District. The soil samples were collected from that area  
8 comprises two sources of irrigations viz well water and bore well water and soybean crop which  
9 were taken in these fields. The mean value of pH 7.79 was recorded with ground-water irrigation  
10 and EC in ground-water irrigated soil 0.81 dS m<sup>-1</sup>.The organic carbon contains in ~~ground-water~~  
11 ~~groundwater~~ irrigated soil with high SAR and RSC was lower by 35.19 per-cent less than mean  
12 value of organic carbon. The lowest available nitrogen, phosphorus, and potassium status were  
13 185.44, 15.65 and 178.80 kg ha<sup>-1</sup> respectively were obtained with the application of ground-water  
14 of high RSC and SAR. The highest accumulation of heavy metal in soil 1.16 mg kg<sup>-1</sup> lead, 1.30  
15 mg kg<sup>-1</sup> Cobalt, 1.19 mg Kg<sup>-1</sup>Nikel and 0.037 mg kg<sup>-1</sup>Cadmium were present in soil with the  
16 application of high SAR and RSC irrigation water. Due to continuous and injudicious irrigation  
17 with poor quality ~~ground-water-groundwater~~ adversely affect the physical and chemical properties  
18 of soil-.

19  
20 **Key-words:** SAR, RSC, ~~Ground~~water, Physicho-Chemical

21  
22 **INTRODUCTION**

23  
24 Water is one of the limiting factors for agricultural development in developing countries in  
25 order to meet the growing demand of the increasing population. Water is one of the ~~precious~~  
26 ~~valuable~~ sources of nature and required for crop production, ~~whereas an invaluable basic natural~~  
27 ~~resource for human being it sustains life on the earth, w~~Water is being used for several purposes  
28 viz. drinking, Irrigation, hydro-electric production, Industries, transport, sanitation, recreation etc.  
29 However, the ~~largest-most abundant~~ use is for Irrigation which enables to increase the farm

30 | production. A part of rain-water percolates in the ground through joints and cracks in the rocks  
31 | and is known as ground-water. It is a major-significant source of Irrigation in India. About 60% of  
32 | the Irrigated land in the country depends on groundwater (Shah *et al.*, 2000). In Maharashtra the  
33 | total geographical area of 30.77 Mha spread over in thirty-one districts out of which about 59.2  
34 | %( 18.31 M ha) area is under crops and 20.8% under forests. The irrigated area is 11.4% of the  
35 | gross cropped area in the state.

36 | Irrigation is practiced in those parts of the world where rainfall is not sufficient to support  
37 | crop growth or where the rain does not fall when the plants need water. The objective of irrigation  
38 | is to supply plants with water, as needed, to increase yields. An irrigation project should take  
39 | water use efficiency into account as well as the economy involved. When applying irrigation  
40 | water, excesses as well as shortages, as well as shortages, should be avoided (Varallyay, 1977).

41 | The history of irrigated agriculture has shown that irrigation can cause severe deterioration  
42 | of soil productivity. Many early civilizations, whose rise was supported by the productivity of  
43 | irrigated agriculture, were thought to fall as a result of problems caused by irrigation (Gulhati and  
44 | Smith, 1967).

45 | The most common reasons for the failure of irrigation projects are associated with  
46 | waterlogging, salinization, and alkalization. These problems appear gradually and are influenced  
47 | by the quality and quantity of irrigation water, condition of the irrigated land, and other soil  
48 | environmental factors. As the problems develop, they may be recognized by the failure of the  
49 | irrigated land to maintain high yields. If allowed to continue, the problems may become so severe  
50 | that the irrigated land will no longer be productive (Kovda, 1973).

51 | Salt problems and waterlogging are caused by lack of adequate drainage, poor quality water, or  
52 | improper management practices or any combination of the three. These problems, in addition to  
53 | damaging plants, cause deterioration of some desirable soil properties

54 | In India, ground-water-groundwater is major threat to irrigated agriculture. The  
55 | problem has been increasing gradually with the expansion of irrigation facilities associated with  
56 | the faulty water use management. This problem is appearing at a fast rate in deep black and  
57 | medium black soils of the central peninsular India.

58 | The problems of soil salinity/sodicity and water logging in canal, bore well  
59 | irrigated Vertisols are of severe and deserve an immediate attention for corrective measures,  
60 | therefore monitoring of salinity/sodicity in such irrigation areas is essential and hence the detail

Comment [RD1]: Citation required

Comment [RD2]: Citation required

61 knowledge of properties of dry land soils and their inter-relationship will be useful for managing  
62 salt affected soils of similar areas elsewhere in the state for sustainable crop production. It was felt  
63 necessary to take up the study on the assessment of soil salinity and sodicity status in the  
64 particular area. The ~~location-location~~-specific information based on detailed characterization of  
65 kanholibara villege area soils under irrigation is very much essential for judicious management of  
66 such soils. The irrigation induced salinity/sodicity problem in black clay soil has not been  
67 extensively studied although some attempts have been made and the information is scattered.

**Comment [RD3]:** This sentence is very long. To improve readability, consider breaking this into multiple sentences

**Comment [RD4]:** Concluding remark of introduction is missing

68

## 69 MATERIALS AND METHODS

### 70 *Study area*

71 Geographically, Kanholi-Bara is situated at the latitude 20.93 and longitude 78.84  
72 at the elevation of 321.26 m above sea level and lies under sub-tropical zone, covering an net  
73 cropped area of about 704.71 ha<sup>-1</sup> in Hingana Tehsil of Nagpur district, Maharashtra. Kanholi  
74 Bara is characterized by hot and dry summer and fairly cold winter. This area shows wide diurnal  
75 fluctuation in temperature. The maximum and minimum temperature ranged from 26.9<sup>0</sup>C to  
76 43.0<sup>0</sup>C and 13.7<sup>0</sup>C to 26.6<sup>0</sup>C, respectively, whereas the relative humidity varied from 20 to 72 per  
77 cent during the crop growth period, mean annual rainfall is about 1,566.3 mm.

78 Surface soil samples were collected with free survey where there is a differentiation in  
79 soil. A composite soil sample (0-15 cm) depth, after harvesting of soybean crop was taken from  
80 the different spots of the experimental area were taken. These samples were stored in polythene  
81 bags then processed and labeled as per the need for laboratory analysis. Physical properties of the  
82 soils, such as particle size distribution were determined by the international pipette method (Klute  
83 and Dirksen, 1986). The bulk density was determined by core method described by Yaalon  
84 (1957). The hydraulic conductivity was measured by constant head method described by Richards  
85 (1954). Chemical properties like pH and EC of the soil suspension (1:2 ratio) was determined by  
86 the methodology of Jackson (1973). For the determination of soil organic carbon (SOC), the  
87 modified Walkley and Black wet oxidation method was used (Walkley and Black, 1934; Jackson,  
88 1973). Nitrogen was determine by alkaline potassium permanganate method described by Subbiah  
89 and Asija (1956). Available P was estimated calorimetrically as per Olsen method whereas

90 Available potassium in soil was extracted by Neutral ammonium acetate solution and potassium  
91 was determined using flame photometer (Jackson, 1967). Subsequently available Sulphur by  
92 Turbidimetric method given by Chesnin and Yien (1951).

93 The exchangeable cations like calcium of soils were determined using methods outlined by  
94 described by Jackson (1967). Heavy metals (Pb, Co, Cr, Cd) in the di-acid extract was determined  
95 by u atomic absorption spectrophotometer (ASS) as described by Page *et al.* (1982).

96 About 20 ~~ground-water-groundwater~~ samples from irrigated field of the farmers from study  
97 area were collected and on the basis of laboratory analysis of ~~ground-water-groundwater~~ samples  
98 results 15-20 sites of the farmers fields. The chemical parameters like pH, EC, Ca<sup>2+</sup>, Mg<sup>2+</sup>, Na<sup>+</sup>,  
99 CO<sub>3</sub><sup>-</sup>, HCO<sub>3</sub><sup>-</sup> were analysed using standard methods given by Richards 1954. SAR and RSC was  
100 determined to study suitability of water for irrigation.

101

102

## RESULTS AND DISCUSSION

### 103 1. Chemical composition and quality of irrigation water

104 Quality of irrigation water is one of the main factors that affect the physical and chemical  
105 properties of soil and ultimately, the crop growth, quality, and yield. The irrigation water must be  
106 free from excess soluble salts and from the concentration of specific substances that may create soil  
107 quality problems such as salinity, sodicity, permeability, and specific ion toxicity. Sometimes the  
108 source of irrigation creates hazards to soil quality. In the semi-arid and arid regions, irrigation is  
109 essential for successful crop production. But the main source of irrigation is groundwater (well  
110 and bore well) which is usually saline and sodic and contain toxic heavy metals when industrial  
111 areas are surrounded by them. And contain varying degree of salt concentration and their  
112 continuous application affects crop growth, quality, and yield. The analysis of irrigation water  
113 from sources of the study area for its chemical composition and to know the quality is necessary  
114 to its suitability for irrigates soils. The composition of water sample collected from well and bore  
115 well at different places are presented in (Table 1).

116

117 Table 1. Chemical Composition of Irrigation Water.

118

Sample No.	pH	Ec dS m <sup>-1</sup>	Cations			Anions		SAR	RSC	Ca/Mg	NO <sub>3</sub> -N (mg L <sup>-1</sup> )
			Ca <sup>++</sup>	Mg <sup>++</sup>	Na <sup>+</sup>	CO <sub>3</sub> <sup>-</sup>	HCO <sub>3</sub> <sup>-</sup>				
1.	7.69	0.77	2.42	0.91	8.00	1.63	3.45	3.50	1.75	3.31	5.3
2.	7.75	0.88	2.02	0.81	8.15	1.72	3.76	3.95	2.65	2.50	5.7
3.	7.77	0.89	1.97	0.80	8.24	1.72	3.79	3.87	2.47	2.49	5.8
4.	7.84	1.59	1.96	0.69	8.58	1.76	4.10	4.21	2.92	2.09	7.2
5.	7.73	0.87	2.05	0.82	8.20	1.70	3.64	3.86	2.38	2.78	5.6
6.	7.71	0.79	2.06	0.83	8.18	1.69	3.58	3.84	1.95	1.95	5.4
7.	7.98	1.67	1.70	0.60	9.40	1.76	4.12	4.95	3.21	2.84	8.1
8.	7.55	0.61	2.88	0.97	6.23	1.55	3.11	2.53	0.81	2.97	3.2
9.	7.79	1.38	1.84	0.88	8.42	1.75	3.89	4.06	2.74	2.46	5.8
10.	7.65	0.72	2.42	0.87	7.44	1.68	3.56	3.27	1.45	2.83	4.8
11.	7.59	0.64	2.61	0.92	6.66	1.60	3.29	2.83	1.24	2.79	3.9
12.	7.61	0.66	2.61	0.92	6.66	1.60	3.29	3.08	1.36	2.84	4.0
13.	7.81	1.49	1.84	0.88	8.42	1.75	3.89	4.07	2.87	2.46	7.1
14.	7.63	0.71	2.62	0.94	6.59	1.57	3.23	3.21	1.40	2.80	4.3
15.	7.87	1.62	1.81	0.63	8.48	1.66	3.90	4.33	3.12	2.87	7.3
16.	7.99	1.78	2.95	0.89	7.38	1.77	3.48	5.33	3.59	2.83	8.2
17.	7.67	0.73	2.55	0.90	7.68	1.66	3.45	3.30	1.66	2.66	4.8
18.	8.10	1.83	1.35	0.55	9.49	1.78	4.16	5.49	4.04	2.45	8.5
19.	7.91	1.65	1.86	0.64	8.86	1.68	3.99	4.47	3.17	2.91	7.8
20.	7.57	0.63	2.84	0.95	6.49	1.56	3.21	2.66	0.98	2.99	3.6

119 The pH of these water samples ranged from 7.55 to 8.1 while electrical conductivity from 0.61 to  
120 1.83 dS m<sup>-1</sup>. The sodium adsorption ratio (SAR) varied between 2.53 and 5.49; the maximum value  
121 was observed in sample no. 18 and [the](#) minimum value observed in sample no. 8. The carbonate was  
122 found ranges from 1.55 to 1.78 while bicarbonate ranged from 3.11 to 4.16. The residual sodium  
123 carbonate (RSC) was above the normal range in water samples of the study area. The irrigation  
124 water containing RSC more than 2.5 me L<sup>-1</sup> is not suitable for irrigation purposes (Richards,  
125 1954) The soluble Ca/Mg ratio of this water sample ranged from 1.95 to 3.31. The maximum value  
126 was observed in sample no 1 and minimum value was observed in sample no. 6. [Similar A similar](#)  
127 result was observed in Kadu (1997).

128 Whereas the Nitrate nitrogen content in water ~~is~~ varies from 3.2 to 8.5 the highest value  
129 was observed in sample No. 18(8.5) followed by 8.2 and 8.1 in sample No.16 and sample No.7  
130 respectively and lowest value was observed in sample No.08 (3.5).this water samples are under  
131 second class (Intensity of problem is moderate)..Very frequently ~~ground water~~ [groundwater](#)  
132 contains high amount of nitrate. When such type of irrigation water is applied on soils  
133 continuously, various properties of soils are affected, Similar result was closely [pointed](#) by  
134 Almasri and Kaluarachchi (2004).

135 **As per the quality criteria of irrigation water given by U. S. Salinity Laboratory (Richards, 1954),**  
136 **the studied water samples were medium in salinity and low in sodium, hence classified as C2S1**  
137 **and can be used for irrigation.**

138

## 139 2. Physico-chemical properties of soils of study area

140 The Particle [size-Size](#) Distribution data showed that all the soils have [a](#) high amount of  
141 clay compared to sand and silt fractions since the soil developed by basaltic parent material  
142 produce high amount clay (Eswaran *et al.* 1988). Bulk density of these soils varied from 1.32 to  
143 1.51 Mg m<sup>-3</sup>. Comparative low values of bulk density in the study area soils can be ascribed to  
144 high clay content and dominated by smectitic clay mineral, which is expanding type of clay  
145 mineral (Bharambe *et al.* 1999). The saturated hydraulic conductivity (HC) of these soils ranged  
146 from 1.07 to 2.26 cm hr<sup>-1</sup>. Low Hydraulic conductivity indicates poor structure and drainage of

147 soil. Ahmed and Wester (1988) reported that high clay content and exchangeable sodium  
148 responsible for low hydraulic conductivity (Table 2).

149 Soil reaction was low to moderately alkaline (pH 7.55 to 8.1). These soils are non-saline as  
150 indicated by the electrical conductivity, which ranged from 0.61 to 1.83 dS m<sup>-1</sup> at 25o C, but  
151 more accumulation of salts was observed in surface layer of these soils. Organic carbon content  
152 (0.70 to 1.46 g kg<sup>-1</sup>) was moderate to high; Organic matter has been found to be more or less  
153 uniformly distributed in the first meter of the profile in some Indian Vertisols. For example,  
154 organic C and total N in the profile of a deep Vertisols at the ICRISAT Similar result were closely  
155 painted by Dulal (1965); Singh *et al.* (1991), presented in (Table 3.)

156 The available Nitrogen content of soils varied from 185.44 to 300.01kg ha<sup>1</sup>. (Table 3. ) The  
157 available Phosphorus content of soils varied from 15.65 to 19.45 kg ha<sup>1</sup> The available Potassium  
158 content of soils varied from 178.80 to 330.8kg ha<sup>-1</sup> Similar results ~~were~~ also noted by Singh  
159 (1988). While the available Sulphur content of soils varied from 11.56 to 12.51kg ha<sup>1</sup> respectively  
160 were obtained with the application of ~~ground-water~~ groundwater of high RSC and SAR. similar  
161 result were closely confirmative by Patangray et al (2018)

Comment [RD5]: Explain why ?

162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172

Table No. 2. Physical properties of soils influence by ~~ground-water~~ groundwater irrigation

Sample No.	Bulk density	Particle size analysis %			Texture	Hydraulic conductivity (cm hr <sup>-1</sup> )
		Sand	Silt	Clay		
1.	1.39	14.57	32.13	53.30	Clay	1.77
2.	1.43	12.81	31.81	55.38	Clay	1.49
3.	1.42	14.81	29.34	55.85	Clay	1.56
4.	1.46	13.12	28.25	58.63	Clay	1.28
5.	1.41	12.23	34.12	53.65	Clay	1.63
6.	1.40	16.21	28.33	50.02	Clay	1.70
7.	1.49	12.81	18.13	59.06	Clay	1.07
8.	1.32	25.75	31.34	42.91	Clay	2.26
9.	1.44	11.85	31.12	57.03	Clay	1.42
10.	1.37	21.65	28.33	50.02	Clay	1.91
11.	1.34	10.58	27.72	61.70	Clay	2.12
12.	1.35	15.21	34.17	50.62	Clay	2.05
13.	1.45	25.28	39.75	34.97	Clay	1.35
14.	1.36	13.12	28.25	58.63	Clay	1.98
15.	1.47	16.12	27.11	56.77	Clay	1.21
16.	1.50	10.12	31.45	58.43	Clay	1.00
17.	1.38	14.57	32.13	53.30	Clay	1.84
18.	1.51	11.82	30.17	58.01	Clay	0.93
19.	1.48	11.85	31.12	57.03	Clay	1.14
20.	1.33	18.23	27.38	54.39	Clay	2.19



Table No. 3. Chemical properties of soil Influenced by ground-water-groundwater irrigation

Sample No.	Soil reaction (pH)	EC (dS m <sup>-1</sup> )	Organic carbon (g kg ha <sup>-1</sup> )	Available N (kg ha <sup>-1</sup> )	Available P (kg ha <sup>-1</sup> )	Available K (kg ha <sup>-1</sup> )	Available S (kg ha <sup>-1</sup> )	Exchangeable(Ca <sup>++</sup> ) (coml. (p+) kg <sup>-1</sup> )
1.	7.70	0.712	1.18	257.80	18.05	274.30	12.17	29.05
2.	7.83	0.825	1.02	233.68	17.26	242.60	11.97	28.25
3.	7.83	0.795	1.06	239.71	17.45	250.90	12.01	28.45
4.	7.91	0.933	0.90	215.59	16.64	218.40	11.81	27.29
5.	7.75	0.766	1.10	245.74	17.67	258.70	12.06	28.65
6.	7.72	0.739	1.15	251.77	17.86	266.50	12.11	28.85
7.	8.08	1.070	0.78	197.50	16.04	194.60	11.67	27.05
8.	7.49	0.543	1.46	300.01	19.45	330.8	12.51	30.45
9.	7.86	0.858	0.99	227.65	17.05	234	11.92	28.05
10.	7.65	0.660	1.27	269.86	18.46	290.60	12.26	29.46
11.	7.59	0.585	1.38	287.95	19.05	314.7	12.41	30.06
12.	7.62	0.609	1.34	281.92	18.87	306.3	12.36	29.85
13.	7.89	0.895	0.94	221.62	16.85	226.30	11.87	27.85
14.	7.63	0.634	1.30	275.89	18.65	298.50	12.31	29.65
15.	7.96	0.977	0.86	209.56	16.45	210.70	11.76	27.45
16.	8.10	1.118	0.73	191.47	15.86	186.90	11.61	26.84
17.	7.68	0.686	1.22	263.83	18.27	282.40	12.21	29.25
18.	8.13	1.167	0.70	185.44	15.65	178.80	11.56	26.65
19.	7.98	1.025	0.82	203.53	16.26	202.50	11.71	27.26
20.	7.52	0.566	1.42	293.98	19.27	322.5	12.46	30.25

183 **Table No. 4 Heavy metal content of Soil:**

Sample No.	Heavy Metal Content (mg kg <sup>-1</sup> )			
	Pb	Co	Ni	Cd
1.	0.90	1.03	1.01	0.026
2.	0.98	1.11	1.09	0.028
3.	0.96	1.09	1.07	0.028
4.	1.04	1.17	1.15	0.030
5.	0.94	1.07	1.05	0.027
6.	0.92	1.05	1.03	0.027
7.	1.10	1.24	1.23	0.033
8.	0.76	0.89	0.87	0.023
9.	1.00	1.13	1.11	0.029
10.	0.86	0.99	0.97	0.025
11.	0.80	0.93	0.91	0.024
12.	0.82	0.95	0.93	0.024
13.	1.02	1.15	1.13	0.029
14.	0.84	0.97	0.95	0.025
15.	1.06	1.19	1.17	0.030
16.	1.13	1.27	1.26	0.035
17.	0.88	1.01	0.99	0.026
18.	1.16	1.30	1.29	0.037
19.	1.08	1.21	1.19	0.031
20.	0.78	0.91	0.89	0.023


184 **Table No. 5. Correlation coefficient and simple regression equation between properties of soil with quality of irrigation water.**

185

Sr.No.	Parameters	X	R	Regression Equation
1	Hydraulic conductivity	SAR	-0.9761	Y = -0.490x + 3.479 R <sup>2</sup> = 0.9528
2	Hydraulic conductivity	RSC	-0.9907*	Y = -2.218x + 5.827 R <sup>2</sup> = 0.9816
3	SAR	RSC	0.9734	Y = 1.095x - 1.920 R <sup>2</sup> = 0.9476
4	SAR	Bulk Density	0.9761	Y = 0.070x + 1.145 R <sup>2</sup> = 0.9528
5	RSC	Bulk Density	0.9907*	Y = 0.063x + 1.270 R <sup>2</sup> = 0.9816
6	SAR	Soil Reaction (pH)	0.9833*	Y = 0.288x + 6.918 R <sup>2</sup> = 0.9669
7	SAR	EC	0.9825*	Y = 0.228x + 0.068 R <sup>2</sup> = 0.9654
8	SAR	Sulphur per cent	0.9858*	Y = -0.075x + 1.097 R <sup>2</sup> = 0.9718
9	SAR	Exchangeable calcium	0.9696	Y = -0.154x + 29.12 R <sup>2</sup> = 0.011

186 Note- All "r" values are significant at 1 per cent level. \*Showing "r" values are significant at 5 per cent level.

UNDER PEER REVIEW



187 **3. Heavy metal content of Soil Influenced by ~~ground water~~ groundwater Irrigation:**

188 Pb content in soil sample varies from 0.76 to 1.16 mg kg<sup>-1</sup> low amount of Pb 0.76 mg kg<sup>-1</sup>(sample  
189 no. 8) presented in (Table 4). and high amount of Pb 1.16 mg kg<sup>-1</sup> (sample No.18) followed by  
190 1.13 and 1.10 mg kg<sup>-1</sup> in sample No. 16 and sample No.7 was due to content of Co, Cr, Fe, Mn,  
191 Ni and Zn were associated with parent rocks and corresponded to the first principal component  
192 called the lithogenic component. A significant correlation was found between lithogenic metals  
193 and some soil properties such as soil organic matter similar result were reported by Mico *et al.*  
194 (2006)

195 Cd content in soil sample varies from 0.023 to 0.037 mg kg<sup>-1</sup> whereas Co content in soil sample  
196 varies from 0.89 to 1.30 mg kg<sup>-1</sup> While Ni content in soil sample varies from 0.87 to 1.29 mg kg<sup>-1</sup>  
197 these heavy metal like Cd, Co and Ni was found to be low amount in (sample no. 8) and high  
198 amount in (sample No.18) followed by in sample No. 16 and sample No.7 respectively is due to  
199 farmers around industrial areas are using effluents or contaminated river/well water for irrigation  
200 purpose. Since these effluents contain high amount of trace elements and other pollutant heavy  
201 metals, which hazardous to the soil and crop. Similar result was observed in Patel *et al.* (2004).

202 **4. Relationship of ~~Ground Water~~ Groundwater Irrigation on Physico-Chemical Properties**  
203 **of Soils.**

204 Some Serious effects occur on physical and chemical properties of the soils of study area  
205 due to improper and over irrigation by farmers.

206 The relationship between Hydraulic conductivity and SAR showed Negative significant  
207 correlation ( $r = 0.97$ ). presented in (Table 5) indicate that Hydraulic conductivity of soil decrease  
208 with increase in SAR of irrigation water. Similar result was observed in Vaidya *et al.* (2007). The  
209 relationship between RSC and Bulk Density showed significant positive relationship ( $r = 0.99$ )  
210 which means that RSC of ~~ground water~~ groundwater impact on soil Bulk density. Bulk density of  
211 soil increased with increasing RSC of Irrigation water. Similar result was observed in Malewar  
212 and More (1988), also Porosity of the soil decreased with the increasing sodicity level (RSC) of  
213 the irrigation water. The dispersion ratio and soil strength, however, showed an increasing trend  
214 with increasing RSC of the irrigation water. Addition of gypsum had significantly improved  
215 infiltration rate and porosity of the soil. Yadav and Kumar (2004)

**Comment [RD6]:** This sentence is very long.  
To improve readability, consider breaking this  
into multiple sentences.

216 The relationship between clay with SAR and RSC shows positive correlation ( $r=0.97$ ) It  
217 means as the SAR increases in soils, the RSC also increases. SAR is positively correlated with  
218 RSC and EC Singh and Marok (1980). the problem of RSC was associated with low salinity (EC  
219 below  $3 \text{ dS m}^{-1}$ ). They further noted multiple correlations between water quality parameters and  
220 soil characteristics and showed salinity built in soil was positively correlated with salinity of water  
221 while pH was influenced by EC, RSC, and SAR. Chauhan *et al.* (1990), whereas the relationship  
222 between SAR with Soil reaction (pH) shows Signiant positive correlation ( $r = 0.98$ ). It indicates  
223 that as the SAR increases the soil reaction (pH) increases. pH increases with increase in SAR of  
224 irrigation water Chauhan *et al.* (1990)

225 The relationship between SAR with Electrical Conductivity shows a significant positive  
226 correlation ( $r = 0.98$ ) there is positive correlation of SAR with Electrical conductivity. Singh and  
227 Marok (1980). While the relationship between SAR with Sulphur per cent and Exchangeable  
228 calcium shows significant positive correlation ( $r = 0.98$ ) respectively. It showed that as the SAR  
229 of irrigation water- increases the Sulphur and Exchangeable calcium of soil also increases, and the  
230 relationship between SAR and Bulk density showed positive correlation( $r=0.97$ ) indicate that  
231 Bulk density increases with increase in SAR of Irrigation water.

## 232 Conclusion

233 The Kanholi Bara village of Hingna Tahsil in Nagpur district situated 44 km away from  
234 Nagpur, Surrounded by industries from some side, in Kharif farmers of the region taking soybean  
235 as a main crop and giving imbalanced fertilizers. In rabbi farmers are taking Gram and providing  
236 irrigation and again use imbalanced fertilizer without doing soil and ~~ground-water-groundwater~~  
237 testing. The Impact of Imbalanced fertilizer use contaminated ~~ground-water-groundwater~~ was also  
238 noticed by other government agencies to carry out the research study in the area. In view of above  
239 facts and analytical data of soil, ~~ground-water-groundwater~~ from 20 different farms. Now it can be  
240 concluded that, erratic use of ~~ground-water-groundwater~~ with contamination adversely affects the  
241 soil physical properties. Further, the availability of nutrients also affected. Continuous use of such  
242 medium RSC water further deteriorates physical condition of soil which directly influences the  
243 soil fertility status.

244

245

246 **References**

- 247 Ahmed N. and Webster J. L. 1988 Improvement and management of salt affected Vertisols  
248 (Scotland clays) using irrigation and organic and inorganic soil amendments in Barbados.  
249 West Indies. Trans. Intl. Workshop swells shrink soils. Oxford and IBH, New Delhi: 232-  
250 233.
- 251 Almasri M. N. and Kaluarachchi J. J. 2004 Assessment and management of long-term nitrate  
252 pollution of ~~ground-water~~ groundwater in agriculture-dominated watersheds and their  
253 interaction effect on early growth of plant root and their transfer to human food chain., J.  
254 environ. Sci. health, Vol. 34(4):681-708
- 255 B Harambe P. R., Kadam S. G., Shinde S. D. and Shelke D. K. 1999 Characterization of soils of  
256 Majalgaon canal command area, Water management center. Marathwada Agriculture  
257 University, Parbhani, Maharashtra., J. Indian Soc. Of Soil Sci. Vol.47:749-759.
- 258 Chauhan R. P. Bhudayal C.P. and Vishwoi R. K. 1990 Quality assessment of under~~ground-water~~  
259 groundwater of semi-arid tract and their impact on soil. J. Indian Soc. Soil Sci. Vol.  
260 39:516-519.
- 261 Chesnin L. and Yien C. H. 1951 Turbid metric determination of available sulphur in soil. Soil Sci.  
262 Soc. of America Proceedings Vol.15:149-151.
- 263 Dulal R. 1965 Dark clay Soils of Tropical and Subtropical regions., FAO Agri. Dev. Paper No. 83,  
264 Rome, Italy 16 P.
- 265 Gulhati, N. D., and W. C. Smith. 1967. Irrigated agriculture: An historical review. In R. M.  
266 Hagan, H. R. Haise, T. W. Edminster, and R. C. Dinauer, (eds.). Irrigation of Agricultural  
267 Lands. Agronomy, 11:3-11. Am. Soc. Agron., Madison, Wis.
- 268 Jackson M. L. 1967 Soil chemical analysis, prentice hall of India private limited New Delhi.
- 269 Kadu P.R. 1997 Soils of Adasa watershed, their geomorphology, formation, characterization and  
270 land evaluation for rational land use. Ph. D. Thesis (unpub.), Dr. Panjabrao Deshmukh  
271 Krishi Vidyapeeth, Akola.
- 272 Kovda, V. A. 1973. Some effects of irrigation and drainage on soils, Irrigation, Drainage and  
273 Salinity. FAO/UNESCO. An International Source Book. p. 387-408 .

- 274 Klute A, Dirksen C. Hydraulic conductivity and diffusivity laboratory methods in methods of soil  
275 analysis Part-I. In: Klute, A. (ed) Agron Monograph 9, Madison, Wisconsin, 1986, 716-  
276 719.
- 277 Malewar G.U. and More S.D. 1988 Soil plant behavior in relation to salt affected Vertisols of  
278 Maharashtra. Retrospect's and prospects A Review. J. Maharashtra Agric.  
279 Univ.Vol.13:249-253.
- 280 Mico C., Recatala L., Peris M. and Sanchez J. 2006 Assessing heavy metal sources in Agriculture  
281 soils of a European Mediterranean area by multivariate analysis, Chemosphere,  
282 Vol.65(5):863-872.
- 283 Page A.L., Miller R.H. and Keeney D.R. 1982 Methods of soil analysis, part 2, Chemical and  
284 microbiological properties, Amer. Soc. of Agron, Madison, Wisconsin, USA.
- 285 | Patangray A. J., Kuchanwar O. D. and Ingle S. N. 2018 Assessment of [ground water-groundwater](#)  
286 quality and its effect on physico-chemical properties of soils under Pench irrigation project in  
287 Nagpur district, Maharashtra. International Journal of Chemical Studies.Vol.6 (4) pp 1559-1564
- 288 Patel K.P., Pandya R.R., Maliwal G.L., Patel K.C., Ramani V.P. and George V. 2004 Heavy  
289 Metal Content of Different Effluents and their Relative Availability in Soils Irrigated with  
290 Effluent Waters around Major Industrial Cities of Gujarat., Journal of the Indian Society  
291 of Soil Science. Vol. 52(1):89-94.
- 292 Richards L. A.1954 Diagnosis and Improvement of saline and alkaline soils., handbook, USDA  
293 Handbook No. 60., USDA, Washington D. C..
- 294 Singh K., Bhandari A.R and Tomar K. P. 1991 Morphology, genesis and classification of some  
295 soils of northwest Himalayas. Journal of the Indian Society of Soil science.
- 296 Subbiah B. V. and Asija G. L. 1956 A Rapid Procedure for the estimation of available nitrogen in  
297 soil, Current sciences Vol.25:257-260.
- 298 Varallyay, G. 1977. Soil water problems related to salinity and alkalinity in irrigated lands, ^ E. B.  
299 Worthington (ed.). Arid Land Irrigation in Developing Countries. Pergamon Press,  
300 Oxford. p. 251-264.
- 301 Yaalon D. H. 1957 Problèmes of Soil testing on Calcareous Soils. Plant and Soil. Vol.8(3):257-  
302 287.
- 303 Yadav and Kumar P. 2004 Studies on the long term effect of sodic waters, FYM and gypsum on  
304 the productivity of various cropping systems and physicochemical characteristics in light  
305 textured soil, Chaudhary Charan Singh Haryana Agricultural University.
- 306 Walkley A, Black AI. An examination of degtijareff method for determining soil organic matter  
307 and a propose modification of chromic acid titration method. Soil Science. 1934; 37:29-  
308 38.



309

310

311

UNDER PEER REVIEW

