

## Effects of *Moringa oleifera lam.* leaf powder on *Bifidobacteria* and *Escherichia coli* in the gut of albino rats

### ABSTRACT

**Aim:** This study was carried out to determine the effect of dried *Moringa oleifera* leaves on *Bifidobacteria* and *Escherichia coli* in the gut of albino rats.

**Location:** The rats were habituated under laboratory conditions at the Animal house of the Department of Zoology, Faculty of Science, University of Ibadan, for two weeks in order to adapt to the environmental conditions during the experiment.

**Duration of study:** The rats were exposed to the Moringa feed for four weeks.

**Design of study:** There were five groups in all. The 5 to 6 weeks old rats were fed with *Moringa oleifera* powder supplement except the control groups.

**Method:** No supplement of Moringa feed was administered to Group A while Group B received streptomycin antibiotics.

Groups C, D and E received dried leaf supplement of *Moringa oleifera* (DMO) 1.25 g/kg body weight (2.5%), 2.5 g/kg body weight (5%) and 5.0 g/kg body weight (10%) respectively.

**Results:** *E. coli* counts increased from  $2.3 \times 10^4$  to  $2.6 \times 10^4$  colony-forming units per gram (cfu/g) in group E, from  $2.2 \times 10^4$  to  $3.0 \times 10^4$  cfu/g in group B; but reduced from  $4.1 \times 10^4$  to  $3.7 \times 10^4$  cfu/g in group D and from  $5.4 \times 10^4$  to  $3.9 \times 10^4$  cfu/g in group C between day 20 and day 28. As from day 8, the isolates from the non-control groups were resistant to the *Moringa oleifera* extract except *E. coli* isolates in both 5% and 10% Moringa groups on day 8 with 6 mm zone of inhibition each. The rate of *Bifidobacteria* viable counts increase in group E was expressed as  $P = 0.05$  at the beginning of the experiment unlike *E. coli* counts where there was a decrease.

**Conclusion:** The *Moringa oleifera* leaf alters the microbiota in the gut, a situation which sends impulses to the brain. Thus, the *Moringa oleifera* leaf powder is a potential prebiotic for probiotics like *Bifidobacteria*, and as well induce changes in the gut-brain axis.

**Keywords:** *Moringa oleifera*, *Escherichia coli*, *Bifidobacteria*, prebiotic, gut.

### INTRODUCTION

In the sub-Himalayan areas of Afghanistan, India, Pakistan, Bangladesh, *Moringa oleifera* is found to be widely grown, likewise in the tropics. The leaves, bark, flowers, fruit, seeds, and root are used to make medicine. "Tired blood", also known as anemia, is treated using *Moringa oleifera*. Other diseases like arthritis and rheumatism are also treated using *Moringa oleifera* [19, 20]. Myriads of infections, ailments and diseases are also treated by the use of *Moringa oleifera*. These ailments and diseases include constipation, epilepsy, stomach pain, stomach and intestinal ulcers, asthma, cancer, intestinal spasms, headache, heart problems, high blood pressure, diabetes, diarrhea, kidney stones, fluid retention, thyroid disorders. Furthermore, bacterial, fungal, viral, and parasitic infections are not excluded [12, 13].

*Moringa oleifera* contains various nutrients and important classes of food such as proteins, vitamins, and minerals. A very good characteristic of *Moringa oleifera* is its antioxidant ability because it protects cells from damage. Interestingly, all parts of the *Moringa oleifera* are edible. Over the years, humans have been consuming all parts of the *Moringa oleifera* tree. Various chemicals namely alkaloids, proanthocyanidins, cinnamates, flavonoids and anthocyanins have been reported to be found in the *Moringa oleifera* tree [8, 14]. Thus, the effect of *Moringa oleifera* leaf powder on albino rats could involve actions against oxidants and inflammations with probable mechanisms of action which will be evaluated in this study. *Moringa oleifera* is potentially active against free radicals [7].

A laboratory albino rat is a rat of the species *Rattus norvegicus* (brown rat) which is bred and kept for laboratory analysis and research in numerous fields across the medical and health sciences [17]. Arguably, Wistar rats were the first set of rats to be developed for the purpose of research and stand as model organisms. Distinct characteristics such as high activity rate, long ears for hearing sensitivity, makes it preferable than other type of rats for research.

*Bifidobacterium* is a genus of Gram-negative, non-motile, often branched Anaerobacter. They are ubiquitous and inhabit major areas in the gut and tissues of humans and animals. Some of the major areas they inhabit are the gastrointestinal tract, vagina, mouth of mammals, including humans; in an endo-symbiotic relationship. *Bifidobacteria* are one of the

Formatted: Tab stops: Not at 0.5"

Formatted: Normal (Web)

57 common probiotics and major genera of bacteria that constitute a good fraction the colon flora in mammals [6, 18]. In the  
58 gut of humans, there exists a microbiota of organisms which include beneficial organisms like a few strains of *Escherichia*  
59 *coli* (*E.coli*), a type of coliform bacteria which has a few species that act in the synthesis of some vitamins. However, some  
60 strains of *E.coli* produce toxins and cause diarrhea in humans. An example is the O157:H7 strain. The gastrointestinal  
61 micro-ecosystem is always fluctuating leading to an altered microbiota which disrupts the intestinal microbial balance  
62 exposing the compromised host to opportunistic infections [17].

Formatted: Font: 10 pt

64 The *Moringa oleifera* leaf has since become an important food supplement worldwide because it possesses anti-  
65 inflammatory and antioxidant properties [2]. The *Moringa oleifera* leaf powder's beneficial and bactericidal effect on  
66 organisms in the gut representative of *Bifidobacteria*, a Gram-positive anaerobe as well as a probiotic and *Escherichia*  
67 *coli*, a Gram-negative facultative aerobe as well as a prominent coliform respectively explains the need for this study.  
68 Previous works have been done to show the effect of *Moringa oleifera* leaves extract at different concentrations on growth  
69 of studied probiotic bacteria which showed that the growth of all studied probiotic bacteria was affected by the *Moringa*  
70 *oleifera* leaf extract [3]. Furthermore, Abeer, *et al.* [3] reported that increasing the concentration of *Moringa oleifera* leaves  
71 extract from 0 to 8 % led to increase in the probiotic bacterial growth at 37 °C for 24 hours of incubation time. The aim of  
72 this study therefore, is to examine and analyze the effect of *Moringa oleifera* leaf powder on the population of  
73 *Bifidobacteria* spp. and *Escherichia coli* in the gut of Wistar Albino rats.

Formatted: Font: (Default) Arial

## 75 MATERIALS AND METHODS

### 77 Moringa leaf powder preparation:

78 Approximately 500 g of fresh tender leaves of *Moringa oleifera* were harvested from Orita Challenge suburb of the city of  
79 Ibadan in Oyo State, Nigeria. The leaves were authenticated at the Botany Department of the Faculty of Agriculture,  
80 University of Ibadan. The *Moringa* leaves were washed in water to remove dirt and later washed in 1-% saline solution to  
81 remove microbes and washed again with fresh water [4]. Water was allowed to drain for about 15 minutes. The *Moringa*  
82 *oleifera* leaves were dried in air at 25 to 28-°C, turned over at intervals with gloves and kept away from sun rays for 7  
83 days. The *Moringa oleifera* dried leaves were processed into powder form and kept in well-covered containers to prevent  
84 air [9]. One hundred grams of the dry powder was obtained, put in a dry container and stored in a cool dry place.

### 86 Animal Grouping:

87 Groups of five (n=5) *Rattus norvegicus* albino rats of weight range 170 to 230 g were used and named as follows:

- 88 • Group A (normal control) – fed with normal feed diet (50 g/kg body weight per day per rat)
- 89 • Group B (experimental control) – received streptomycin 40 mg/kg body weight/day per rat
- 90 • Group C – received dried leaf supplement of *Moringa oleifera* (DMO) 1.25 g/ kg body weight/day per rat (2.5 %  
91 Moringa feed)
- 92 • Group D – received dried leaf supplement of *Moringa oleifera* (DMO) 2.5 g/kg body weight/day per rat (5 % Moringa  
93 feed)
- 94 • Group E- received dried leaf supplement of *Moringa oleifera* (DMO) 5.0 g/kg body weight/day per rat (10 % Moringa  
95 feed).

Formatted: Font: 10 pt

96 Exposure to these treatments was done after the acclimatization period. Thorough close physical examination as well as  
97 temperature reading was done to ensure the rats were healthy.

99 **Ethical Approval:** The Animal Ethics Committee of the Department of Zoology, Faculty of Science, University of Ibadan,  
100 gave approval for the purchase of the rats with receipt number 1452592 and housing of the rats in the Animal House of  
101 the Department. A veterinary Doctor was also involved in the monitoring and analysis of the rats throughout the period.

### 103 Bacteria counts:

104 The determination of bacteria counts in the feces was performed. Viable fecal bacteria counts were determined before  
105 exposure and determined at 4-day intervals up to the 28th day of exposure. Feces were collected in sterile containers,  
106 weighed and suspended in 10 ml of 0.9-% saline solution. This was shaken vigorously for 10 to 20 minutes to allow the  
107 larger particles to settle below. About 1 ml of the suspensions was serially diluted 10-fold and appropriate dilutions were  
108 plated in duplicates on nutrient agar and incubated at 37-°C for 24 to 48 hours both aerobically and anaerobically. On the  
109 28th day of exposure, the gastrointestinal tract of animals from each group was cut open and samples were taken from  
110 the duodenum, ileum, ascending colon and descending colon of the intestines. Swabs of the intestinal parts were taken  
111 after they were cut open.

### 113 Animal housing and feeding:

114 The rats (aged between 5 to 7 weeks) were housed five per cage. Cages (24 x 18 x 12 cm) were made of plastic and  
115 metal gauze cover. The animals were habituated under laboratory conditions at the animal house of the Department of

116 Zoology, Faculty of Science, University of Ibadan, for two weeks in order to adapt to the environmental conditions during  
117 the experiment. They were fed with standard diet 50 g/kg body weight per day per rat, and water was provided *ad libitum*  
118 (without measurement) [15].

#### 119 Isolation of *E. coli*:

120 Thirty-seven grams of Eosin methylene blue (EMB) Agar (HMK Ltd), 52 g of MacConkey agar (Biotec Ltd) and 28 g of  
121 nutrient agar (HMK Ltd) were dissolved in 1l of distilled water, swirled and sterilized by autoclaving for 15 minutes at 121  
122 °C. The prepared media was allowed to cool to about 45-°C and 20 ml volumes of the liquid medium was poured  
123 aseptically into sterilized petri dishes and allowed to cool before inoculation with suspected colonies of *E. coli*.  
124

#### 125 Isolation of *Bifidobacteria*

126 Three selectively modified Bifidobacteria media (BFM), selective media recommended for the isolation of the  
127 *Bifidobacterium* spp. from tissues, feces or stool specimens were used for the isolation and identification of *Bifidobacteria*  
128 spp.: BFM 1, BFM 2 and BFM 3.

129 **Bifidobacterium Medium (BFM 1)** was specially composed with the following ingredients in grams per litre: peptone  
130 special (23.0), sodium chloride (5.0), glucose (5.0), L-cysteine hydrochloride (0.3), starch soluble (1.0), agar (15.0), with a  
131 final pH of 5.5±0.2 (at 25 °C).  
132 [6].

133 **Bifidobacterium Medium (BFM 2)** was specially composed with the following ingredients in grams per litre: MRS Agar  
134 (25.0), L-cysteine hydrochloride (5.0) [6]

135 **Bifidobacterium Medium (BFM 3)** was specially composed with the following ingredients in grams per litre: peptone  
136 (5.0), sodium chloride (5.0), lactulose (5.0), L-cysteine hydrochloride (0.5), starch soluble (2.0), tryptone (15.0), meat  
137 extract (2.0), yeast extract (7.0), peptone(5.0), riboflavin (0.001), thiamine chloride HCl(0.001), methylene blue (0.016),  
138 lithium chloride (2.0), propionic acid (5 ml) (added after sterilization at 55-°C) at a final pH ( at 25-°C) of 5.5 (with 10 N  
139 NaOH).

140 NB: Lactulose is the main carbon source.

141 Methylene blue, propionic acid and lithium chloride are inhibitors of other bacteria.

142 The low pH inhibits *Enterobacter*. **BFM 3 is a novel media composition specifically used for this study.**

#### 143 Sub-culturing:

144 The distinct colonies from agar plates were cultured on freshly prepared agar plates using proper streaking techniques.

145 Pure isolates were sub-cultured on prepared nutrient agar slants in McCartney bottles at 37-°C overnight

#### 146 Cultural characteristics of organisms

147 Distinct colonies from the plates were observed and classified based on the cultural characteristics such as shape,  
148 surface, elevation, colour, opacity, consistency and edges on the agar plate.

#### 149 Biochemical tests

150 The tests involved the use of the indole, methyl red, Voges-Proskauer and citrate (IMViC) tests as well as catalase, urease  
151 and sugar fermentation tests.

152 The test required to identify *Bifidobacteria* is the fructose-6-phosphate phospho-ketolase (F6PPK). The F6PPK detection  
153 was used for the identification of *Bifidobacteria* were the isolates were grown anaerobically for 42 hours in a 20 ml broth  
154 culture at 37-°C. The broth was centrifuged for 3 minutes to harvest the cells at 14 000 × g. The centrifuged broth formed  
155 a pellet of harvested cells which was washed twice with a phosphate buffer (0.05 M, pH 6.5, cysteine 500 mg/l) and  
156 ruptured in ice for 2 minutes and mixed with 0.25 ml each of sodium fluoro-iodoacetate and 7 fructose-6-phosphate  
157 solution. The reaction was incubated for 30 minutes at 37-°C and 1.5 ml of hydroxymine chloride (pH 6.5) was added. At  
158 room temperature, 1 ml each of 15-% tricarboxylic acid, 4 M hydrochloric acid and iron III chloride hexahydrate was  
159 added. The tube was shaken vigorously after the addition of each solution. A reddish-violet color indicated the presence of  
160 fructose -6- phosphate phosphoketolase characteristic of *Bifidobacteria* spp. The result was negative if the colour  
161 remained yellow [6].  
162

## 163 RESULTS

164 The Gram-staining reaction revealed short Gram-negative rods which were further identified as *E.coli*. indole, methyl-red,  
165 citrate (IMViC) biochemical and sugar-fermentation tests results confirmed the presence of *E. coli*. The Gram-staining  
166 reaction also revealed Gram-positive rods of various sizes and shapes, in single and chains further identified as  
167 *Bifidobacteria* of characteristic V-shape and Y-shape otherwise known as 'palisade' arrangements. Fructoso-6-phosphate  
168 phospho-ketolase test results for *Bifidobacteria*. The F6PPK test result for *Bifidobacteria* was positive if there is a reddish-  
169 violet colour immediately after shaking the tube which indicates the presence of the fructoso-6-phosphate phospho-

177 ketolase enzyme characteristic of *Bifidobacteria* spp. The F6PPK result was negative if the colour does not change from  
 178 yellow to reddish-violet [6].  
 179 ❖ *Bifidobacteria* strains were not really affected by the streptomycin but were resistant to it unlike the *E. coli* strains  
 180 that were susceptible to streptomycin. Variation in counts of *E. coli* and *Bifidobacteria* in fecal samples in group B shows  
 181 that *E. coli* was susceptible to streptomycin. Pacheco *et al.* also confirms that there is a low-level resistance of *E. coli* to  
 182 streptomycin because of the protein-freezing molecules in streptomycin [18]. In group C, D and E, the rats were fed with  
 183 2.5%, 5 %, and 10-% *Moringa oleifera* rat feed respectively (i.e 1.25 g/kg body weight, 2.5 g/kg body weight and 5.0 g/kg  
 184 body weight respectively) between days 1 to 28. Table 1 shows the proximate analysis of the *Moringa oleifera* leaf powder  
 185 indicating the protein content, ash content and other nutrients. Phytochemical analysis also showed the presence of  
 186 flavonoids and saponins  
 187  
 188

Formatted: Space After: 0 pt, No bullets or numbering, Tab stops: 0", Left

189 **Table 1:** Proximate analyses of *Moringa oleifera* leaf powder:


Formatted: Font: Not Bold

Formatted: Font: Not Bold

| Parameter                | Calculated nutrients values |
|--------------------------|-----------------------------|
| Dry matter, DM (%)       | 90.92                       |
| Crude protein, CP (% DM) | 31.5                        |
| Ether extract, EE (% DM) | 14.8                        |
| Crude fiber, CF (% DM)   | 37.4                        |
| Lysine (% DM)            | 0.94                        |
| Methionine (% DM)        | 0.42                        |
| Ash Content(% DM)        | 9.0                         |
| Calcium, Ca (% DM)       | 1.05                        |
| Phosphorus, P (% DM)     | 0.69                        |
| Vitamin B <sub>1</sub>   | 0.09                        |
| Vitamin B <sub>2</sub>   | 0.05                        |
| Vitamin B <sub>3</sub>   | 0.8                         |
| pH                       | 6.27                        |

191  
 192 Figure 1 shows the growth of *Bifidobacteria* in the non-control groups and the antimicrobial effects of *Moringa* on  
 193 *Bifidobacteria* in the respective groups. Figure 2 shows the growth of *E. coli* in the non-control groups and the  
 194 antimicrobial effects of *Moringa* in the respective groups.  
 195

Formatted: Justified, Indent: Left: 0", First line: 0"




196  
 197 **Figure 1:** The growth of *Bifidobacteria* in non-control groups and the antimicrobial effects of *Moringa* in the respective  
 198 groups.


Formatted: Indent: Left: 0", First line: 0"



212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237




However, Figures 3 and 4 shows the comparative variation in counts of *Bifidobacteria* and *E. coli* respectively in faecal samples between days 1 to 28 which reveals a decrease in the *E. coli* and *Bifidobacteria* viable counts up to day 8 in groups C, D and E. *Bifidobacteria* counts increased by more than 50-% between day 12 and day 16 with group E having the highest percentage increase but became relatively stable between day 20 and 28. *E.coli* counts increased only by 13 % in group E, by 26-% in group B; but reduced by 9.7-% in group D and 27.7-% in group C between day 20 and day 28. The effect of *Moringa oleifera* leaves extract at different concentrations on growth of studied probiotic bacteria (i.e *Bifidobacteria* spp.) showed that the growth of all studied probiotic bacteria was affected by the *Moringa oleifera* leaf extract [3]. Furthermore, Abeer, *et al.* [3] reported that increasing the concentration of *Moringa oleifera* leaves extract from 0 to 8-% led to increase in the probiotic bacterial growth at 37-°C for 24 hours of incubation time. The statement above is in tandem with the result obtained in this research work where much increase in *Bifidobacteria* counts were observed at 10-% *Moringa oleifera* concentration (in group E) than all other groups between days 12 and 16.



**Figure 3:** *Bifidobacteria* fecal counts in all groups between days 1 and 28.

However, the antibacterial activity exhibited on some bacterial isolates by *Moringa oleifera* could be as a result of the presence of flavonoids and tannins, since these phytochemicals are reported to confer antibacterial activity [2, 18]. Figure 3 shows the antibacterial effect of *Moringa oleifera* on *E. coli* especially on the rats in the 2.5-% *Moringa oleifera* feed group. The *E. coli* and *Bifidobacteria* counts in fecal samples in all groups between days 1 to 28 is shown in Table 3. Group A without Moringa feed supplement is the control group. Group B were given with normal rat feed and streptomycin 40 mg/kg body weight.

- Formatted: Font: 9 pt, Font color: Text 1
- Formatted: Font: 9 pt
- Formatted: Font: 9 pt, Font color: Text 1
- Formatted: Font: 9 pt
- Formatted: Font: 9 pt, Font color: Text 1
- Formatted: Font: 9 pt
- Formatted: Font: 9 pt, Font color: Text 1
- Formatted: Font: 9 pt
- Formatted: Font: 9 pt, Font color: Text 1
- Formatted: Font: 9 pt
- Formatted: Font: 9 pt, Font color: Text 1
- Formatted: Font: 9 pt, Font color: Text 1
- Formatted: Font: 9 pt
- Formatted: Font: 9 pt, Font color: Text 1
- Formatted: Font: 9 pt
- Formatted: Font: 9 pt, Font color: Text 1
- Formatted: Font: 9 pt
- Formatted: Font: 9 pt, Font color: Text 1
- Formatted: Font: 8 pt



238  
239  
240  
241  
242  
243  
**Figure 4:** *E. coli* fecal counts in all groups between days 1 and 28.

244  
245  
246  
**Table 3:** Antimicrobial activity of *Moringa oleifera* and streptomycin on faecal isolates of *E.coli* and *Bifidobacteria*.

|        | ZONES OF INHIBITION (mm) |   |  |   |               |    |                            |   |               |    |                          |   |                |    |                           |   |                |   |     |   |
|--------|--------------------------|---|--|---|---------------|----|----------------------------|---|---------------|----|--------------------------|---|----------------|----|---------------------------|---|----------------|---|-----|---|
|        | Group A<br>(Control)     |   | Group B<br>(Antibiotics)<br>[Positive control] |   |               |    | Group C<br>(2.5 % Moringa) |   |               |    | Group D<br>(5 % Moringa) |   |                |    | Group E<br>(10 % Moringa) |   |                |   |     |   |
|        | <i>E. coli</i>           |   | Bfd  |   | <i>E.coli</i> |    | Bfd                        |   | <i>E.coli</i> |    | Bfd                      |   | <i>E. coli</i> |    | Bfd                       |   | <i>E. coli</i> |   | Bfd |   |
|        | S                        | M | S  | M | S             | M  | S                          | M | S             | M  | S                        | M | S              | M  | S                         | M | S              | M | S   | M |
| Day 1  | 14                       | 8 | 6  | 6 | 14            | 8  | 6                          | 6 | 14            | 10 | 8                        | 6 | 16             | 10 | 6                         | 6 | 16             | 8 | 6   | 6 |
| Day 4  | 14                       | 8 | 8  | 6 | 16            | 8  | 6                          | R | 14            | 6  | 6                        | R | 16             | 6  | 6                         | R | 18             | 6 | 6   | 6 |
| Day 8  | 20                       | 8 | 6  | R | 18            | 18 | R                          | R | 16            | R  | R                        | R | 18             | 6  | R                         | R | 18             | 6 | R   | R |
| Day 12 | 20                       | 6 | 6  | 6 | 20            | R  | R                          | 6 | 20            | R  | R                        | R | 18             | R  | R                         | R | 24             | R | R   | R |
| Day 16 | 18                       | 8 | 6  | 6 | 12            | R  | R                          | R | 14            | R  | R                        | R | 14             | R  | R                         | R | 14             | R | R   | R |
| Day 20 | 18                       | 6 | 6  | 6 | 12            | R  | R                          | R | 14            | R  | R                        | R | 20             | R  | R                         | R | 14             | R | R   | R |
| Day 24 | 20                       | 6 | N  | N | 12            | R  | R                          | R | 16            | R  | R                        | R | 20             | R  | R                         | R | 20             | R | R   | R |
| Day 28 | 20                       | N | N  | N | 10            | R  | R                          | R | 14            | R  | R                        | R | 20             | R  | R                         | R | 16             | R | R   | R |

244  
245  
246  
*E.coli*-*Escherichia coli*, Bfd-*Bifidobacteria*, M-*Moringa oleifera*, S-Streptomycin, R-Resistant, N-Negligible(< 6)


Formatted: Indent: Left: 0", Tab stops: 0", Left + Not at 0.5"

247 Table 4 reveals that at 50 mg/l for *Moringa oleifera* and 0.64 mg/l for streptomycin, *E. coli* is strongly inhibited while  
 248 *Bifidobacteria* is slightly inhibited. Figure 5 shows the effect of the *Moringa oleifera* leaf powder on *Bifidobacteria* and *E. coli*  
 249 in mean and standard deviation form. Table 5 reveals a significant increase in both *Bifidobacteria* in group E and *E. coli*  
 250 in group C with high maximum limits of the organisms in the respective groups using the Dunnet's test.  
 251

252 **Table 4:** Determination of minimum inhibitory concentration of *Moringa oleifera* and streptomycin.

|   | <i>Moringa oleifera</i><br>(mg/l) |       |       |      |
|---|-----------------------------------|-------|-------|------|
|   | *50                               | 25    | 12.5  | 6.25 |
| <i>E. coli</i> (Zones of inhibition)        | 24 mm                             | 16 mm | 8 mm  | 0 mm |
| <i>Bifidobacteria</i> (Zones of inhibition) | 6 mm                              | 0 mm  | 0 mm  | 0 mm |
|   | <i>Streptomycin</i><br>(mg/l)     |       |       |      |
|   | *0.64                             | 0.32  | 0.16  | 0.08 |
| <i>E. coli</i> (Zones of inhibition)        | 32 mm                             | 20 mm | 12 mm | 0 mm |
| <i>Bifidobacteria</i> (Zones of inhibition) | 6 mm                              | 0 mm  | 0 mm  | 0 mm |

254 \*Concentrations where there is significant inhibition.  
 255



256 **Figure 5:** Effect of *Moringa oleifera* leaf Powder on *Bifidobacteria* and *E. coli* counts in all Groups in "Mean ± Sd" form

257 a= Statistically significant compared to the control group.

258 b= Statistically significant compared to the control and antibiotics group.

259 c= Statistically significant compared to the control, antibiotics and 2.5 % Moringa group.

260 d= Statistically significant compared to the control, antibiotics, 2.5 % and 5 % Moringa group

Formatted: Centered

Formatted: Centered

Formatted: Centered

Formatted: Indent: Left: 0", First line: 0"

Formatted: Font: 9 pt



274 | **Table 5:** Comparison ratio of *Bifidobacteria* and *E. coli* counts after various treatments of *Moringa oleifera* leaf-powder  
 275 | using Dunnett's test.  
 276 |

| Group | <i>Bifidobacteria</i> |            | <i>E. coli</i> |            |
|-------|-----------------------|------------|----------------|------------|
|       | Min. limit            | Max. limit | Min. limit     | Max. limit |
| C     | -8.4                  | 3.6        | -3.17          | 11.27      |
| D     | -7.85                 | 4.15       | -6.28          | 8.16       |
| E     | -3.02                 | 8.98       | -8.58          | 5.86       |

277 | NB: Groups A and B are not included because they are negative and positive controls respectively.  
 278 |  
 279 |

**Comment [SaA1]:** Why never display the results?

## 280 | DISCUSSION

281 | The *Moringa oleifera* is a very important source of micro and macro nutrients, including dietary fiber. It has respectable  
 282 | antioxidant and anti-inflammatory potency. As a single plant it contains almost all nutrients from all other herbs combined.  
 283 | It is a potential source of tocopherols, pro-vitamin A, vitamin C, calcium, protein and minerals. The *Moringa oleifera* leaf is  
 284 | the most nutritious part of the plant containing significant quantities of crude protein, vitamins and minerals. The  
 285 | percentage dry weight of crude protein of the *Moringa oleifera* leaf powder used in this study (31.5-% dry matter) is higher  
 286 | than that of other vegetable leaves. Abiodun *et al.* [2] reported that the *Moringa oleifera* leaf powder has 25.29 % protein  
 287 | dry weight of which no other vegetable has close to such amount.  
 288 |

289 |  
 290 | Over the years, people have made use of plants and herbs to cure several illnesses and diseases like fever, wounds and  
 291 | bruises, constipation, weight management, cardiovascular diseases and many others. The effect of *Moringa oleifera*  
 292 | leaves extract at different concentrations on growth of studied probiotic bacteria (i.e *Bifidobacteria* spp.) showed that the  
 293 | growth of all studied probiotic bacteria was affected by the *Moringa oleifera* leaf extract [3]. Furthermore, Abeer, *et al.* [3]  
 294 | reported that increasing the concentration of *Moringa oleifera* leaves extract from 0 to 8 % led to increase in the probiotic  
 295 | bacterial growth at 37-°C for 24 hours of incubation time. Likewise in this study, the increase in the *Moringa oleifera* feed  
 296 | across the groups also resulted into corresponding increases in bacteria counts, especially probiotic *Bifidobacteria*.  
 297 | *Bifidobacteria* has been known to prevent the incidence of inflammatory bowel diseases (IBD) like ulcerative colitis and  
 298 | Crohn's disease in the gut of humans and animals over the years [1, 16]. Thus, the increase in the load of *Bifidobacteria*  
 299 | in the gastrointestinal tract tends to reduce the rate of IBD occurrences [6].  
 300 |

**Formatted:** Tab stops: Not at 0.5"

301 | The statement by Abeer, *et al.* [3] is in tandem with the result obtained in this research work where much increase in  
 302 | *Bifidobacteria* counts were observed at 10-% *Moringa oleifera* concentration (in group E) than all other groups between  
 303 | days 12 and 16. Abeer *et al.* [3] also stated that optimum growth was recorded for all probiotic bacteria at 8-%  
 304 | concentration of the *Moringa oleifera* leaves extract at 37-°C for 24 hours incubation time. *L. jonsonii* and *B. adolescentis*  
 305 | exhibited a higher growth *L. casei* and *B. lactis*, respectively. The presence of essential amino acids in the *Moringa*  
 306 | *oleifera* leaves improved the growth of the organisms [3]. The antibacterial activity exhibited on some bacterial isolates by  
 307 | *Moringa oleifera* could be as a result of the presence of flavonoids and tannins, since these phytochemicals are reported to  
 308 | confer antibacterial activity [2, 18]. In this study, there was a considerable reduction in the amount of both *E. coli* and  
 309 | *Bifidobacteria* between day 1 and day 8; thereafter, *E.coli* counts increased only by 13-% in group E, by 26 % in group B;  
 310 | but reduced by 9.7-% in group D and 27.7-% in group C between day 20 and day 28. The study shows the highest  
 311 | reduction of *E. coli* at 2.5-% *Moringa oleifera* concentration.  
 312 |

313 | Only at 10-% Moringa powder concentration did *Bifidobacteria* grow maximally, while *E. coli* counts were low; but at 2.5-%  
 314 | concentration, there were low *Bifidobacteria* counts but high *E. coli* counts. This typically reveals the importance of  
 315 | dosage and concentration in herbal therapy and medicinal plant intake as it were in pharmaceutical drugs administration.  
 316 | It also shows that the 2.5-% moringa powder concentration was not effective enough in inhibiting the growth of *E. coli* and  
 317 | exhibiting the growth of *Bifidobacteria*, while the 10-% moringa leaf powder concentration had optimum inhibitory effect on  
 318 | *E. coli* and served as a boost for probiotic *Bifidobacteria*. The powder form of the moringa leaf mixes well with food and  
 319 | quickly starts to break down as the food chyme undergoes an enzymatic process in the mouth. In addition, the noticeable  
 320 | hyperactivity signs in the albino rats and the bidirectional neurohumoral communication system between the brain and the  
 321 | gut suggest the use of moringa leaf powder as a potential modulator of the gut flora connected with the correlating effect

on the vagus nerves which later sends the information about the intestines to the brain. Further studies can be done on how the moringa leaf supplement induces changes in the gut and the systematic effects it has on the brain. The correlation between the microbiota and human emotions could be checked in further studies putting into consideration possible feedback loops in the gut-brain axis.

Furthermore, this study reveals the *Moringa oleifera* leaf powder is being used as a modulatory tool against probable disruption of gut microbiota by certain factors causing intestinal microbial imbalance which exposes the compromised host to opportunistic infections. All animals in the groups fed with the *Moringa oleifera* leaf powder supplement were found to be healthier and stronger throughout the days of the experiment. The animals in groups C, D, and E appeared to be more active especially animals in group E with 10-% *Moringa oleifera* leaf supplement.

The hyperactivity in these groups of animals exposed to the *Moringa oleifera* leaf powder could be linked to the biochemical signals between the gastrointestinal tract (GIT) and the central nervous system (CNS) apparently described as the gut-brain axis. This explains that changes in the microbiota of the GIT immediately triggers neurotransmitters to the parts of the nervous system connected with the gut which includes the vagus nerve, CNS, enteric nervous system, autonomic nervous system and hypothalamic-pituitary-adrenal (HPA) axis. Drugs and food always cause microbiota changes which further cause changes in the levels of cytokines, which further affect the brain functions [11].

## CONCLUSIONS AND RECOMMENDATIONS

The present study suggests that the *Moringa oleifera* leaf's mechanism of action in *E. coli* involves anti-propulsive and antisecretory effects. *Moringa oleifera* can be used to create high activated carbons which are able to sequester and remove cyanobacterial microcystin-LR quite effectively and the leaf extract also appears to be capable of suppressing cyanobacterial growth as 20-to 160 mg of *Moringa oleifera* extract per liter of water is able to suppress growth of *Microcystis aeruginosa* and cause the colony count to decline [7].

Therefore, moringa dried leaf supplement is suitable for the growth of beneficial organisms like *Bifidobacteria* in the gastrointestinal tract. Beneficial bacteria of intrinsic antibiotic resistance could also be boosted by *Moringa oleifera* plant leaf to restore the gut microbiota after antibiotic treatment. *Moringa oleifera* leaf supplement is suitable for improving the intestinal microbial balance thus could serve as a modulating tool for the immune system [4, 10]. Therefore, 10 % moringa leaf in the powder form is recommended as a prebiotic for adults per meal per day for a substantial boost in the load of *Bifidobacteria* and other probiotics in the gastrointestinal tract and for an increase in activity as observed in this study.

## REFERENCES

1. Abd El-Hack, A. E., Alagawany, M., Elrys A. S., Desoky, E. M., Hala M. N. T., Elnahal, A.S.M., et al. Effect of forage *Moringa oleifera* L. (Moringa) on animal health and nutrition and its beneficial applications in soil, plants and water purification. *Agriculture* 2018; 8:145.
2. Abiodun, B. S. Adediji, A. S.; Taiwo, O.; Gbenga, A. Effects of *Moringa oleifera* root extract on the performance and serum biochemistry of Escherichia coli challenged broiler chicks. *J.Agric. Sci.* 2015; 60,505–513.
3. Abeer, E. A., Amer, B. A., Abd El-Salam, Aida S. S. Effect of *Moringa oleifera* leaves extract as a growth factor on viability of some encapsulated probiotic bacteria. *World Journal of Dairy & Food Sciences* 2014; 9 (2): 86-94.
4. Akhouri, S., Prasad, A. & Ganguly, S. Immunomodulatory effect of *Moringa oleifera* leaf extract in broiler chicks. *Indian Vet. J.*, 2014a; 91(2): 52-54.
5. Akhouri, S., Prasad, A. & Ganguly, S. Experimental study of *Moringa oleifera* leaf extract in aqueous and powder formulations on immune response of chicks by nitroblue tetrazolium reduction test. *Int. J. Chem. Pharma. Sci.*, 2014b; 2(5): 849-851.
6. Arboleya S, Watkins C, Stanton C, Ross RP. Gut *Bifidobacteria* populations in human health and aging. *Frontiers in microbiology.* 2016; 7:1204.
7. Choudhary, M. K., Bodakhe, S. H., Gupta, S. K. Assessment of the antiulcer potential of *Moringa oleifera* root-bark extract in rats. *J Acupunct Meridian Stud.* 2013; 4:34
8. Deshmukh, P., Sharma, R. K., Sharma, V., & Pankaj J. Immunomodulatory activity of *Moringa oleifera* in albino rats. *Journal of Animal Research:* 2015; 5(2)277-281.
9. Divya, S., Mandal, A. B., Biswas, A., Yadav, A. S. & Biswas, A. K. Effect of dietary *Moringa oleifera* leaves powder on growth performance, blood chemistry, meat quality and gut microflora of broiler chicks. *Animal Nutrition and Feed Technology* 2014; 14: 349-357
10. Elabd, E. M.Y., Morsy, S.M., Elmalt, H. A. Investigating of *Moringa oleifera* role on Gut microbiota composition and inflammation associated with obesity following high fat diet feeding. *Maced J Med Sci.* 2018; 6(8):1359-1364.
11. Filaretova, L. & Bagaeva, T. The realization of the gut-brain interactions with corticotrophin-releasing factor and glucocorticoids. *Current Neuropharmacology.* 2016; 14(8): 876-881.

- 382 12. Gopalakrishnan, L.; Doriya, K.; Kumar, D.S. *Moringa oleifera*: A review on nutritive importance and its medicinal  
383 application. *Food Sci. Hum. Wellness*, 2016; 5, 49–56
- 384 13. Lu, W.; Wang, J.; Zhang, H.J.; Wu, S.G.; Qi, G.H.. Evaluation of *Moringa oleifera* leaf in laying hens: Effects on  
385 laying performance, egg quality, plasma biochemistry and organ histopathological indices. *Ital. J. Anim. Sci.* 2016;  
386 15, 658–665
- 387 14. Madukwe E. U., Ezeugwu J. O. & Eme P. E. Nutrient composition and sensory evaluation of dry *Moringa oleifera*  
388 aqueous extract. *Inter J Basic Appl Scie*; 2013; 13(03):100.
- 389 15. Mun'im A, Puteri MU, Sari SP, Azizahwati. Anti-anemia effect of standardized extract of *Moringa oleifera* Lam.  
390 leaves on aniline induced rats. *Pharmacogn J*; 2016; 8(3):255-8.
- 391 16. O'Callaghan, A. & van Sinderen, D. *Bifidobacteria and their role as members of the human gut microbiota.*  
392 *Frontiers in Microbiology* 2016; 7:925
- 393 17. Ojo, N.A., Mbaya, Y.P., Simon, J., Sodipo, O.A., Adawaren, E.O., Yahi, D., *et al.* Effect of extract of *Moringa*  
394 *oleifera* leaves on leukocytic response in rats. *Vom Journal of Veterinary Science.* 2015; 10: 85 - 95
- 395 18. Pacheco O. R., WallMedrano, A., Go-I, M. G, Ramos, C. M. G, Ayala, Z. J. F, González, A. G. A. Effect of  
396 phenolic compounds on the growth of selected probiotic and pathogenic bacteria. *Letters in applied microbiology.*  
397 2018; 66(1):25-31
- 398 19. Sahay, S., Yadav, U., Srinivasamurthy, S. Potential of *Moringa oleifera* as a functional food ingredient: A review.  
399 *International Journal of Food Science and Nutrition.* 2017; 2(5) 31-37.
- 400 20. Suzana, D., Suyatna, F. D., Azizahwati, A. R., Santi P. S., Abdul Mun'im. Effect of *Moringa oleifera* leaves extract  
401 against hematology and blood biochemical value of patients with iron deficiency anemia. *J Young Pharm*, 2017;  
402 9(1): 79-84  
403  
404