

Seasonal and intra-seasonal changes in major rainfall indicators in the Ivorian cocoa zone in a context of global warming: the case of the Gôh and Lôh-Djiboua regions in west-central Côte d'Ivoire

Worldwide there is a context of global warming. This in methods. You can consider to change the title.

ABSTRACT

Aims: To understand the evolution of seasonal and intra-seasonal rainfall indicators involved in cocoa production in the Centre-West, one of the cocoa basins in Côte d'Ivoire, in order to propose technical routes adapted to the climatic conditions of the region.

Study design: Collection and processing of daily rainfall data collected by rain gauges at Divo and Gagnoa stations from 1946 to 2015.

Place and duration of studies: Divo Cocoa Research Station of the National Center for Agricultural Research, between January 2017 and June 2019.

Methodology: The break years in the time series were detected at both stations from the KhronoStat software. Key seasonal and intra-seasonal rainfall indicators for cocoa were determined and studied using the Intat+ version 3.36 agro-meteorological statistical analysis software. The behaviour of these rainfall descriptors was analysed during each sub-period by comparison. These key indicators are the dates of the beginning of the main useful rainy season (URS), the dates of the end of the URS, the duration of the URS, the cumulative rainfall and the intra-seasonal dry sequences which are the agroclimatic parameters directly influencing the main flowering of the cocoa tree on which 90% of the harvest depends.

Results: Seasonal and intra-seasonal rainfall indicators in Gagnoa and Divo are generally declining after the break-up years detected in 1972 in Divo and 1966 in Gagnoa. After the breaks, the starts of the useful Rainy Season are later and the ends are earlier than before, which results in a shortening of the post-break length of the URS and a reduction in cumulation.

intra-seasonal in both localities. Conversely, the maximum dry sequences of URS show a slight increase after the break-up dates. The station of Gagnoa was less affected by the rainfall recession than that of Divo where the downward trend in seasonal and intra-seasonal rainfall events is more severe.

Conclusion: The scarcity of seasonal and intra-seasonal rainfall events in Divo and Gagnoa is more drastic in Divo than in Gagnoa, but it remains low and does not yet pose a significant threat to cocoa production in the Gôh and Lôh-Djiboua regions.

Keywords: Seasonal and intra-seasonal changes, major rainfall indicators, cocoa farming area, Central West, Côte d'Ivoire

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74

1. INTRODUCTION

One of the greatest scourges facing humanity today and which future generations will inevitably have to face is not of viral or bacterial origin, but of environmental origin. Indeed, global warming appears to be the main concern of humankind in the 21st century. This is why each year a major supranational conference is organized to consider concerted and effective action by all countries of the world against the common enemy, whose sometimes extreme manifestations (storms, cyclones, tsunamis, torrential rains, floods, prolonged droughts, forest fires, bush fires, etc.) are becoming increasingly regular and widespread. In West Africa in general and Côte d'Ivoire in particular, climate change has long been reflected in an unprecedented reduction in rainfall amounts and river flows, as well as in a disruption of seasonality and rainfall patterns [1, 2, 3, 4], before also resulting in increasingly recurrent flooding due to a return to better rainfall conditions from the 2000s onwards [5, 6]. Climate variability has affected not only the rainfall regime but also the hydrological and plant resources [7] on which the country remains closely dependent. However, despite the fundamental importance of agriculture in the Ivorian economy, very few studies have been carried out on the involvement of climate factors in agricultural production; most research is devoted solely to the impact of climate variability and change on hydrological resources on the one hand and on groundwater resources on the other hand [2, 1, 8, 9, 10, 11, 12, 13, 14, 15, 16]. As Ivorian agriculture is predominantly rainfed, climate instability resulting from permanent changes in both monthly and annual rainfall amounts, as well as the sometimes unpredictable behaviour of climatic events within rainy seasons, is likely to disrupt agricultural yields and the incomes of the populations that depend on them. Cocoa farming, which is the spearhead of the country's economy, is unfortunately not spared by this situation. As a reminder, since 1977, Côte d'Ivoire has been the world's largest producer of cocoa, accounting for more than 40% of world production. Cocoa production accounts for 40% of export earnings at the macroeconomic level and contributes 10% to GDP formation. On the social level, about 600,000 farm managers provide a livelihood for nearly 6 million people [17]. Cocoa is grown mainly by small farmers on an area of more than two million hectares producing 1.2 million tonnes of merchant cocoa annually. Cocoa is therefore the driving force of the Ivorian economy. Faced with the predominant role of cocoa farming in the national economy, this study aims to analyse the evolution and potential impact of seasonal and climatic events. ?
intra-seasonal factors determining cocoa production in the Centre-West, which is one of the bastions of cocoa in Côte d'Ivoire. The ultimate objective is to enable the various actors in the cocoa sector to put in place more resilient measures to deal with climatic hazards.

2. METHODOLOGY

2.1 DESCRIPTION OF THE STUDY AREA

The study was carried out in the Centre-West of Côte d'Ivoire, in the regions of Lôh-Djiboua and Gôh, -Djiboua whose respective chief towns are Divo and Gagnoa. It is located in the second cocoa loop (1960-1970), between latitudes 5°22' and 6°26' N and longitudes 4°58' and 6°34' W and covers an area of 10792 km² (Figure 1). These regions belong to the district of Gôh-Djiboua, which is located in the humid tropical zone [18] where rainfall fluctuates between 1200 and 1600 mm per year [19] and is divided into four seasons: a major rainy season from March to June, a small dry season in July and August, a small rainy season from mid-September to mid-November, and a major dry season from December to February. The average humidity of 85% is subject to strong seasonal variations. The average temperature is 27°C and varies annually between 19° and 33°C. The duration of annual exposure is about 1800 to 2000 hours. The predominant climax is the semi-deciduous dense rainforest. The soils are moderately to highly

75 desaturated ferrallitic [20, 21]. The humus horizon is thin, but rich in organic matter, weakly acidic and well
76 structured under primary forest. These soils are suitable for perennial crops such as coffee and cocoa.

77 78 **2.2. Data used and methods of analysis**

79 80 **2.2.1. Data used**

81
82 To carry out this study we used daily rainfall data from Divo and Gagnoa over the period 1946-2015. The
83 rainfall database used comes from the meteorological service of the Sustainable Soil Management and
84 Water Management Programme of the Centre National de Recherche Agronomique (CNRA); but also
85 from the historical database of the Office pour la Recherche Scientifique des Territoires d'Outre-Mer
86 (ORSTOM), now the Institut de Recherche pour le Développement (IRD).

87 88 89 **2.2.2. Methods of analysis**

90 91 **2.2.2.1. Period selection, criticism and data filling**

92
93 The temporal window chosen for this study (70 years) has the advantage of having fairly homogeneous
94 data with complete annual series, with the exception of certain years when some data are missing,
95 particularly on the Divo station, where the incomplete data of 1946 were replaced by those of the Tiassalé
96 station, only 60 km away. This recent and fairly long database provides objective and more representative
97 trends of the current climatic conditions of the departments studied.

98 99 100 **2.2.2.2. Methods for determining breaks within interannual rainfall series**

101
102 The "KhronoStat" software, designed by HydroSciences Montpellier and freely accessible on the SIEREM
103 website (<http://www.hydrosciences.org/spip.php>) [22] was used to detect possible breaks in time series. A
104 break can be generally defined by a change in the probability law of the time series at a given time, most
105 often unknown [11]. This program includes many specific tests of a change in the behaviour of the variable
106 in the time series. The detection of breaks within time series required the application of a set of methods,
107 including the Pettitt test [23], the Buishand "U" method [24], the Bayesian procedure of Lee and Heghinian
108 [25] and the Hubert segmentation procedure [26]. It is at the end of the application of these various tests
109 that a failure date detected by the majority of the tests was chosen.



110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130

Centre-West region

Côte d'Ivoire ?

131

132



133

134

135

136

137

138

139

140

141

142

143

144

145


146

4°58' W

6°34' W

N. 07.0

N. 07.0



4°58' W

6°34' W

N. 77.0

N. 77.0

Figure 1: Presentation of the study area ?

the cocoa tree

Rewrite.

147 **2.2.2.3. Determination of key seasonal and intra-seasonal rainfall indicators in cocoa production**

148
149 The productivity of a cultivated cocoa tree necessarily requires regular growth, abundant flowering and
150 fruiting as well as well-distributed foliar outbreaks throughout the year. To do this, it must be in favourable
151 climatic conditions, obeying the following criteria: (1) the annual rainfall amounts are between 1200 and
152 1500 mm[27] but a minimum annual threshold of 1200 mm is sufficient to consider its establishment in a
153 region[28]; (2) the annual cumulative rainfall during the high rainy season is greater than 700 mm[29]. It is
154 this rain that triggers the first flowering of the cocoa tree for the main harvest from September to January;
155 (3) the duration of the dry season is less than 3 months[27] otherwise the cumulative rainfall during this
156 period must be greater than 70 mm[30]. (4) because of the weak rooting of its lateral roots, 20 days
157 without rain are sufficient to lose the crop[31]. As part of this work, we analyzed the seasonal and intra-
158 seasonal rainfall descriptors that are critical to the success of cocoa farming, using the Instat+v.3.37
159 software. These are the start and end dates of the major rainy season or Useful Rainy Season (URS), the
160 length of this season, as well as the cumulative rainfall and maximum dry sequences during the major
161 rainy season.
162

163 **2.2.2.3.1. Definition of the parameters studied**

164
165 **2.2.2.3.1.1. Start and end dates of the great rainy season in Guinea area**

166
167 The start and end dates of the URS (Useful Rainy Season) are automatically determined by the Instat+
168 v3.37 software. The determination of these dates takes into account rainfall, the value of potential
169 evaporation (FTE) and the useful soil reserve.
170 The approach used for this study is based on both the work of [32] and that of
171 [33] and [34] who adapted Sivakumar's method ([35],[36],[36],[37]) to Ivorian climatic realities. The
172 following criteria for determining the length of rainy seasons in the Guinean zone under bimodal conditions
173 have been established:

- 174 - the date of the beginning of the main useful rainy season is after¹ February, when the amount of
175 rain collected in 2 consecutive days is at least equal to 20 mm without a dry sequence of more
176 than 7 days in the following 20 days;
- 177 - the end date of this season corresponds to the first day after¹ July when soil capable of containing
178 70 mm of available water is completely exhausted by a daily loss of evapotranspiration of 4 mm;
179 i.e. when the water balance is zero.

180
181 **2.2.2.3.1.2. Duration of great rainy season**

182
183 The duration of each rainy season is obtained by differentiating between the start and end dates of the
184 seasons.
185

186 **2.2.2.3.1.3. Cumulative rainfall**

187
188 Seasonal rainfall totals are the sum of the rainfall amounts recorded during a rainy season. It represents
189 the amount of rain collected during the agricultural season.
190

191 **2.2.2.3.1.4. Maximum dry sequences**

192
193 A dry sequence is defined as the number of consecutive rain-free days with a height greater than the
194 minimum value (1 mm) of the smallest of the classes of daily precipitation amounts proposed by the
195 international standards defined by the World Meteorological Organization [38]. The different classes are
196 defined according to the number of rainy days with a height between: 1 and 10 mm (P1); 10 and 30 mm
197 (P2); 30 and 50 mm (P3); >50 mm (P4).
198
199

200
201
202
203
204
205
206
207
208
209
210

3. RESULTS

3.1. DETECTION OF YEARS OF BREAKS IN THE RAINFALL SERIES

Most of the tests identified rainfall breaks in 1972 at Divo station and in 1966 at Gagnoa station (Table 1). These results indicate a specific change in the average in the rainfall series of the departments studied.

Table 1: Breaks in the rainfall series established by the various tests

Stations	Failure tests								Date indicated by the majority of tests
	Pettitt		Lee and Heghinian		Buishand		Hubert		
	Year	Proba.	Year	Proba.	Year	Proba.	Year	Proba.	
Divo	1972	0,001	1955	0,65	1972	0,99	1953 ; 1956	0,1	1972
Gagnoaa	1966	0,00678	1966	0,1968	1966	0,99	1954 ; 1965	0,1	1966

211 Proba : Probability
212
213

3.2. SEASONAL AND INTRA-SEASONAL MUTATIONS OF THE MAIN CLIMATIC VARIABLES OF THE COCOA CROP YEAR

3.2.1. Start of the Useful Rainy Season (URS)

Before the break-up years, the season began on average on 7 March in Divo and 8 March in Gagnoa. On the other hand, after the breakdowns, URS begins on average on 17 March in Divo and on 7 March in Gagnoa. The onset of URS is therefore on average 10 days later in Divo (current period) than before 1972, unlike in Gagnoa, where the season is currently one day earlier than in the period before 1966 (Table 2).

Table 2: Average start dates for URS in Divo and Gagnoa

Station	DIVO		GAGNOA	
	1946-1971	1972-2015	1946-1965	1966-2015
Period of time	March 7	March 17	March 8	March 7
Average	March 7	March 17	March 8	March 7
Standard deviation	21	27	18	19
Coefficient of variation (%)	31	35	26	28

227
228
229
230

3.2.2. End of the Useful Rainy Season (URS)

Before the breaks, the season ends on average on 27 August in Divo and on 8 August in Gagnoa. After the breaks, the end of the season occurs on average on July 27 in Divo, i.e. one month earlier than the period before the break. It is also, on average, one day earlier (August 7) than the pre-break period in

234 Gagnoa. The end of the season that occurred, on average, much earlier in Gagnoa (19 days earlier than
 235 in Divo) now takes place much later (11 days later than in Divo) (Table 3).

236
 237
 238
 239

Table 3: Average end dates of URS in Divo and Gagnoa

Station	DIVO		GAGNOA	
	1946-1971	1972-2015	1946-1965	1966-2015
Period of time	1946-1971	1972-2015	1946-1965	1966-2015
Average	August 27th	July 27th	August 8	August 7
Standard deviation	55	14	29	40
Coefficient of variation (%)	23	7	13	18

240
 241
 242
 243

3.2.3. Lengths of the URS

244 On average, during the period 1946-1971 in Divo, the main rainy season lasted 173 days (\approx 5 months 23
 245 days) while during the post-rupture period (1972-2015), it did not exceed 132 days
 246 (\approx 4 months 12 days); a 41-day shortening (\approx 1 month 11 days). On the other hand, in Gagnoa, URS has
 247 an average lengthening of 4 days compared to the period before rupture (1946-1965). It went from 154
 248 days (\approx 5 months 4 days) to 158 days (5 months 8 days). Before the breaks, the useful rainy season was
 249 on average longer in Divo, but this shortening by more than one month reversed the trend (Table 4).

250
 251
 252

Table 4: Average length of URS in Divo and Gagnoa

Station	DIVO		GAGNOA	
	1946-1971	1972-2015	1946-1965	1966-2015
Period of time	1946-1971	1972-2015	1946-1965	1966-2015
Average	173 days	132 days	154 days	158 days
Standard deviation	63	31	29	44
Coefficient of variation (%)	37	24	19	29

253
 254
 255

3.2.4. Intra-seasonal rainfall totals for URS

256 During the two sub-series before the breaks observed at the two stations, the average Divo accumulation
 257 is 1168 mm while the Gagnoa accumulation is 886 mm. However, during the two post-rupture periods, the
 258 average intra-seasonal rainfall totals are 701 mm in Divo and 826 mm in Gagnoa respectively. The
 259 average of the cumulative heights, which was higher at Divo, experienced a clear post-breakdown
 260 regression to reach almost the required threshold. Gagnoa experienced a smaller reduction (Table 5).
 261 Regardless of the station and the observation period, the cumulative threshold (700 mm) required to meet
 262 the cocoa tree's water requirements during URS is exceeded (Table 5).
 263

264
 265
 266
 267
 268
 269
 270
 271
 272
 273

274 Table 5: Average cumulative rainfall (mm) of URS in Divo and Gagnoa
 275

Station	DIVO		GAGNOA	
	1946-1971	1972-2015	1946-1965	1966-2015
Period of time	1946-1971	1972-2015	1946-1965	1966-2015
Average	1168	701	886	826
Standard deviation	670	211	216	257
Coefficient of variation (%)	57	30	24	31

276
 277 **3.2.5. Maximum dry sequences**

278
 279 **3.2.5.1. Maximum intra-seasonal dry sequences of URS**

280
 281 Before ruptures, dry sequences within the Useful Rain Season last on average 8 days in Divo and 7 days
 282 in Gagnoa during URS. On the other hand, the maximum post-rupture dry sequences of URS are up by 1
 283 day in Divo and Gagnoa, where they are worth 9 days and 8 days respectively. It can be seen that in Divo,
 284 dry sequences greater than 20 days and 30 days are almost non-existent before rupture (respectively 1%
 285 and 0%) while during the post-rupture period, 5% of years contain dry sequences of more than 20 days
 286 and 3% have dry episodes of more than 30 days. In Gagnoa, on the other hand, the importance of non-
 287 rainfall episodes of more than 20 days in URS remains unchanged after the break (2% of years) (Table 6).
 288

289 Table 6: Descriptive statistics of the maximum dry sequences of URS in Divo and Gagnoa
 290

Station	DIVO		GAGNOA	
	1946-1971	1972-2015	1946-1965	1966-2015
Periods of time	1946-1971	1972-2015	1946-1965	1966-2015
Maximum	25	56	22	29
Minimum	2	2	2	2
Average	8	9	7	8
Standard deviation	4	8	4	4
Coefficient of variation (%)	59	80	63	58
SS ≤ 20 days (%)	99	95	98	98
SS ≤ 30 days (%)	100	97	100	100
SS ≤ 60 days (%)	100	100	100	100
SS ≤ 90 days (%)	100	100	100	100

291 **SS:** dry sequences
 292
 293
 294

295 **4. DISCUSSION**

296
 297 The various statistical tests detected major rainfall accidents in the rainfall series of the two localities
 298 studied. As a reminder, many studies have identified changes in stationarity in African hydroclimatic series
 299 during the 20th century, especially those corresponding to a sudden decrease in precipitation in the late
 300 1960s in the Sudano-Sahelian zone ([39], [40], [41], [42], [43]) and in the Guinean and Sudano-Guinean

(39-43)

[32, 44, 5, 14, 45]

301 zone ([32], [44], [5], [14], [45]). This is why, [46] stated that "the sudden inflection point was observed in
302 1970 making it the pivotal year between two periods of distinct rainfall patterns". In our case, these break-
303 up dates were identified in 1972 in Divo and in 1966 in Gagnoa. These dates are perfectly consistent with
304 the break-up years indicated by [44] and [47] for the same locations. These ruptures are generally part of
305 the period designated for the majority of West African countries by [2] and [8]. Indeed, by studying the
306 evolution of the time series of 33 rainfall stations in West Africa in the Sahel zone, Sudano-Guinean and
307 Guinean, [40] have highlighted significant ruptures mostly between 1968 and 1972 [48]. This is the same
308 observation in our study because although the majority of authors agree that the rupture occurred in Côte
309 d'Ivoire around 1970, this year is only given as an indication [5]. Indeed, it has been designated as a
310 pivotal year in the evolution of time series in West Africa because it corresponds to the break-up date of
311 most stations in the West African region. However, there are several stations that experience a break at
312 dates other than 1970, but which are close to it. After the years of disruption, a general rainfall recession
313 set in in the regions of Gôh and Lôh-Djiboua. These post-breakdown climatic provisions are corroborated
314 by the work of [49] who, in a study on the impact of climate variability on coffee and cocoa production in
315 central-eastern Côte d'Ivoire, which was the first cocoa loop, showed that the rainfall series in the
316 departments of Daoukro, Bocanda, Agnibilékro, M'bahiakro and Abengourou show a dry or deficit period
317 after the breaks. in the

318 The study of the evolution of the high rainy season in cocoa production, commonly referred to as the
319 "Useful Rainy Season or URS" in this work and the distribution of its main descriptors throughout the
320 season was essential insofar as several authors ([29], [50], [51], [52]) have shown that it is the rains
321 received during this period of the year that trigger the first flowering of the cocoa tree on which 90% of the
322 harvest generally occurs between September and January depends. A better knowledge of the behaviour
323 of these climatic variables makes it possible to better understand their impact on cocoa production in the
324 study sector. The evaluation of the onset of URS during the wet and dry sub-periods on either side of the
325 break-up years reveals that the post-break-up start of the season in Divo is 10 days later than the pre-
326 break period (17 March compared to 7 March). The same trend is observed in Gagnoa with a one-day
327 post-rupture delay (7 March instead of 8 March). This late post-rupture start is the same as that observed
328 by [32] in southern Togo, a cocoa-producing country with similar climatic conditions to the Ivorian forest
329 south that hosts our study area. During their study, these authors demonstrated that the arrival of the
330 potentially useful high rainy season is later in the 1970-2000 period (post-rupture) than in the pre-break
331 period (1950-1969). Indeed, they observed that early coastal rains that began, on average, from 15 to 28
332 March in the period preceding the rupture (1950-1969) are now observed only from 29 March. According
333 to [53], a delay of at least 25 days in the beginning of the rainy seasons was observed in the Sudano-
334 Sahelian zone of Nigeria during the 1983 drought, which was a year of extreme drought in West Africa.
335 This delay even reaches 40 to 50 days in some parts of Nigeria during the same year. The work of [54] in
336 relation to the impact of rainfall variability on the water balance of pineapple-grown soils in southern Benin
337 has also shown that the seasons start with a delay of 5 to 25 days. Unlike the start of the useful rainy URS
338 season, which is late after breaks, the end of URS in Divo and Gagnoa have an early start of one month
339 and one day respectively. These results are consistent with those of ([55], [5], [34]).

340 Concerning the length of the useful rainy season, it can be seen that the later start and earlier end of the
341 main rainy season observed after the rainfall accidents (1966 and 1972) in the two stations analysed
342 necessarily leads to a post-rupture shortening of this season. This is why the duration of URS has been
343 reduced from 173 to 132 days in Divo (shortened by more than a month). This post-rupture regression is
344 corroborated by the work of [32] during which the statistical analysis revealed a narrowing of the duration
345 of potentially useful URS rainy seasons, due to a delay in their installation and/or early termination. This result
346 is also in agreement with a study by [56] in relation to the identification of rainy season start and end dates
347 in Senegal and East Africa. In this study, these experts demonstrated that in Senegal, significant trends in
348 start and end dates indicate a shortening of the rainy season between 1950 and 1992. Other authors such
349 as [14] have reached the same conclusion in the N'zi catchment area, a tributary of the Bandama River in
350 Côte d'Ivoire. Unlike Divo, we observe that in Gagnoa, on the other hand, URS increased by an average
351 of 4 days. Two main reasons could explain this phenomenon. The first one which is more plausible is the
352 one highlighted by [6] which showed that since the early 2000s, a new period of high rainfall has appeared
353 in many localities in humid tropical climates. The second reason is that put forward by Chaouche [57] who
354 demonstrated during his work in Sudano-Sahelian Africa, that the rainy season does not always follow a
355 trend of reduction parallel to that of the annual height.

This is a long sentence which is the point? Use period

1

this author

356 In terms of rainfall totals recorded within the URS, we have shown that the average intra-seasonal totals
357 collected at Divo and Gagnoa stations (respectively 874 mm and 843 mm) are well above the threshold
358 totals required by the cocoa tree during the useful rainy season. Nevertheless, we note a worrying trend
359 towards a reduction in these accumulations after breaks (701 mm after compared to 1168 mm before in
360 Divo and 826 mm after compared to 886 mm before in Gagnoa). This ability to regress is confirmed by
361 [50] who, by analyzing the sub-period 1978-2007 noted that rainfall averages tended to become
362 insufficient (less than 700mm) to meet the cocoa tree's water needs. Indeed, the latter discovered very
363 low average rainfall totals (only 164.7 mm in Divo and 652.1 mm in Gagnoa).
364 For the maximum dry sequences of URS, the respective averages after breaks are only 9 and 8 days in
365 Divo and Gagnoa. In addition, during the post-rupture period there are only 5% of years in Divo and 2% of
366 years in Gagnoa that have maximum dry sequences exceeding 20 days without rain that could be harmful
367 to cocoa tree productivity. Similarly, the increase in post-rupture intra-seasonal dry-season sequences [58]
368 of only one day during the useful rainy season is not likely to disturb the flowering of cocoa trees.

370 5. CONCLUSION

371
372 This study, initiated in the Centre-West of Côte d'Ivoire, which is one of the main cocoa production areas
373 in the country, updated the distribution of the main seasonal and intra-seasonal rainfall indicators
374 influencing cocoa production in the Gôh and Lôh-Djiboua regions. The various analyses show that Divo
375 and Gagnoa have a clear downward trend after the rainfall breaks identified in 1966 and 1972. This trend
376 towards rainfall depreciation had a direct impact on key seasonal and intra-seasonal indicators during the
377 periods 1972-2015 in Divo and 1966-2015 in Gagnoa. Thus, the start of the Useful Rainy Season is now
378 later and the endings are earlier. This leads to a shortening of URS overall in these departments, which is
379 accompanied by an intra-seasonal decrease in cumulations and an increase in maximum dry sequences
380 during URS. This seasonal and intra-seasonal degradation of agro-climatic variables remains insignificant
381 and is therefore not yet a limiting factor for the sustainability of cocoa production in this historical cocoa
382 bastion. However, if the degeneration of these descriptors continues, these areas could become marginal,
383 which is why improved varieties that are more resistant to climatic hazards, particularly drought, should be
384 developed and disseminated in these regions.

387 COMPETING INTERESTS

388
389 "The authors stated that there are no competing interests."

391 CONSENT (IF APPLICABLE)

392
393 Not concerned

395 ETHICAL APPROVAL (IF APPLICABLE)

396
397 Not concerned

399 REFERENCES AND REFERENCES

- 400
401
402 1. Servat E., Paturel J. E., Lubès H., Kouamé B., Ouédraogo M. & Masson J. M. (1997). Climatic
403 variability in humid Africa along the Gulf of Guinea. Part I: detailed analysis of the phenomenon in
404 Côte d'Ivoire, *Hydrol. Sci. J.*, (191), 1-15.
- 405 2. Paturel J. E., Servat E., Kouamé B., Lubès H., Ouédraogo M. & Masson J.M. (1997). Climatic
406 variability in humid Africa along the Gulf of Guinea. Part II: an integrated regional approach.
407 *Hydrol. Sci. J.*, (191), 16-36. *No bold*
- 408 3. Brou Y. T., Servat E. & Paturel J. E. (1998). Human activities and climate variability: case of the
409 Ivorian southern forest, *IAHS Publ*, (252), 365-373.

bold?

- 410 4. **Soro G. E., Anouman D. G. L., BI T. G., Srohorou B. Savannah I.** (2014). Characterization of
411 meteorological drought sequences at various weather scales in Sudanese type climates: case of
412 the extreme northwest of Cote d'Ivoire. *LARHYSS Journal* (18),
413 1112-3680.
- 414 5. **Noufé D.** (2011) Changement hydroclimatique et transformation de l'agriculture : exemple des
415 paysanneries de l'Est de la Côte d'Ivoire, Thèse de doctorat de géographie, hydrosciences
416 Montpellier, Université Paris 1 Panthéon-Sorbonne, Paris, 375p.
- 417 6. **Noufé D., Lidon B., Mahé G., Servat E., Brou Y.T., Koli B. Z. & Chaléard J.-L.** (2015). Climate
418 variability and rainfed maize production in eastern Côte d'Ivoire. *Hydrol. Sci., J.* 56 (1), pp. 152-
419 167.
- 420 7. **Aguilar L. A. A. A.** (2009). Impact of recent climate variability on Senegal's niayas ecosystems
421 between 1950 and 2004. Thesis; Université du Québec à Montréal, Québec, 185p.
- 422 8. **Servat E., Paturel J. E., Kouamé B., Travaglio M., Ouédraogo M. & Boyer J. F.** (1998).
423 Identification, characterization and consequences of hydrological variability in West and Central
424 Africa. *IAHS, 252*, pp 323-337.
- 425 9. **Servat E., Paturel J. E., Lubès-Niel H., Kouamé B., Masson J. M., Travaglio M. & Marieu B.**
426 (1999). Different aspects of rainfall variability in non-Sahelian West and Central Africa. *Journal of*
427 *Water Sciences*, 12 (2): 363-387
- 428 10. **Ardoin-Bardin S., Lubès-Niel H., Servat E., Dezetter A. Boyer J.F.** (2003) "Analysis of the
429 persistence of drought in West Africa: characterization of the situation in the 1990s", *IAHS*
430 *Publication, vol. 278*, 223-228.
- 431 11. **Ardoin-Bardin S.** (2004). Hydroclimatic variability and impacts on water resources in large river
432 basins in the Sudano-Sahelian zone. Doctoral thesis. University of Montpellier II. 437p.
- 433 12. **Goula B. T. A., Kouassi V. J. & Savane I.** (2006). Impacts of climate change on water resources
434 in humid tropical zones: Case of the Bandama catchment area in Côte d'Ivoire. *African Agronomy,*
435 *Vol. 18, No. 1*, pp: 1-11.
- 436 13. **Kouassi A.M., Kouamé K.F., Goula B.T.A., Lasm T., Paturel J.E. & Biémi J.** (2008).
437 Influence of climate variability and land use change on the rain-flow relationship based on a
438 global modelling of the N'zi (Bandama) catchment area in Côte d'Ivoire. *Revue Ivoirienne des*
439 *Sciences et Technologie, Vol. 11*,
440 pp. 207-229.
- 441 14. **Kouassi A. M., Kouamé K. F., Koffi Y. B., Dje K. B., Paturel J. E. & Oulare S.** (2010). Analysis
442 of climate variability and its influences on seasonal rainfall patterns in West Africa: the case of the
443 N'zi (Bandama) catchment area in Côte d'Ivoire", *Cybergeog. European Journal of Geography,*
444 *Environment, Nature, Paysage*, published online on 07 December 2010, modified on 10
445 December 2010. URL: <http://cybergeog.revues.org/23388>. Accessed May 09, 2011.
- 446 15. **Fossou R.M.N., Soro N., Traoré B., Lasm T., Sambou S., Soro T., Orou K. R., Cissé T.M., &**
447 **Kane A.** (2014). Climate variability and its impact on surface water resources: the case of the
448 Bocanda and Dimbokro stations, East Central Ivory Coast in West Africa, *Africa science*, 10(4),
449 pp.118-134.
- 450 16. **Fossou R. M. N., Lasm T., Soro N., Soro T., Soro G., De Lasme O.Z., Baka D., Onetie, O.Z.** &
451 **Orou R.** (2015). Climate variability and its impact on groundwater resources: the case of the
452 Bocanda and Dimbokro stations in central eastern Côte d'Ivoire (West Africa). *Larhyss Journal*. 21
453 : 97-120
- 454 17. **Tano A. M.** (2012). Cocoa crisis and strategies of producers in the sub-prefecture of Méadji in
455 southwestern Côte d'Ivoire. PhD from the University of Toulouse, UMR Dynamiques Rurales, 261
456 p.
- 457 18. **Eldin M.** (1971). Le climat In le milieu naturel de la Côte d'Ivoire. pp. 73-108.
- 458 19. **Kouamé B., Koné D. & Yoro G.**, (2006). Rainfall in 2005 and 2006 in the southern half of Côte
459 d'Ivoire: In. *The CNRA in 2006*, pp. 12-13.
- 460 20. **Perraud A.** (1971): Soils. In: *The natural environment of Côte d'Ivoire*. ORSTOM thesis n°50.
- 461 21. **Oswald M.**, 1997. Reconstruction of a society through several crises: the rural Bété society.
462 Thesis of 3rd cycle. Institut Nationale Agronomique Paris-Grignon, Paris, Volume 1, pp. 146-180.
- 463 22. **Boyer J.F., Dieulin C., Rouche N., Cres A., Servat E., Paturel J.E. & Mahé G.** (2006). SIEREM
464 an environmental information system for water resources. *IAHS publication, vol. 308* p: 19

Why no bold?

- 465 23. Pettitt A. N. (1979). A non-parametric approach to the change-point problem. Applied statistics, 28, 126-135. DOI : 10.2307/2346729
- 466
- 467 24. Buishand T. A. (1984). Tests for detecting a shift in the mean of hydrological time series. Journal of Hydrology, 58, 51-69.
- 468
- 469 25. Lee A. F. S. & Heghinian S. M. A (1977). Shift of the mean level in a sequence of independent normal random variables: A Bayesian approach. Technometrics, 19 (4), pp. 503-506.
- 470
- 471
- 472 26. Hubert P., Carbonel J. P. & Chaouche A. (1989). Segmentation of hydrometeorological series. Application to rainfall and flow series in West Africa. Journal of Hydrology, 110, 349-367.
- 473
- 474 27. Brou Y.T., NGoran J.A.K., Bicot S & Servat E. (2003). Climate risk and agricultural production in Côte d'Ivoire: effect of rainfall variations on cocoa production. In: Proceedings of the 14th International Conference on Cocoa Research (Accra, Ghana, 18-23 October 2003): 259-267.
- 475
- 476
- 477 28. Mossu G. (1990). The cocoa tree. The tropical agriculture technician. Maisonneuve and Larousse. ACCT. 159 p.
- 478
- 479 29. Mian K.A. (2007). Contribution to the agro-climatic analysis of the cocoa growing area in Côte d'Ivoire. Final thesis, AGRHYMET Regional Centre, 64p.
- 480
- 481 30. Dian B. (1978). Geographical aspects of the coffee and cocoa pair in the Ivorian economy. NEA, Abidjan-Dakar. 111 p.
- 482
- 483 31. Lachenaud P. (1992): Factors of fruiting in cocoa trees. Influence on the number of seeds per fruit (*Theobroma Cacao* L.), Grignon : INA, 1992, *Doctoral thesis*, 188 P.
- 484
- 485 *Italics* 32. Adewi E., Badameli K. & Dubreuil V. (2010). Evolution of potentially useful rainy seasons in Togo from 1950 to 2000. Climatology 7, 89-107
- 486
- 487 33. Sarr B. (2007). User manual for agrometeorological engineers. AGHYMET Regional Centre, Niamey, 72p.
- 488
- 489 34. Goula B. T., Srohourou B., Brida, A., N'zué K. A. & Goroza G. (2010). Determination and variability of growing seasons in Côte d'Ivoire. Int J Eng Sci, 2(11), 5993-6003
- 490
- 491 35. Sivakumar M. (1988). Predicting rainy season potential from the onset of rains in the southern sahelian and soudanian climatic zone of West Africa. Agricultural and forest meteorology, 42, 295-305.
- 492
- 493
- 494 36. Balme M., Wales S. Lebel T. (2005). Start of the rainy season in the Sahel: variability at hydrological and agronomic scales analysed from EPSATNiger data. Drought, 16(1), 15-22.
- 495
- 496 37. Stern R., Rijks D., Dale I., Knock J., (2006). INSTAT+ for Windows V3.036 Statistical Services Center, University of Reading: Reading.
- 497
- 498 38. O.M.M. (1990). Guide des pratiques climatologiques, (100). Geneva, Switzerland, 515 p.
- 499
- 500 39. Sircoulon J. (1987). Variation in river flows and lake levels in West Africa since the beginning of the 20th century. In: The Influence of Climate Change and Climatic Variability on the Hydrologic Regime and Water Resources (Proc Vancouver Symposium August 1987). IAHS Publication, No. 168, pp. 13-25.
- 501
- 502
- 503 40. Hubert P., Carbonel J. P. & Chaouche A. (1989). Segmentation of hydrometeorological series. Application to rainfall and flow series in West Africa. Journal of Hydrology 110, pp 349-367.
- 504
- 505 41. Demarée G.R. & Nicolis C. (1990). Onset of Sahelian drought viewed as a fluctuation-induced transition. Quaterly Journal of the Royal Meteorological Society, 116, 221-238.
- 506
- 507 42. Moron V. (1993). Precipitation variability in tropical Africa north of the equator (1933-1990) and relationship with ocean surface temperatures and atmospheric dynamics. Doctoral thesis, University of Burgundy, Climate Research Centres, Dijon, 219 pages.
- 508
- 509
- 510 43. Nicholson S.E. (1993): An overview of African rainfall fluctuations in the last decade. Journal of climate, 6, 1463-1466.
- 511
- 512 44. Soro T. D., Soro N., Oga Y. M. S., Lasm T., Soro G., Ahoussi K. E. & Biémi J. (2011). Climate variability and its impact on water resources in the square degree of Grand-Lahou (South-West Côte d'Ivoire). Physio-Géo - Géographie Physique et Environnement, (5), 55-73. DOI: 10.4000/physiogeog.158
- 513
- 514
- 515
- 516 45. Afouda F., Salako A.P.M. & Yabi I. (2014). Intra-seasonal instability of rains during the main agricultural season in the commune of Kétou in Benin, Revue de Géographie du Laboratoire Leïdi - ISSN 0851 - 2515 - N°12, December 2014
- 517
- 518

why italics?

- 519 46. **Klassou S. D.** (1996) : *Recent climato-hydrological trends and their consequences on the*
520 *environment: the example of the Mono River basin (Togo-Benin)*. PhD thesis, University of
521 Bordeaux III, 472 p.
- 522 47. **Brou Y. T.** (2005). Climate, socio-economic changes and landscapes in Côte d'Ivoire. Thesis on
523 the synthesis of scientific activities presented with a view to obtaining the Habilitation à Diriger des
524 Recherches, Université des Sciences et Techniques de Lille, France, 212p.
- 525 48. **Niasse M., Afouda N. A. & Aman A.** (2004): Reducing West Africa's vulnerability to climate
526 impacts on water resources, wetlands and desertification. Elements of a regional strategy for
527 preparedness and adaptation, 71p.
- 528 49. **Kanohin F., Bachir M.S., Aké G. E. & Savané I.** (2012). Climate variability and coffee and cocoa
529 production in humid tropical areas: the case of the Daoukro region (Central-Eastern Côte d'Ivoire).
530 International Journal of Innovation and Applied Studies, (2)1:194-215
- 531 50. **Kassin K.E., Koné D., Kouamé B., Yoro G.R. & Assa A.** (2008). Rainfall variability and
532 prospects for cocoa replanting in West Central Côte d'Ivoire. Journal of Applied biosciences 12;
533 633-641.
- 534 51. **Kassin K.E., Kouamé B., Coulibaly K., Tahy G. M., N'Guessan W.P., Aka A. R., Assi M. E.,**
535 **Guiraud B. H., Koné B. & Yao G.F.** (2018). Prospects for sustainable cocoa farming from the
536 rainfall balance in the last thirty years at Lôh-Djiboua and Gôh post-pioneers regions, Côte
537 d'Ivoire, (9)11:188-198
- 538 52. **Dibi Kangah P.A. & Mian K.A.** (2016). Agroclimatic analysis of the cocoa zone in Côte d'Ivoire.
539 Journal of Geography of the University of Ouaga I Prof Joseph KI-ZERBO, (2)5:45-68
- 540 53. **Ouédraogo M.** (2001). Contribution to the study of the impact of climate variability on water
541 resources in West Africa. Analysis of the consequences of persistent drought: hydrological
542 standards and regional modelling. PhD thesis, University of Montpellier II, France. 486p.
- 543 54. **Houssou C.M.V., Hounsou M.B., Ulrich S.C.Y.A., Houssou C.S. & K.E. Agbossou** (2016).
544 Rainfall variability and impact on soil water balance under pineapple cultivation, International
545 Journal of Innovation and Applied Studies, (15)4:830-845
- 546 55. **Dékoula C. S., Kouamé B., N'goran E. K., Yao F. G., Ehounou J. N. & Soro N.** (2018). Impact of
547 rainfall variability on the growing season in the cotton production area of Côte d'Ivoire. European
548 Scientific Journal, ESJ, 14(12):143-156
- 549 56. **Camberlin P., Okoola R., Diop M. & Valimba P.** (2003). Identification of rainy season start and
550 end dates: Applications in East Africa and Senegal. *Publication of the International Climate*
551 *Association* (15): 295-303.
- 552 57. **Shoe A.** (1988). Structure of the rainy season in Sudano-Sahelian Africa. Thesis of the Ecole
553 Nationale Supérieure des Mines de Paris, 263 p.
- 554 58. **Balliet R., Saley B.M., Eba E.L.A., Sorokoby V.M., N'Guessan B.H.V., N'dri O.A., Djè K.B. &**
555 **Biémi J.** (2016). Evolution of extreme rainfall extremes in the Gôh Region (Central West Côte
556 d'Ivoire), European Scientific Journal, (12)23 : 74-87
- 557
558
559