

**GROWTH AND VOLUME ESTIMATES OF TEAK (*Tectona grandis* Linn F.) IN
KANYA FOREST PLANTATION, KEBBI STATE, NIGERIA**

ABSTRACT

This study was conducted in order to estimate growth and volume production of Teak (*Tectona grandis*) in Kanya Forest Plantation, Nigeria. The plantation was divided in to six strata-based age classes (A=38, B=37, C=36, D=35, E=34, F=28, years). Five plots were randomly selected from each stratum. Trees within each plot were enumerated and measured. variables measured include total height, diameter at the base, middle, top, and diameter at the breast height were taken from 30 temporary sampled plots of 25x25m approximately from the center, 180 dominant trees were selected from 712 trees. Descriptive statistic was used to summarize the results while inferential statistic(correlation) was used to establish relationship growth and yield variables. Basal area and volume of sampled trees were computed using Excel as well as scatter plots, correlation analysis was achieved using SPSS statistical package version 20. The results of growth and yield values obtained from the dominant trees are (B=249.312m³/ha, D=196.128m³/ha, F=134.976m³/ha, C=119.328m³/ha, E=100.320m³/ ha and A=86.976m³/ha). The results showed that B was(37years) the best and A(38years) was the poorest. The results of correlation showed positive relationships with most of the tree growth and yield characteristics but negative relationships exist between age and some parameters that is to say as the age increases those parameters are decreasing.

Key words: Volume, Site index, Site productivity, Basal area and DBH

1. INTRODUCTION

Forest stand productivity is largely defined as site quality which expresses the growth potential of the species, and it is influenced mainly by forest soils [1, 2]. According to Skovvsgaard and Vanclay [3] Forest Site productivity is a quantitative estimate of the potentials of a given site to produce wood/timber or biomass for a particular species. For instance, site index (SI) or height of the specific population of the dominant and co-dominant trees at reference age is a widely accepted measure of site productivity in forestry [4]. In forestry, site productivity emphasizes the timber or biomass production capability as a major site indicator for site regardless of its ecosystem concept. The concept of site classification has long and rich history in agriculture and forestry. Alternative approaches have been developed for productivity site, depending on the intended purpose. For instance, plant communities or even attribute of single plants have been used as relative indicators of productivity potentials of an ecosystem sometimes refers to as

35 “phytometers”. Site index is an important proxy of site quality and has been used in many
36 conceptual and simulation models of ecosystem dynamics.

37 Continuous depletion of forest resources in Nigeria is on the increase as a result of high demand
38 of wood and wood products, this result in a situation where the resources can no longer meet
39 current demands and the future needs of the teaming population. Consequently, there has been a
40 shift from tropical natural forest management to management of plantation of mainly exotic
41 species in Nigeria [5]. Sustainable forest management require information on the growing stock,
42 such information serves as a guide to the forest managers for evaluating and allocating forest
43 area for exploitation. In timber production, estimations of the growing stock are often expressed
44 in terms of volume, which can be estimated from easily measurable dimensions of the tree [6, 7].
45 In current forest research, the requirement to encompass this new paradigm involves an
46 increasing need for precise estimate of forest structure and biomass, potential productivity or
47 forest growth [8] and modeling on different scales from stand to landscape level. In this regard, a
48 deep knowledge of forest productivity of the state is essential to develop forestry and land use
49 plan and policies [9]. The main objective of this study estimates the volume production of Teak
50 in Kanya Forest Plantation and specifically to determine basal area, volume growth in relation to
51 specific sites and to establish relationships between tree measurable parameters and stand age.


52 Teak (*Tectona grandis* Lf.) occurs naturally only in India, Myanmar, the Loa's People's
53 Democratic Republic and Thailand. It is, however, naturalized in Java and Indonesia [10]. It is
54 also planted throughout tropical Asia, many parts of tropical Africa, and some parts of Latin
55 America [10,11]. Nigeria was the first country outside Asia where teak was introduced between
56 1889 and 1902. [12,13,10,5]. The first teak seed was imported into Nigeria from India while
57 subsequent ones came from Myanmar. The first 750 ha of teak plantation was established in
58 1890 at the Olokemeji forest reserve in the then Western Nigeria, now part of Ogun State 273
59 [5,13,10] There were about 651 ha of teak trees at premier teak plantation site in Nigeria, the
60 Olokemeji forest reserve, alone in 1997 [13]. By the year 2000, there were about 132,500ha in
61 tropical Africa [11].

62 Teak is almost found in all northern states with the exception of few, such as Sokoto Maiduguri
63 and Yobe etc.,.With about 70,000 ha, Nigeria has the largest (52.7%) teak plantation in Africa
64 Common local uses of teak timber include furniture making, joinery and general carpentry
65 works, floor parquet production, flush door manufacturing, as poles for electricity transmission
66 and land telephone lines, as struts in buildings, and as beams in bridge construction [14,15,16].

67 **2.0 MATERIALS AND METHODS**

68 **2.1 The Study Area**

69 The study was conducted in Kanya Forest Plantation in Danko Wasagu Local Government,
70 Kebbi State is located on Latitude 11.339⁰N to 11.348⁰ and Longitude 5.606⁰E to 5.641⁰E,
71 occupying about 4,208km². It is bordered in the South by Sakaba Local Government, in the West
72 by Zuru Local Government both in Kebbi State and in the North by Bukkuyum Local
73 Government Area of Zamfara State. Danko Wasagu has an estimated population of about
74 265,271 people [17]. The vegetation falls under Northern Guinea Savannah. The topography is
75 said to be flat or low land with fertile soil covered by sandy soils, sometime coarse in texture
76 with fadama and alluvial plain suitable for agricultural activities. The weather is marked by
77 single rainy season and long dry season; the average rainfall is 720mm, the rainy season is about
78 four to five months, the mean temperature ranges from 31⁰C and 38⁰C. From the month of
79 November to February cold weather is usually experienced due to the dry harmattan wind and
80 from March to May, the weather is generally hot and wet as in the tropics [18].



81
82 Figure 1. Map of Kanya Forest Plantation

83

84 **2.2 Sampling Procedure**

85 The area was stratified in to different age classes based on the years of establishment (1979,
86 1980, 1981, 1982, 1983, and 1989) on which five temporary sample plots of 25 x 25m
87 (0.0625ha) were marked at random from each age block close to the center. Measurements were
88 taken on all trees within the selected plots. Stand age was obtained from plantation records.

89 **2.3 Data Collection**

90 The data obtained include:

91 Counting and recording of individual trees per plot, Measuring the total height of six dominant
92 trees in all selected plots using Haga Altimeter (this represented the 100 largest trees per ha),
93 Diameter at breast height (DBH) of all individual trees was measured at 1.3m above ground
94 level. Flexible measuring tape was used to determine the circumference of the boles, Diameters
95 at three different points (Base, middle, Top) were determined with the aid of Spiegel Relascope.

96 **2.4 Data Computations and Analysis**

97 The data collected were organized and screened for analysis.

98 Descriptive statistic was used to summarize the results while inferential statistic (correlation) was
99 used to establish relationships between growth and yield variables. Basal area and volume of
100 sampled trees were computed using Excel as well as scatter plots, correlation analysis was
101 achieved using SPSS statistical package version 20.

102 **2.5 Basal area computation**

103 The basal area for each sampled tree was determined using the formula suggested by Husch *et. al*
104 [6]

$$105 \quad BA = \frac{\pi D^2}{4} \quad (1)$$

106 Where: BA = Basal area in m², D = Diameter at breast height (m), π= Pi (3.142)

107 Basal area per plot was obtained by adding the basal area of all individual trees within the plot.

108 Basal area per hectare for each age series was determined by first summing the basal areas of the
109 30 sample plots selected from the age series and finding their mean, then multiplying the mean
110 basal area per plot by the number of sample plots per hectare which is 16.

111 2.5 Volume estimation

112 The stem volume of each mean tree was estimated using the Newton's formula [6]. The formula
113 is expressed as:

$$114 V = \pi h \left(\frac{D_b^2 + 4D_m^2 + D_t^2}{24} \right) \quad (2)$$

115 Where: V = Stem volume in (m³), D_b = Diameter (m) at the base of the tree, D_m = Diameter (m)
116 at the middle of the tree, D_t = Diameter (m) at the top of the tree, h = Total height of the tree (m).

117 3.0 RESULTS

118 3.1 Growth and Yield Variables

119 The data collected include all the individual trees (712) measured from 30 plots selected at
120 random. The parameters computed are summarized and presented in Table 1. In the summary,
121 the mean, minimum, maximum values together with standard error and standard deviation are
122 also presented in order to see the data distribution pattern.

Table 1: Growth and Yield Characteristics/Variables ทำไม้ฐานน้อยกว่าส่วนกลาง

| Variables | Min | Max | Mean | SEM | SD |
|---------------------|-------|-------|-------|-------|-------|
| Db(cm) | 7.1 | 55.7 | 27.9 | 0.24 | 6.44 |
| Dbh(cm) | 6.6 | 48.1 | 22.9 | 0.19 | 5.19 |
| Dm(cm) | 5.5 | 45.0 | 20.6 | 0.19 | 5.08 |
| Dt(cm) | 5.0 | 35.0 | 14.9 | 0.18 | 4.72 |
| H(m) | 4.85 | 28.25 | 12.96 | 0.23 | 3.89 |
| BA(m ²) | 0.01 | 0.94 | 0.25 | 0.01 | 0.17 |
| V(m ³) | 0.060 | 5.190 | 0.716 | 0.024 | 0.651 |

123 Note: Db = Diameter at the base; Dbh = Diameter at the breast height; Dm = Diameter at the middle; Dt = Diameter at the top; H
124 = Height; BA = Basal area and V = Volume; Min = Minimum; Max = Maximum; SEM = standard error of mean and SD =
125 Standard deviation

126 The summaries of growth and yield characteristics of 180 sampled dominant trees are
127 presented in Tables 2 and 3. Mean, minimum and maximum values of Dbh, height, BA and
128 volume are recorded for all the age series. The standard error of the mean was also attached to
129 all the mean values in order to see the variability distribution of the sampled data from the
130 population.

131

132

133

134 **Table 2: Summary Statistics of Dominant Trees (Sampled Trees)**

| Age (years) | Plots | Trees | Dbh (cm) | | | Height(m) | | |
|----------------|-------|-------|----------|-------|------------|-----------|-------|------------|
| | | | Min | Max | Mean* | Min | Max | Mean* |
| 38 | 5 | 6 | 12.51 | 36.98 | 23.77±0.29 | 9.85 | 15.25 | 15.61±0.44 |
| 37 | 5 | 6 | 20.53 | 27.05 | 25.10±0.75 | 11.30 | 19.60 | 15.19±0.51 |
| 36 | 5 | 6 | 19.26 | 37.91 | 26.62±0.93 | 10.70 | 20.00 | 15.58±0.42 |
| 35 | 5 | 6 | 16.23 | 37.91 | 30.07±1.39 | 11.55 | 19.60 | 22.61±0.46 |
| 34 | 5 | 6 | 19.89 | 48.09 | 24.91±0.89 | 18.80 | 28.25 | 15.07±0.39 |
| 28 | 5 | 6 | 16.87 | 39.15 | 25.59±0.41 | 12.90 | 19.80 | 16.06±0.29 |

135 *Mean± standard error

136 **Table 3: Summary of yield characteristics of Dominant Trees (Sites Trees)**

| AC | P | Trees | Basal Area (m ²) | | | Volume (m ³) | | | | |
|----|---|-------|------------------------------|------|-----------|--------------------------|-------|-------|------------|-------------------|
| | | | Min | Max | Mean | Mean BA/ha | Min | Max | Mean | Mean volume/ha |
| A | 5 | 6 | 0.01 | 0.11 | 0.04±0.01 | 4.29 | 0.240 | 0.980 | 0.906±0.04 | 86.976 |
| B | 5 | 6 | 0.03 | 0.16 | 0.50±0.03 | 18.03 | 0.610 | 4.310 | 2.597±0.20 | 249.312 |
| C | 5 | 6 | 0.29 | 1.11 | 0.11±0.04 | 10.34 | 1.260 | 5.630 | 1.243±0.18 | 119.328 |
| D | 5 | 6 | 0.02 | 1.11 | 0.08±0.01 | 7.20 | 0.480 | 5.470 | 2.043±0.18 | 196.128 |
| E | 5 | 6 | 0.03 | 0.18 | 0.05±0.01 | 4.70 | 1.150 | 5.300 | 1.045±0.08 | 100.320 |
| F | 5 | 6 | 0.02 | 0.12 | 0.14±0.02 | 13.14 | 0.580 | 2.290 | 1.406±0.08 | 134.976 |

137 *Mean± standard error

138 3.2 Basal Area and Volume Accumulation at Different Dbh Size Classes

139 Basal area and volume production at different Dbh classes are presented in Table 4. The lowest
 140 and highest basal area were 0.73m² and 174.77m² which was recorded from Dbh class 41-45cm
 141 and 46-50cm, respectively. The lowest and highest volume recorded were 4.46m³ and 509.821m³
 142 from Dbh class (41-45cm and 45-50cm) respectively.

Table 4: Basal Area and Volume Accumulation at Different Dbh Size Classes

| Dbh Class (cm) | Basal Area(m ²) | Volume (m ³) |
|----------------|-----------------------------|--------------------------|
| 05-09 | 4.01 | 6.941 |
| 10-15 | 42.08 | 91.290 |
| 16-20 | 65.46 | 180.803 |
| 21-25 | 44.64 | 152.952 |
| 26-30 | 6.51 | 28.244 |
| 31-35 | 8.30 | 32.371 |
| 36-40 | 2.79 | 12.442 |
| 41-45 | 0.73 | 4.460 |
| 46-50 | 174.77 | 509.821 |

143 **3.3 Basal Area and Volume by Height Classes**

144 Basal area and volume growth based on the height classes are presented in Table 5. The lowest
 145 and highest BA values were 26.54 m² and 174.77m², the lowest and highest volume were
 146 111.310m³ and 509.82m³ recorded from 13-16 m and 25-28 m classes, respectively

Table 5: Basal Area and Volume at Different Height Classes

| Height Classes(m) | Basal Area(m ²) | Volume (m ³) |
|-------------------|-----------------------------|--------------------------|
| 05-08 | 68.81 | 150.140 |
| 09-12 | 58.08 | 185.272 |
| 13-16 | 26.54 | 111,310 |
| 17-20 | 36.81 | 133.331 |
| 21-24 | 27.30 | 113.023 |
| 25-28 | 174.77 | 509.821 |

147 **3.4 Relationship between variables**

148 Table 6. Shows correlation coefficients between tree variables and age of the plantation in which
 149 the relationships between measured variable/parameters were positive and significant, while the
 150 relationship between the age and some variables showed the negative relationship with exception
 151 of basal area and volume which showed positive correlation.


152 **Table 6: Correlation matrix for growth and yield variable of trees in the study area**

| | H(m) | DB(cm) | DBH(cm) | DM(cm) | DT(m) | BA(m ²) | VOL.(m ³) | AGE |
|-----------------------|----------|----------|---------|----------|----------|---------------------|-----------------------|----------|
| H(m) | 1 | 0.457** | 0.556** | 0.606** | 0.591** | 0.181** | 0.529** | -0.188** |
| DB(cm) | 0.457** | 1 | 0.817** | 0.734** | 0.549** | 0.301** | 0.520** | 0.150** |
| DBH(cm) | 0.556** | 0.817** | 1 | 0.853** | 0.683** | 0.432** | 0.601** | -0.096* |
| DM(cm) | 0.606** | 0.734** | 0.853** | 1 | 0.765** | 0.384** | 0.618** | -0.204** |
| DT(m) | 0.591** | 0.549** | 0.683** | 0.765** | 1 | 0.671** | 0.790** | -0.167** |
| BA(m ²) | 0.181** | 0.301** | 0.432** | 0.384** | 0.671** | 1 | 0.846** | 0.141** |
| VOL.(m ³) | 0.529** | 0.520** | 0.601** | 0.618** | 0.790** | 0.846** | 1 | 0.018** |
| AGE | -0.188** | -0.150** | -0.096* | -0.204** | -0.167** | 0.141** | 0.018** | 1 |

153 **Correlation is significant at the 0.01 level (2-tailed) *Correlation is significant at the 0.05 level (2-tailed)


154 Volume distribution in the reserve is said to be more concentrated in trees with Dbh ranging
 155 from 12-32cm with the highest volume accumulation recorded between 0.100-1.200m³. The trees
 156 are said to be sparsely distributed when recording increase in Dbh i.e from 32-50cm, volume
 157 production above 32cm ranges from 1.2-5.0 (Fig 2). Figure 3 presents volume distribution in the

158 reserve based on height and was more concentrated in trees with height ranging from 5.2-20.5m
159 with the highest volume accumulation recorded between 1.000-2.000 m³. The trees were sparsely
160 distributed when recording increase in height i.e. above 20.5m. Figure 4 shows dominant height
161 distribution and was more concentrated in trees with Dbh ranging from 15-30cm and sparsely
162 distributed above 30. Figure 4 shows dominant height and volume distribution of dominant trees.
163 Volumes of dominant trees were found within 0.100-1.200m³ sparsely distributed above 1.200m³



164


165 **Figure 2: Volume accumulation of measured trees at different Dbh**



166

167 **Figure 3: Volume Accumulation of measures trees at Different Height**

168



169

170 **Figure 4. Dominant height of 180 sampled trees based on Dbh.**

171 **4.0 DISCUSSION**

172 **4.1 Growth and yield characteristics**

173 Summary statistics of 180 sampled trees (dominant Dbh and height) were presented depicting
 174 low dbh and height values considering the age of the plantation and were as a result of poor
 175 management. Similar research was conducted by Onyekwelu [19], when Developing Site Index
 176 Curves for Opepe (*Nauclea didderichii*) Plantation in Southwestern Nigeria who reported
 177 slightly higher values of dominant height as well as Dbh, this could be as a result of variation in
 178 the ecosystem and the species involved. Akindele [20] also constructed similar site index curve
 179 for *Tectona grandis* (Teak) in the Dry High Forest Areas of Southwestern Nigeria. The highest
 180 dominant height and dbh reported were less than what was obtained in this study, this could be as
 181 a result of variation in age of the plantation and difference in location, Dominant stand height is a
 182 good predictor of growth, because size is biologically more significant than chronological age as
 183 a causal variable, especially in trees, where meristems are constantly renewed [21].

184 The mean basal area/ha reported in this study is lower than that obtained by Garcia, Mwangi [22,
 185 23], mean volume/ha obtained in this study is said to be higher than what was obtained by [24]
 186 this may be as a result of differences in silvicultural practices, location as well as soil factors in
 187 the study area. [25] reported high range of basal area than that obtained in this study. The low
 188 basal area was as a result of lack of silvicultural management. The findings revealed that the
 189 basal area increases with the increase in age except for the aged teak affected by thinning

190 operations. Many researches on Teak volume were reported by different studies at different age
191 classes, for instance at the age of 16 years the volume reported by [26] is far better than the value
192 reported in this study, this variation might have been influenced by climate variability, rainfall as
193 well as soil fertility of the site. [27] reported 40 years old Teak produced volume less than the
194 value obtained in this research. The appropriate method of quantifying volume of a stand is
195 necessary at different age classes and site because volume differ with location, silvicultural
196 activities, site classes and age. Tree volume provides valuable information on supply of both
197 industrial wood and hence identifying sustainable management of forests and woodland
198 ecosystems [28, 29]. Dbh classes 40-45cm, 36-40 and 05-09 recorded lower basal area and
199 volume which could be attributed to fewer number of stems compared to other Dbh classes. In
200 this research, the summation of volume of the second and the third Dbh class was less than that
201 presented by [30] for the same specie which they obtained from similar Dbh class, this might be
202 as a result of climatic variability, site, soil as well as silvicultural operations involved. [31] in
203 Northern Thailand reported similar Dbh class which disagreed with this research.

204 **4.2 Relationships between growth variables**

205 Pearson correlation analysis of the stand variables with age revealed that, there was high
206 association between tree characteristics such as diameter at the breast height, height as well as
207 volume. Plantation ages revealed negative relationship with the rest of the variables with the
208 exception of basal area and volume growth which showed positive relationship. There was
209 significant and positive correlation with most of the tree growth and yield characteristics, this
210 coincides with the findings of [32] and [33]. For instance, tree height-DBH, height-volume,
211 DBH-volume and basal area-volume displayed a positive correlation. Also, correlation analysis
212 was observed by [34] in Developing Site Index Equation and Curves for Site Quality Assessment
213 of *Pinus Caribea* Monoculture Plantation in South Western Nigeria. They discovered a high
214 linear relationship between tree age and other growth characteristics such as Dbh, total height,
215 and merchantable height as well as slenderness coefficient, these varies with association
216 displayed by age and other parameters in this research, Dbh, Height, Db, Dm, Dt showed
217 negative relationships and this indicates that as they approached that age (plantation age) these
218 parameters decreases. Appropriate silvicultural treatment such as thinning and pruning be done
219 on regular basis to avoid unnecessary nutrient uptake competition.

220

221 **CONCLUSION**

222 Growth and yield production of *Tectona grandis* was investigated in this research. Basal area of
223 sampled trees are as follows according to magnitude B=18.03m²/ha, F=13.14m²/ha,
224 C=10.34m²/ha, D=7.20m²/ha, E=4.72m²/ha, A=4.29m²/ha with B having the highest and the
225 lowest. The yield values obtained from the dominant trees are (B=249.312m³/ha,
226 D=196.128m³/ha, F=134.976m³/ha, C=119.328m³/ha, E=100.320m³/ ha and A=86.976m³/ha).
227 Conclusively site B was (37years) as the best site for *Tectona grandis* and A(38years) was the
228 poorest which is as a result of soil variations within the study site. The results of correlation
229 showed positive relationships with most of the tree growth and yield characteristics but negative
230 relationships exist between age and some parameters

231 **REFERENCES**

- 232 1. Skovsgaard J, Vanclay JK. Forest site productivity: a review of the evolution of dendrometric
233 concepts for even- aged stands. *Forestry*; 81(1): 13-31.
- 234 2. Avery TE, Burkhart HE. Forest measurements. Fifth Edition. Waveland Press Inc. New York,
235 US; 2015
- 236 3. Vanclay JK. Modelling forest growth and yield: Applications to mixed tropical forests.
237 Wallingford, UK: CAB International; 1994.
- 238 4. Kayahara GJ, Klinka K, Marshall PL. Testing site index-site-factor relationship for predicting
239 *Pinus contorta* and *Picea engelmannii* _ *P. glauca* productivity in central British Columbia,
240 Canada. *Forest Ecology and Management*. 1998; 110 (1–3):141– 150.
- 241 5. Formecu. Nigerian Forestry Statistics. Compiled by Khalique-ur-Rahman and Auofor, RO.
242 Forestry projects monitoring and evaluation unit, Federal Department of Forestry, Ibadan,
243 Nigeria. FOR/MEU/STAT/PUB/NO.2. 1991.
- 244 6. Husch B, Beers TW, Kersaw JA. Forestry inventory,150–160 in *Forest Mensuration*, 4th Ed.
245 John Wiley & Sons, Hoboken, NJ; 2003.
- 246 7. Akindele SO. Volume Prediction from Stump Diameters of *Gmelina arborea* (Roxb trees in
247 Akure forest reserve, Nigeria. *Nigerian Journal of Forestry*. 2003;33(2): 116–23.
- 248 8. Tickle PK, Coops NC, Hafner SD, Bago Science Team. Assessing forest productivity at local
249 scales across a native eucalypt forest using a process model, 3PG-SPATIAL. *Forest Ecology and*
250 *Management*. 2001;152: 275–291.
- 251 9. Landsberg J. Modelling forest ecosystems: state of the art, challenges and future directions.
252 *Canadian Journal of Forest Research*. 2003;33: 385–397. 2003
- 253 10. Pandey D, Brown C. Teak: a global overview. *Unasylva*.2000; 51:18-26
- 254 11. Krishnapillay B. Silviculture and management of teak plantations. *Unasylva: An*
255 *International Journal of Forestry and Forest Industries*.2000; 51:14-2

- 256 12. Home JEM. Teak in Nigeria. Nigerian Information Bulletin (New Series) No. 16.1966
- 257 13. Oluwalana SA. An economic assessment of the existing teak and *Gmelina* plantations in
258 Ogun State, Nigeria. The Nigerian Journal of Forestry.1997; 27(2):40-47.
- 259 14. Omole AO. Small diameter teak (*Tectana grandis*) processing for export market and Its
260 effects on national economy. Paper presented in the Seminar, Department of
261 Agricultural Engineering, University of Ibadan, Nigeria.1996:18
- 262 15. Ezekiel OA. The effects of fuelwood removals from Gambari forest reserve teak plantation
263 on the environment. A Seminar Paper (Unpublished) Presented in the Department of Agricultural
264 Engineering, University of Ibadan, Nigeria. 1997:17
- 265 16. Areghan SE. sawmilling industry in Nigeria: a case study of the Industrial Development Unit
266 of Forestry Research Institute of Nigeria. A Seminar Paper (Unpublished) Presented in the
267 Department of Agricultural Engineering, University of Ibadan, Nigeria. 2001:24
- 268 17. National Population Commission. *Provisional Census Figure*. Abuja Nigeria. 2006:1-3.
- 269 18. Girma SA. Agro-climatology of Millet Production in Desert Fringe Zone of Nigeria, A Case
270 Study of Kebbi State. M.Sc. dissertation.; Federal University of Technology Minna, Niger
271 state.2008;1-97.
- 272 19. Onyekwelu JC. Site index curves for site quality assessment of *Nauclea diderrichii*
273 monoculture plantations in Omo forest reserve, Nigeria. Journal of Tropical Forest Science.
274 2005;17(4): 532-542.
- 275 20. Akindele SO. Development of a site index equation for teak plantations in south-western
276 Nigeria. Journal of Tropical Forest Science.1991;4(2):162–169.
- 277 21. García O. A simple and effective forest stand mortality model. Mathematical and
278 Computational Forestry & Natural-Resource Sciences. 2009;1(1):1–9.
- 279 22. Mwangi RA. Volume and biomass estimation models for *Tectona grandis* grown at longuza
280 forest plantation, Tanzania. M.Sc. Dissertation University of Agriculture. Morogoro,
281 Tanzania.2015.
- 282 23. Ige PO, Akinyemi GO. Site Quality Assessment for *Tectona grandis* Linn.f Plantations in
283 Gambari Forest Reserve, Nigeria. Journal of Forestry Research and Management. 2015; 12:58-
284 67.
- 285 24. Sunanda C, Jarayaman KM. Prediction of Stand attributes of even aged Teak Stands using
286 multilevel models. Forest Ecology and Management. 2006; 236:1–11.
- 287 25. KFRI. Volume distribution for *Tectona grandis* L.f in Kerala. Kerala Research Report.
288 [<http://docs.kfri.res.in/KFRI-RR/KFRI-RR201.pdf>] December, 2011.
- 289 26. Chambers JQ, Santos J, Ribeiro RJ, Higuchi N. Tree damage, allometric relationships and
290 aboveground net primary production in a tropical forest. Forest Ecology and Management. 2001;
291 152:73–84.

- 292 27. Mugasha WA, Eid T, Bollandås O.M, Malimbwi RE, Chamshama SAO, Zahabu E, Katani
293 JZ. *Allometric models for prediction of aboveground biomass of single trees in miombo*
294 *woodlands in Tanzania*. The Proceedings of the First Climate Change Impact Adaptation and
295 Mitigation (CCIAM) Programme Scientific Conference, Blue Pearl Hotel, Dar Es Salaam,
296 Tanzania 2nd and 3rd January. 2001;8-17.
- 297 28. Prasad V, Lalnundaga K, Iremkini JV. Growing stock variation in different *teak* (*Tectona*
298 *grandis*) *forest stands of Mizoram, India*. Journal of Forestry Research. 2008;19(3): 204-208.
- 299 29. Gajasemi J, Jordan C. Decline of Teak Yield in Northern Thailand: Effects of Selective
300 Logging on Forest Structure. BIOTROCA.1990;22(2): 114-118.
- 301 30. Adekunle VAJ. Nonlinear regression models for timber volume estimation in natural forest ecosystem
302 southwest, Nigeria. Research Journal of Forestry. 2007;1(2): 40-54 doi: 103923/rjf. 2007. 40.54.
- 303 31. Adeyemi AA. Site Quality Assessment and Allometric Models for Tree Species in Orban
304 Forest, Nigeria. Journal of Sustainable forestry. 2016;36(4): 280-289.
- 305 32. Oyebade BA, Osho JS, Adesoye PO. Development of Site Index Equations for Site Quality
306 Assessment of *Pinus Carribea* Monoculture Plantation in South Western Nigeria. Journal of
307 Forestry and Environment and Environmental Science. 2014;30(40): 215-321.