

Severity of Asthma and Serum Vitamin D Levels in Asthmatic Children

ABSTRACT

Introduction Vitamin D deficiency has been rediscovered as a public health problem worldwide. It has been postulated that vitamin D deficiency explains a portion of the asthma epidemic. The purpose of this study was to assess the serum vitamin D levels in children with asthma as compared with the non-asthmatic population and to investigate the association of serum vitamin D levels with the severity of asthma.

Methods & Materials We measured serum 25-hydroxyvitamin D (25-OH vitamin D) levels in 50 children with mild intermittent to moderate persistent asthma at the time of enrollment and 50 age- and sex-matched non-asthmatic children in a case-control study. The independent sample t-test, χ^2 test, and spearman correlation coefficient were used to analyze the data.

Results Vitamin D level was 13.6 ± 1.1 ng/ml and 19.2 ± 1.8 ng/ml in asthmatic and non-asthmatic individuals, respectively. The mean (\pm SD) levels of serum vitamin D were statistically significant between asthmatic and non-asthmatic individuals. Females had lower levels of vitamin D than males. Asthma severity was inversely associated with serum vitamin D level.

Conclusion Findings of the present study demonstrate the significance of vitamin D in asthma. Sufficient doses of vitamin D should be administered to pregnant mothers and the babies monitored for symptoms of wheezing or asthma during childhood. Clinical trials are needed to definitively answer questions about the role of vitamin D in asthma.

Keywords: Asthma, Vitamin D, Vitamin D deficiency, 25-OH vitamin D

1. INTRODUCTION

Asthma is the most prevalent chronic respiratory disease in all age groups, affecting 300 million people worldwide. In children, asthma is the most common chronic disease. The prevalence of asthma has been increased in recent years. According to ISAAC global asthma prevalence raised from 11.1 to 11.6% in children and 13.2 to 13.7% in adolescents [1].

In Iran, the prevalence of asthma among children under 18 years of age varies from 1.26% to 11.6% depending upon sex, ethnicity, geography, and other factors [2]. Many factors can influence the severity of asthma, such as viral respiratory infections, male gender, food allergy, atopy, and air pollution. The role of these factors has been clearly proven over the past years [3]. Some recent studies have been conducted to determine the role of some vitamins and microelements in asthma occurrence and prevention [4-6].

Among the vitamins, the roles of vitamins C and D in the occurrence of asthma have been studied more seriously. Low levels of vitamin C and α -carotene are believed to be risk factors for asthma occurrence [5]. According to Chambers and Hawrylowicz, supplementary vitamin D had an anti-inflammatory effect on the lungs and also increased the level of IL10, an anti-inflammatory mediator [7]. Vitamin D has a regulatory role in other inflammatory processes and diseases [8]. Some studies showed that low levels of vitamin D increase mortality in patients with breast and lung cancers [9-10]. Low levels of vitamin D are also correlated with some autoimmune disorders, such as type I diabetes mellitus [11] multiple sclerosis [12], rheumatoid arthritis [13], and inflammatory bowel diseases [14].

34 Vitamin D has an important role in the suppression of the inflammatory response of Th2 cells
35 in the lungs. In animal models, it can induce surfactant synthesis and stimulate lung
36 maturation [15]. The role of vitamin D and the synthesis of surfactant in the human fetus is
37 more complex [16]. The risk of wheezing can be lowered in infants whose mothers have
38 taken high doses of vitamin D during pregnancy [17].

39 Some studies have shown that vitamin D can be effective in regulating immune responses
40 and can affect fetal lung growth [18]. Low levels of maternal serum vitamin D during
41 pregnancy can increase the risk of occurrence of asthma during childhood [19]. Vitamin D
42 supplementation can reduce asthma attack requiring systemic corticosteroids for treatment
43 [20]. However, some studies have presented contradictory results regarding the relationship
44 between serum vitamin D levels and allergic diseases. Taking supplementary vitamin D can
45 increase allergic diseases in children and an increased risk of asthma and allergies with a
46 high level of vitamin D was reported [21]. This study was designed to determine and
47 compare the levels of serum vitamin D in children with asthma as compared with those of a
48 control group. Moreover, the relationship between the severity of asthma and the level of
49 serum vitamin D was investigated.

50
51

52 **2. MATERIAL AND METHODS**

53

54 In this cross-sectional case-control study, 50 asthmatic children, aged 2–18 years, were
55 enrolled via simple random sampling. The patients were referred to the Asthma and Allergy
56 Clinic of Afzalipour Medical Center of Kerman University of Medical Sciences. Asthma was
57 diagnosed and classified according to the National Asthma Education and Prevention
58 Program's Expert Panel Report 3 [22].

59 Patients who suffered from disorders interfering with vitamin D metabolism or kidney disease
60 and those who were taking antiepileptic drugs, supplementary vitamin D or other vitamins
61 were excluded from the study. Fifty participants referring to the out-patient pediatric clinic of
62 Afzalipour Medical Center due to simple and uncomplicated disorders, e.g., common cold
63 and otitis media were selected as the control group. For each case, one control subject was
64 randomly selected and matched to the index case according to age and gender. For each
65 person, a data collection sheet containing a name, age, gender, asthma severity, and other
66 demographic data was filled out. The study was approved by the Research Ethics
67 Committee of the University, and written informed consent was obtained from all cases and
68 controls. A 5-mL sample of peripheral venous blood was collected from each individual. After
69 the separation of serum by centrifugation at 2000 rpm, the level of 25-OH vitamin D was
70 measured using a commercial ELISA kit (DLD Diagnostica, GmbH) according to the
71 manufacturer's instructions. Serum levels of 25 (OH) vitamin D were divided into three
72 groups, including deficient (<20 ng/ml), insufficient (20–30 ng/ml), and sufficient (\geq 30 ng/ml)
73 [23-24]. SPSS v.15 (SPSS Inc., Chicago, IL.) was used for statistical analysis of data. The
74 independent sample t-test and chi-square analysis were used for comparison of the two
75 groups. The Spearman correlation coefficient was calculated to identify the correlation
76 between serum 25-OH vitamin D level and severity of asthma. $P < 0.05$ was considered to
77 indicate statistical significance.

78

79 **3. RESULTS**

80

81 Vitamin D level was 13.6 ± 1.1 ng/ml and 19.2 ± 1.8 ng/ml in asthmatic and non-asthmatic
82 individuals, respectively. The mean (\pm SD) levels of serum vitamin D were statistically
83 significant between asthmatic and non-asthmatic individuals ($P = 0.01$). Amongst all
84 participants, 52% of cases ($n = 26$) and 62% ($n = 31$) of controls were male. The mean
85 (\pm SD) age was 6.8 ± 0.4 years for cases and 7.6 ± 0.5 years for the controls. The serum
86 vitamin D levels were lower in patients with moderate persistent asthma than in other

87 patients (Table 1), but this difference was not statistically significant. However, there was a
 88 significant negative correlation between serum vitamin D level and asthma severity ($r =$
 89 0.242 , $P = 0.015$).

90
 91
 92

Table 1. Serum vitamin D levels in asthmatic and non-asthmatic individuals according to age

Age	Vitamin D level (ng/mL) Asthmatics	Vitamin D level (ng/mL) Non-asthmatics	P value
Below 5 years	17.4 ± 1.3	25.7 ± 3.8	0.05*
5–10 years	11.5 ± 1.6	15.7 ± 2.1	0.1
Above 11 years	10.3 ± 3.7	16.1 ± 3.1	0.27

93

94 The serum vitamin D levels were 10.6 ± 1.4 ng/ml and 16.3 ± 1.4 ng/ml in female and male
 95 individuals, respectively. The difference between the genders was found to be statistically
 96 significant ($P = 0.009$). Serum vitamin D levels were compared in asthmatic and non-
 97 asthmatic individuals according to age (Table 2). The level of vitamin D was significantly
 98 lower in asthmatic patients under 5 years ($P < 0.05$). However, in older ages, no significant
 99 difference was demonstrated between the two groups. In the case group, children >11 years
 100 had the lowest serum vitamin D levels (10.3 ± 3.7 ng/ml) and children <5 years had the
 101 highest levels (17.4 ± 1.3 ng/ml), a difference that was statistically significant ($P = 0.02$). This
 102 difference was also found to be significant in the control group ($P = 0.04$).

103 Serum vitamin D levels and asthma severity according to National Asthma Education and
 104 Prevention Program's Expert Panel Report (4) shows that a significant ($p=0.04$) relation
 105 between Mild intermittent (15.1 ± 2), Mild Persistent (13.6 ± 1.4) and Moderate persistent
 106 (10.2 ± 3.8).

107 Serum vitamin D levels according to the type of dwelling and childcare situation are shown in
 108 Table 3. The serum vitamin D levels in home-cared children were significantly different in
 109 asthmatic (3.2 ± 1.3 ng/mL) and non-asthmatic subjects (20.9 ± 2.1 ng/mL, $P=0.004$).

110
 111
 112

Table 3. Serum vitamin D level according to childcare situation, type of dwelling, maternal age, mother's education and history of taking vitamin D during infancy.

	Case	Control	P value
Childcare situation			
Kindergarten	14.8 ± 1.8	13.2 ± 2.8	0.65
Home	13.2 ± 1.3	20.9 ± 2.1	0.004
P value	0.5	0.04*	
Type of dwelling			
Villa building	14.1 ± 1.2	19.7 ± 2	0.02
Apartment	10.9 ± 2.6	15.3 ± 1.7	0.23
P value	0.3	0.4	
Taking vitamin D during infancy			
Complete	13.6 ± 1.5	18.3 ± 2.8	0.16

Incomplete	12.9 ± 1.9	17.3 ± 3	0.21
Not taking	15.1 ± 3.1	24.1 ± 2.6	0.046*
P value	0.8	0.4	

113

114

115

116

117

118

119

120

121

122

123

124

125

126

There was statistically significant difference in serum vitamin D levels between asthmatic and non-asthmatic subjects in terms of the type of dwelling. Significantly lower vitamin D concentrations were detected in the asthmatic children living in villa buildings than the control group (P=0.02). However, no significant difference was documented in those living in apartments.

The mean level of vitamin D (95%CI) was measured according to the parental history of providing supplementary doses of vitamin D to children (Table 3). Higher levels of vitamin D were shown in non-asthmatic children who were not taking supplementary vitamin D compared to children with asthma (P=0.046).

4. DISCUSSION

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

This study shows that the serum 25-OH vitamin D level in asthmatic children was lower than those in the controls. The role of vitamin D in the pathogenesis of asthma is not fully understood. Vitamin D can influence immune responses by affecting T-helper type 1 (Th1) and T-helper type 2 (Th2) functions [25]. Polymorphisms in vitamin D receptor-coding genes can correlate with different phenotypes of asthma [26]. There was a significant negative correlation between the serum 25-OH vitamin D level and severity of asthma in this study. Alyasin et al showed the serum 25-OH D3 levels and childhood asthma severity are inversely associated and suggested a direct relationship between pulmonary function test results and vitamin D deficiency in Iranian asthmatic children [27]. In Qatar, with a similar social and climate conditions to Iran, Bener et al found asthmatic children had significantly reduced levels of serum vitamin D compared to the non-asthmatic controls [28]. Brehm et al showed that the severity of asthma and increased markers of allergy are correlated with a low level of serum vitamin D [28].

Litonjua shows that the use of supplementary vitamin D in the diet can play a primary preventive role in the incidence of asthma, reduce the severity of asthma, and improve recovery from steroid-resistant asthma. In this study, there was no significant difference in the serum vitamin D levels in children with a history of taking sufficient amounts of vitamin D in the supplemental form with those who did not take supplementary vitamin D (at least 400 IU per day) during infancy. Failing to remember vitamin D supplementation by mothers is a factor that should be taken into consideration. Studies showed that children who have taken supplementary vitamin D during the first year of life have a lower risk of asthma at 31 years of age, but there was no report of vitamin D levels during this period [29]. A meta-analysis of randomized trials shows the rate of asthma exacerbation can reduce by vitamin D supplementation especially in patients with vitamin D insufficiency [30]. Litonjua (2019) suggested vitamin D supplementation can prevent asthma and wheeze in early life and may help in the treatment of asthma [31].

Gale shows that the prevalence of asthma at the end of the first decade of life has a negative correlation with the serum vitamin D levels of mothers in the third trimester of pregnancy [32]. A deficiency of vitamin D in the prenatal period can influence the development of the fetal lungs and immune system [18].

This study shows that serum vitamin D levels were lower in female than in male patients, this difference may be due to body coverage in female patients. In one study conducted in Turkey, the lowest serum vitamin D levels were found in women who cover their whole body from sunlight [33]. Gender is a significant factor in vitamin D insufficiency as indicated in our

161 findings as well as in a study carried out in Tehran by Rabbani et al. They found the
162 prevalence of vitamin D insufficiency in healthy subjects was 53.6% in girls and 11.3% in
163 boys [24]. A study in Saudi Arabia showed the levels of vitamin D in asthmatic and non-
164 asthmatic females was lower than males which are by the results of the present study [34].
165 Despite higher sunny days in the region throughout the year, these findings can be attributed
166 to the body coverage of females due to the social, cultural and behavioural aspects of this
167 issue.

168 Mirsaeid Ghazi et al showed that children under 10 years of age have higher serum vitamin
169 D levels in contrast to older age groups and the level of serum vitamin D decreases with
170 increasing age [35]. These data are in accordance with our findings. The difference in the
171 serum vitamin D levels in the early years of life in contrast to later years can be due to more
172 attention given by parents to vitamin D supplementation of children in the early years of life
173 and more exposure to sunlight due to more time spent on outdoor activities by this age
174 group.

175 It has been shown that low levels of serum vitamin D are correlated with more severe forms
176 of asthma. Brehm demonstrated that vitamin D insufficiency can develop more severe forms
177 of asthma exacerbation in Puerto Rican children [36]. Tabak et al reported that using some
178 foods, such as fish, containing high amounts of vitamin D can protect children from asthma
179 [37]. Vitamin D deficiency is common in different parts of Iran [38]. Twenty-six percent of
180 primary school children in Isfahan had vitamin D deficiency [39]. In Shiraz 81.3% of healthy
181 children were vitamin D efficient [40]. In Zahedan, southeast of Iran, vitamin D deficiency
182 was reported in 94.7% of apparently health subjects [41]. High prevalence of vitamin D
183 deficiency may be due to the low dietary intake of vitamin D, reduced sunlight exposure and
184 low physical activity [40-41].

185 In conclusion the findings of the present study demonstrate the significance of vitamin D in
186 asthma. The level of serum vitamin D in asthmatic patients was significantly lower than those
187 in controls. The severity of asthma had a significant negative correlation with serum vitamin
188 D levels. Further studies are required to determine the role of vitamin D in the prevention of
189 asthma and in decreasing its severity. To this end, sufficient doses of vitamin D should be
190 administered to pregnant mothers and the babies monitored for symptoms of wheezing or
191 asthma during childhood in future years.

192 REFERENCES

- 193
- 194 1. Ferrante G., La Grutta SL. The burden of pediatric asthma. *Frontiers in pediatrics.*
195 2018;6:186.
- 196 2. Hassanzadeh J, Mohammadbeigi A, Mousavizadeh A, Akbari M. Asthma prevalence in
197 Iranian guidance school children, a descriptive meta-analysis. *Journal of Research in*
198 *Medical Sciences.* 2012;17(3):293.
- 199 3. Liu AH, Covar RA, Spahn JD, Sicherer SH. Childhood Asthma, In: KliegmanRM, Stanton
200 BF, St. Geme JW, Schor NF. *Nelson Textbook of Pediatrics.* 2016, Philadelphia, PP:1095-6.
- 201 4. Somashekar AR, Prithvi AB, Gowda MV. Vitamin D levels in children with bronchial
202 asthma. *Journal of Clinical and Diagnostic Research: JCDR.* 2014;8(10):PC04.
- 203 5. Radia I, Denisc H-than, Wise MR, Serum vitamin levels and the Risk of Asthma in
204 children. *AM Journal Epidemiol* 2004; 159: 351-7.
- 205 6. Brehm JM, Celedón JC, Soto-Quiros ME, Avila L, Hunninghake GM, Forno E, Laskey D,
206 Sylvia JS, Hollis BW, Weiss ST, Litonjua AA. Serum vitamin D levels and markers of severity
207 of childhood asthma in Costa Rica. *American journal of respiratory and critical care*
208 *medicine.* 2009;179(9):765-71.
- 209 7. Chambers ES, Hawrylowicz CM. The impact of vitamin D on regulatory T cells. *Current*
210 *allergy and asthma reports.* 2011;11(1):29-36.
- 211 8. May E, Asadullah K, Zugel U. Immunoregulation through 1, 25-dihydroxyvitamin D 3 and
212 its analogs. *Current Drug Targets-Inflammation & Allergy.* 2004;3(4):377-93.

- 213 9. Crew KD, Gammon MD, Steck SE, Hershman DL, Cremers S, Dworakowski E, Shane E,
214 Terry MB, Desai M, Teitelbaum SL, Neugut AI. Association between plasma 25-
215 hydroxyvitamin D and breast cancer risk. *Cancer Prevention Research*. 2009;2(6):598-604.
- 216 10. Zhou W, Heist RS, Liu G, Asomaning K, Neuberger DS, Hollis BW, Wain JC, Lynch TJ,
217 Giovannucci E, Su L, Christiani DC. Circulating 25-hydroxyvitamin D levels predict survival in
218 early-stage non-small-cell lung cancer patients. *Journal of Clinical Oncology*.
219 2007;25(5):479-85.
- 220 11. Hyppönen E, Läärä E, Reunanen A, Järvelin MR, Virtanen SM. Intake of vitamin D and
221 risk of type 1 diabetes: a birth-cohort study. *The Lancet*. 2001;358(9292):1500-3.
- 222 12. Munger KL, Levin LI, Hollis BW, Howard NS, Ascherio A. Serum 25-hydroxyvitamin D
223 levels and risk of multiple sclerosis. *JAMA*. 2006;296(23):2832-8.
- 224 13. Merlino LA, Curtis J, Mikuls TR, Cerhan JR, Criswell LA, Saag KG. Vitamin D intake is
225 inversely associated with rheumatoid arthritis: results from the Iowa Women's Health Study.
226 *Arthritis & Rheumatism*. 2004;50(1):72-7.
- 227 14. Cantorna MT. Vitamin D and its role in immunology: multiple sclerosis, and inflammatory
228 bowel disease. *Progress in Biophysics and Molecular Biology*. 2006;92(1):60-4.
- 229 15. Nguyen TM, Guillozo H, Marin L, Tordet C, Koite S, Garabedian M. Evidence for a
230 vitamin D paracrine system regulating maturation of developing rat lung epithelium.
231 *American Journal of Physiology-Lung Cellular and Molecular Physiology*. 1996;271(3):L392-
232 9.
- 233 16. Phokela SS, Peleg S, Moya FR, Alcorn JL. Regulation of human pulmonary surfactant
234 protein gene expression by 1 α , 25-dihydroxyvitamin D₃. *American Journal of Physiology-
235 Lung Cellular and Molecular Physiology*. 2005;289(4):L617-26.
- 236 17. Camargo Jr CA, Rifas-Shiman SL, Litonjua AA, Rich-Edwards JW, Weiss ST, Gold DR,
237 Kleinman K, Gillman MW. Maternal intake of vitamin D during pregnancy and risk of
238 recurrent wheeze in children at 3 y of age. *The American journal of clinical nutrition*.
239 2007;85(3):788-95.
- 240 18. Litonjua AA. Childhood asthma may be a consequence of vitamin D deficiency. *Current
241 Opinion in Allergy and Clinical Immunology*. 2009 Jun;9(3):202.
- 242 19. Litonjua AA, Weiss ST. Is vitamin D deficiency to blame for the asthma epidemic?.
243 *Journal of Allergy and Clinical Immunology*. 2007;120(5):1031-5.
- 244 20. Jolliffe DA, Greenberg L, Hooper RL, Griffiths CJ, Camargo Jr CA, Kerley CP, Jensen
245 ME, Mauger D, Stelmach I, Urashima M, Martineau AR. Vitamin D supplementation to
246 prevent asthma exacerbations: a systematic review and meta-analysis of individual
247 participant data. *The Lancet Respiratory Medicine*. 2017;5(11):881-90.
- 248 21. Litonjua AA. Vitamin D deficiency as a risk factor for childhood allergic disease and
249 asthma. *Current opinion in allergy and clinical immunology*. 2012;12(2):179.
- 250 22. National Asthma Education and Prevention Program. Expert Panel Report 3: Guidelines
251 for the diagnosis and management of asthma. NIH pub no 07-4051. Bethesda, MD: National
252 Heart, Lung, and Blood Institute, National Institutes of Health. 2007. Available from:
253 <http://www.nhlbi.nih.gov/guidelines/asthma/>.
- 254 23. Thomas GO, Tutar E, Tokuc G, Oktem S. 25-hydroxy Vitamin D Levels in Pediatric
255 Asthma Patients and its Link with Asthma Severity. *Cureus*. 2019;11(3):e4302.
- 256 24. Rabbani A, Alavian SM, Motlagh ME, Ashtiani MT, Ardalan G, Salavati A, Rabbani B,
257 Rabbani A, Shams S, Parvaneh N. Vitamin D insufficiency among children and adolescents
258 living in Tehran, Iran. *Journal of tropical pediatrics*. 2008;55(3):189-91.
- 259 25. Mahon BD, Wittke A, Weaver V, Cantorna MT. The targets of vitamin D depend on the
260 differentiation and activation status of CD4 positive T cells. *Journal of cellular biochemistry*.
261 2003;89(5):922-32.
- 262 26. Raby BA, Silverman EK, Lazarus R, Lange C, Kwiatkowski DJ, Weiss ST. Chromosome
263 12q harbors multiple genetic loci related to asthma and asthma-related phenotypes. *Human
264 molecular genetics*. 2003;12(16):1973-9.

- 265 27. Alyasin S, Momen T, Kashef S, Alipour A, Amin R. The relationship between serum 25
266 hydroxy vitamin d levels and asthma in children. *Allergy, asthma & immunology research*.
267 2011;3(4):251-5.
- 268 28. Bener A, Ehlayel MS, Tulic MK, Hamid Q. Vitamin D deficiency as a strong predictor of
269 asthma in children. *International Archives of Allergy and Immunology*. 2012;157(2):168-75.
- 270 29. Hyppönen EL, Sovio U, Wjst M, Patel S, Pekkanen J, Hartikainen AL, Järvelinb MR.
271 Infant vitamin d supplementation and allergic conditions in adulthood: northern Finland birth
272 cohort 1966. *Annals of the New York Academy of Sciences*. 2004;1037(1):84-95.
- 273 30. Wang M, Liu M, Wang C, Xiao Y, An T, Zou M, Cheng G. Association between vitamin D
274 status and asthma control: a meta-analysis of randomized trials. *Respiratory Medicine*.
275 2019;150:85-94.
- 276 31. Litonjua AA. Vitamin D and childhood asthma: causation and contribution to disease
277 activity. *Current Opinion in Allergy and Clinical Immunology*. 2019;19(2):126-31.
- 278 32. Gale CR, Robinson SM, Harvey NC, Javaid MK, Jiang B, Martyn CN, Godfrey KM,
279 Cooper C. Maternal vitamin D status during pregnancy and child outcomes. *European*
280 *Journal of Clinical Nutrition*. 2008;62(1):68.
- 281 33. Alagöl F, Shihadeh Y, Boztepe H, Tanakol R, Yarman S, Azizlerli H, Sandalci Ö. Sunlight
282 exposure and vitamin D deficiency in Turkish women. *Journal of Endocrinological*
283 *Investigation*. 2000;23(3):173-7.
- 284 34. Al-Daghri NM, Al-Attas OS, Yakout SM, Alnaami AM, Wani K, Alokail MS. The
285 association of serum 25-OH vitamin D with asthma in Saudi adults. *Medicine*. 2018;97(36).
- 286 35. Ghazi AM, Zadeh FR, Pezeshk P, Azizi F, Cacicedo L. Seasonal variation of serum 25
287 hydroxy D3 in residents of Tehran. *Journal of Endocrinological Investigation*.
288 2004;27(7):676-9.
- 289 36. Brehm JM, Acosta-Pérez E, Klei L, Roeder K, Barmada M, Boutaoui N, Forno E, Kelly R,
290 Paul K, Sylvia J, Litonjua AA. Vitamin D insufficiency and severe asthma exacerbations in
291 Puerto Rican children. *American journal of respiratory and critical care medicine*.
292 2012;186(2):140-6.
- 293 37. Tabak C, Wijga AH, de Meer G, Janssen NA, Brunekreef B, Smit HA. Diet and asthma in
294 Dutch school children (ISAAC-2). *Thorax*. 2006;61(12):1048-53.
- 295 38. Heshmat R, Mohammad K, Majdzadeh SR, Forouzanfar MH, Bahrami A, Ranjbar
296 Omrani GH. Vitamin D deficiency in Iran: A multi-center study among different urban areas.
297 *Iran Journal of Public Health*. 2008;37(1):72-8.
- 298 39. Ardestani PM, Salek M, Keshteli AH, Nejadnik H, Amini M, Hosseini SM, Rafati H,
299 Kelishadi R, Hashemipour M. Vitamin D status of 6-to 7-year-old children living in Isfahan,
300 Iran. *Endokrynologia Polska*. 2010;61(4):377-82.
- 301 40. Saki F, Dabbaghmanesh MH, Omrani GR, Bakhshayeshkaram M. Vitamin D deficiency
302 and its associated risk factors in children and adolescents in southern Iran. *Public Health*
303 *Nutrition*. 2017;20(10):1851-6.
- 304 41. Kaykhaei MA, Hashemi M, Narouie B, Shikhzadeh A, Rashidi H, Moulaei N, Ghavami S.
305 High prevalence of vitamin D deficiency in Zahedan, southeast Iran. *Annals of Nutrition and*
306 *Metabolism*. 2011;58(1):37-41.
- 307

308
309
310