

Original research Articles

IN-VITRO EVALUATION OF ANTIBACTERIAL AND ANTIFUNGAL EFFICACY OF *BOSWELLIA DALZIELII* STEM BARK EXTRACT

ABSTRACT

The efficacy of *Boswellia dalzielii* (Frankincense) stem bark extract on some bacterial and fungal organisms was evaluated for its in-vitro antimicrobial activities against *Staphylococcus aureus*, *Streptococcus pyogenes*, *Escherichia coli*, *Salmonella gallinarium*, *Aspergillus fumigatus* and *Candida albicans*. The research work was carried out in Biochemistry and Microbiology Laboratories of Federal College of Animal Health and Production Technology, National Veterinary Research Institute, Vom. Well diffusion method was carried out on nutrient agar. MIC, MBC and MFC of the test organisms were carried out on nutrient broth. The phytochemistry revealed the presence of saponin, tannin, flavonoids, cardiac glycosides, steroids, terpenes and phenol in ethanol extracts while resin, alkaloid and glycosides were absent in hot water extracts. Alkaloid was also absent in ethanolic extract. The aqueous extract of the plant exhibited neither antibacterial nor antifungal effects against all test organisms used in the study while the ethanolic extract of the plant showed both antibacterial and antifungal effects on the study organisms. The results of this study also showed that the ethanolic extract of *Boswellia dalzielii* stem bark has activity against all bacteria species used in the study (broad spectrum activity). For gram-negative and positive bacteria, *Salmonella gallinarium* and *Staphylococcus aureus* were the most sensitive while *Escherichia coli* and *Streptococcus pyogenes* were the least respectively. *Candida albicans* was more sensitive than *Aspergillus fumigatus*. It was concluded that the test organisms were susceptible to ethanol extracts of the plant and may be good source of antibiotics.

KEY WORDS: *Boswellia dalzielii*, In vitro, Antibacterial, Antifungal, Extract, Susceptible.

INTRODUCTION

Herbal medicine is the oldest form of medicine known to mankind [1, 2]. It was the mainstay of many early civilization and still the most widely practiced form of medicine in the world today [3]. Many people in developing countries still rely on traditional healing practices and medicinal plants for their daily healthcare needs, in spite of the advancement in modern medicine [4]. Traditional medicine which is widespread throughout the world has been recognized by World Health Organization (WHO) as an essential building block of primary health care. According to reports of World Health Organization, 80% of the world's population relies mainly on traditional therapies which involve the use of plants extracts or their active substance [5]. There is abundant undocumented traditional knowledge of herbal remedies used to treat diseases in most cultures [6]. Different traditional healing practices worldwide are designed for either therapeutic or prophylactic use in human or animal diseases [7, 8]. Several studies carried out in Africa, Asia, Europe, Latin America and North America show that plants are routinely used as remedy for animal diseases [9-14]. Historically, it is documented that humans utilize the same herbal preparations that they use to treat their sick animals [15]. In Nigeria, farmers are known to treat animal diseases with herbs and other traditional medical practices before the advent of orthodox medicine [16]. Traditional medical and veterinary practices remain relevant and vital in many areas in Nigeria due to absence or

43 inadequate provision of modern medical services particularly in rural areas [17]. Ethno-
44 veterinary medical practice is widespread among herdsmen and native livestock producers in
45 northern Nigeria. Traditional remedies in this area include plant extracts from different plant
46 parts [18]. Herdsmen in non-industrialized nations of the world still use medicinal plants for
47 the treatment of livestock diseases, either due to lack of access to trained veterinarians and
48 high cost of orthodox medicines, or the held belief that herbal remedies are more efficacious
49 [19].

50 Plants are also potential sources of modern drugs. A recent survey of United Nations
51 Commission for trade and development (UNCTAD) indicated that about 13% of drugs
52 produced within developed countries are derived from plants [20]. Surprisingly, this large
53 quantity of modern drugs comes from less than 15% of the plants, which have been known to
54 have been investigated pharmacologically [21]. Therefore, since there are so many of these
55 naturally occurring substances in plants, it is obvious that the plant kingdom offers better
56 opportunity of providing useful medicinal compounds.

57 *Boswellia dalzielii* (family Burseraceae), commonly known as frankincense tree, abounds in
58 the Savannah regions of West Africa. The plant has several medicinal uses. The decoction of
59 the stem bark is used to treat rheumatism, septic sores, venereal diseases and gastrointestinal
60 ailments [22, 23]. Phytochemical studies of the plant revealed the absence of alkaloids [24],
61 while saponins, tannins, flaonoids, cardiac glycosides, steroids, and terpenes were found to be
62 present [25, 26]. Oil from the leaves of *Boswellia dalzielii* was found to exhibit significant
63 activity against *Staphilicoccus aureus*, *Bacillus subtilis* and *Candida albicans* [27]. The
64 methanolic and aqueous extracts showed antibacterial and antifungal activities [28, 26].
65 Recent studies of the aqueous extract of the stem bark of *Boswellia dalzielii* showed no
66 antimicrobial activity against all the microbes, used, however, produced some anti-ulcer
67 activity [25]. In another recent study, incensole was found to be part of the chemical
68 composition of the stem-bark of *Boswellia dalzielii* [29].

69 Now-a-days the problem of antimicrobial resistance is growing and the outlook for the use of
70 antimicrobial drugs in the future is still uncertain. In general, bacteria have the genetic ability
71 to transmit and acquire resistance to drugs, which are utilized as therapeutic agents [30].
72 Laboratories of the world have found literally thousands of phytochemicals which have
73 inhibitory effects on all types of microorganisms *in vitro* [31]. Unfortunately, development of
74 effective antimicrobial agents has been accompanied by the emergence of drug-resistant
75 organisms due to the irrational and over-use of antibiotics, failure to complete a course of
76 treatment, genetic versatility of microbes and horizontal transfer of resistant genes among
77 bacterial species. All the mentioned factors diminish the clinical effectiveness of antibiotics
78 [32, 33].

79 In recent time, there has been renewed interest on plants as sources of antimicrobial agents
80 due to their use historically and the fact that a good portion of the world's population,
81 particularly in developing countries, n rely on plants for the treatment of infectious and non-
82 infectious diseases [34]. The aim of the research is to determine the susceptibility of some
83 bacterial and fungal organisms to the ethanolic and aqueous plant extracts and also to
84 determine the minimum antibacterial and antifungal concentrations of the plant extracts.

85

86

87

88

MATERIALS AND METHODS

89

METHODOLOGY

90 The Standard qualitative method as described by Sofowora, [35] was used for phytochemical
91 screening of the plant using ethanol and hot water as the solvents in the biochemistry
92 laboratory. Well diffusion and tube dilution methods were used to determine the antimicrobial
93 properties, minimum inhibitory concentration and minimum bacteriocidal concentration
94 concentrations of the plant extract as described by Cheesbrough, [36] while minimum
95 fungicidal concentrations of the plant extract was determined as described by Picman *et al.*
96 [37].

97

PHYTOCHEMICAL SCREENING

98 The herb *Boswellia dalzielii* was obtained from National Veterinary Research Institute, Vom.
99 This herb was identified as *Boswellia dalzielii* by Mr. Okonkwo, a plant taxonomist attached
100 to the Federal College of Forestry, Jos. The powdered stem bark (100g) was extracted
101 exhaustively with petroleum ether 60-80°C in a Soxhlet apparatus for 24hrs. The marc was air
102 dried and re-extracted with ethanol. The aqueous and ethanolic extracts were separately
103 evaporated under reduced pressure to give solid residues weighing 10.76g and 21.82g,
104 respectively. The residues were then subjected to phytochemical screening using standard tests
105 to show the different types of phytochemical constituents present in the stem [35, 38-40].

106

107

SOURCE OF THE ORGANISMS

108 The organisms were collected from the Central Diagnostic Laboratory of the National
109 Veterinary Research Institute Vom and the work was carried out in college Microbiology
110 Laboratory, Federal College Animal Health and Production Technology Vom, Plateau State.
111 Six strains of Microorganisms were collected in their appropriate culture media. These
112 organisms were *Escherichia coli*, *Salmonella gallinarum*, *Staphylococcus aureus*, *Aspergillus*
113 *fumigatus*, *Candida albicans* and *Streptococcus pyogenes*.

114

115

SENSITIVITY TEST USING WELL DIFFUSION METHOD

116 Molten nutrient agar was prepared and 0.2ml of the organism from the broth culture was
117 inoculated into molten nutrient agar and pour plated and was left on the bench to solidify
118 [41]. Six wells were bored using sterile borer. The extract were dispensed into each well using
119 a sterile micropipette at different concentrations of 500mg/ml, 250mg/ml, 125mg/ml,
120 62.5mg/ml and 31.25mg/ml. Gentamycin and Miconazole were used as positive controls for
121 both bacterial and fungal organisms respectively. The plates were incubated at 37°C for 24
122 hours.

123 **DETERMINATION OF MINIMUM INHIBITORY AND MINIMUM**
 124 **BACTERIOCIDAL CONCENTRATIONS**

125 Tube dilution method was used in varying concentration of the liquid media and the extracts in
 126 test tubes at 500mg/ml, 250mg/ml, 125mg/ml, 62.5mg/ml and 31.25mg/ml were dispensed in
 127 tubes, and 0.2ml of the standardized organism was also dispensed in the same tubes. The tubes
 128 were incubated at 37°C for 24 hours, positive control were also setup. The least concentration
 129 without growth gives the MIC. The MIC is then subcultured into a broth culture tubes that
 130 contain no extracts, the lowest concentration that result in no growth of the subcultured is
 131 noted which indicated MBC, [36].

132 **DETERMINATION OF MINIMUM FUNGICIDAL CONCENTRATION**

133 The hyphal growth inhibition test was used to determine the antifungal activity of the plant
 134 extract against fungal strains as previously described Picman *et al.* [37]. Briefly, dilutions of
 135 the test solutions dissolved in vehicle were added to sterile melted PDA at 45°C to give final
 136 concentrations of 500mg/ml, 250mg/ml, 125mg/ml, 62.5mg/ml and 31.25mg/ml of plants
 137 extracts. The resultant solution was thoroughly mixed and approximately 15 mL was poured
 138 onto the petri plate. Plugs of 1 mm of fungal mycelium cut from the edge of actively growing
 139 colonies were inoculated in the center of the agar plate and then incubated in a humid chamber
 140 at 25°C. Control cultures also received an equivalent amount of vehicle. Three replicates were
 141 used for each concentration. Radial growth was measured when the control colonies almost
 142 reached 1.5 cm.

143

144

145 **RESULTS**



146 **Table 1: The phytochemical components of *B. dalzielii* stem bark**
 147 **extracts.**

s/n	Phytochemicals	Observations	
		Ethanol	Hot water
1	Saponin	++	+
2	Tannins	++	+
3	Resins	+	–
4	Alkaloids	–	–

5	Flavonoids	++	+
6	Glycosides	+	-
7	Cardiac glycosides	++	+
8	Steroids	++	+
9	Terpens	++	+
10	Phenol	+++	+

148

149 Key

150

151 - = absent

152 + = slightly present

153 ++ = moderately present

154 +++= heavily present

155

156

157 **Table 2: Antimicrobial activity of the Hot water extracts**

ISOLATES	Concentration in mg/ml						
	500	250	125	62.5	31.25	-ve	+ve
<i>Aspergillus fumigatus</i>	-	-	-	-	-	-	17 
<i>Candida albicans</i>	-	-	-	-	-	-	18
<i>Staphylococcus aureus</i>	-	-	-	-	-	-	10

<i>Streptococcus pyogenes</i>	-	-	-	-	-	-	15
<i>Escherichia coli</i>	-	-	-	-	-	-	9
<i>Salmonella gallinarum</i>	-	-	-	-	-	-	12

158

159

160

161

162



Table 3: Antimicrobial activity of the ethanol extracts

ISOLATES	Concentration in mg/ml						-ve	+ve
	500	250	125	62.5	31.25			
<i>Aspergillus fumigatus</i>	4	4	3	3	2	-	17	
<i>Candida albicans</i>	5	4	3	3	2	-	18	
<i>Staphylococcus aureus</i>	10	6	4	4	3	-	10	
<i>Streptococcus pyogenes</i>	4	3	2	2	1	-	15	
<i>Escherichia coli</i>	9	7	5	2	2	-	9	
<i>Salmonella gallinarum</i>	7	6	5	3	3	-	12	

163

164

165

166

167

168 **Table 4: MIC, MBC AND MFC OF THE EXTRACTS**

ISOLATES	MIC (mg/ml)	MBC (mg/ml)	MFC (mg/ml)
<i>Aspergillus fumigatus</i>	125	NA	125
<i>Candida albicans</i>	125	NA	–
<i>S. gallinarum</i>	62.5	62.5	
<i>S. aureus</i>	62.5	62.5	
<i>E. coli</i>	125	125	
<i>S. pyogenes</i>	250	250	

169

170

171

172 **DISCUSSION, CONCLUSION AND RECOMMENDATION**173 **DISCUSSION**

174 The phytochemical screening of *Boswellia dalzielii* plants showed that it contains saponin,
 175 tannins, flavonoids, cardiac glycosides, steroids, terpenes and phenol in both ethanol and hot
 176 water extraction while resins and glycosides were present only in ethanolic extract but absent
 177 in aqueous extract. Alkaloids was found to be absent in both hot water and ethanol extraction
 178 (Table 1). This is in accordance with Nwinyi *et al.* [27] and Anago *et al.* [42] who
 179 reported the presence of tannin among the phytochemical properties of the plant
 180 and absence of alkaloid in their aqueous and ethanolic extracts respectively.

181 Hassan *et al.* [40] also reported the presence of tannins, saponins, flavonoids, cardiac
 182 glycosides, steroids and terpenes in methanolic extract of the plant.

183 The aqueous extract of the plant exhibited neither antibacterial nor antifungal effects against
 184 all test organisms used in the study (Table 2). This agreed with the report of Nwinyi *et al.*
 185 [27] and Taiwo *et al.* [43] who stated that aqueous extract of the plant has no
 186 antibacterial effect.

187 The ethanolic extract of the plant showed both antibacterial and
188 antifungal effects on the study organisms (Table 3). This also agreed with
189 Olukemi *et al.* [44], Nwinyi *et al.* [27], Numbo *et al.* [45] who reported that ethanolic extract
190 from *Boswellia dalzielii* have antimicrobial property. According to Campbell [46], the presence
191 of substantial level saponin, phenols and tannins in an extract encourage antimicrobial
192 properties.

193 The results of this study showed that the ethanolic extract of the stem bark of *Boswellia*
194 *dalzielii* has activity against some gram-positive and gram-negative bacteria (broad spectrum
195 of activity) [28]. For gram-negative bacteria (Table 4), *Salmonella gallinarium* was the most
196 sensitive while *Escherichia coli* was the least. For gram-positive (Table 4), *Staphylococcus*
197 *aureus* was the most sensitive while *Streptococcus pyogenes* was the least. *Candida albicans*
198 was more sensitive than *Aspergillus fumigatus* (Table 4). In general, this herb was more active
199 with bacteria than fungi (Tables 4). This is due to the complex nature of fungal cell wall which
200 makes entry of drugs and other chemotherapeutic agents extremely [47]. Nwinyi *et al.* [27]
201 stated that presence tannin is responsible the antibacterial activity of *Boswellia dalzielii*
202 ethanolic extract. According to Olukemi *et al.* [44], *Staphylococcus aureus* is very sensitive to
203 *Boswellia dalzielii* ethanolic extract and also reported that gram-negative bacteria are less
204 susceptible to the extract than gram-positive. The result of the study also correlated with the
205 use of the stem bark of *Boswellia dalzielii* by herbal practitioners in Jos to treat gastroenteritis
206 [27].

207

208 CONCLUSION AND RECOMMENDATIONS

209 The phyto-chemistry screening revealed the presence of saponin, tannin, flavonoids, cardiac
210 glycosides, steroids, terpenes and phenol in ethanol extracts while resin, alkaloid and
211 glycosides were absent in hot water extracts. Alkaloid was also absent in ethanolic extract.
212 The aqueous extract of the plant exhibited neither antibacterial nor antifungal effects against
213 all test organisms used in the study while the ethanolic extract of the plant showed
214 both antibacterial and antifungal effects on the study organisms. The results
215 of this study also showed that the ethanolic extract of the stem bark of *Boswellia dalzielii* has
216 activity against some gram-positive and gram-negative bacteria (broad spectrum of activity).
217 For gram-negative bacteria, *Salmonella gallinarium* was the most sensitive while *Escherichia*
218 *coli* was the least. For gram-positive, *Staphylococcus aureus* was the most sensitive while
219 *Streptococcus pyogenes* was the least. *Candida albicans* was more sensitive than *Aspergillus*
220 *fumigatus*. Root, stem and leaves extracts of *Boswellia dalzielii* were recommended to be tried
221 on other microorganisms to ascertain its efficacy. More so, phytotoxicity of *Boswellia dalzielii*
222 should be carried out to determine the possible toxicity of the pharmaco-active ingredients of
223 the plant.

224

225

226

REFERENCES

- 228 1. Eisenberg D, Davis R, Ettner S. Trends in alternative medicine use in the United States
229 1990-1997; results of a follow up survey. *Journal of American Medical Association*.
230 1998; 280:1569-1575.
- 231 2. Kubmarawa D, Akiniyi JA and Okorie DA. Ethnomedicinal survey of the traditional
232 medicine of Lala people of Nigeria. *International Journal of Medicinal Plant and*
233 *Alternative Medicine*. 2013; 1(3): 39-57.
- 234 3. Srivastava J, Lambert J, Victmeyer N. Medicinal Plants: An expanding role in
235 development. *World Bank Technical Paper*. 1996; No. 320.
- 236 4. Ojewole JAO. Evaluation of the analgesic, anti-inflammatory and antidiabetic
237 properties of *Sclerocarya birrea* (A. Rich.) Hochst. Stem bark aqueous extract in mice
238 and rats. *Phytotherapy Research*. 2004; 18:601-608.
- 239 5. World Health Organization. Summary of WHO guidelines for the assessment of herbal
240 medicines. *herbal gram*. 2010; 28:13-14
- 241 6. Raul P, Pedraza M, Manuela P. Animal health care in India. *Information Centre for*
242 *Low External Input in sustainable Agriculture (ILEIA) Newsletter*. 1990; 8(3):22-23.
- 243 7. McCorkle CM. An introduction to ethnoveterinary research and development. *Journal*
244 *of Ethnobiology*. 1986; 129-140.
- 245 8. Mathias ME. Magic, myth and medicine. *Economic Botany*. 1994; 48(1):3-7.
- 246 9. Ali, Z.A: Folk veterinary medicine in Moradabad District (Uttar Pradesh) India.
247 *Fitoterapia*. 1999; 70:340-347.
- 248 10. Passalacqua NG, De Fine G, Guarrera PM: Contribution to the knowledge of the
249 veterinary science and of the ethnobotany in Calabria region (Southern Italy). *Journal*
250 *of Ethnobiology and Ethnomedicine*. 2006; 2:52.
- 251 11. Yinegar H, Kelbessa E, Bekele T, Lulekal E. Ethnoveterinary medicinal plants in Bale
252 Mountains National Park, Ethiopia. *Journal of Ethnopharmacology*. 2007; 112:55-70.
- 253 12. Lans C, Turner N, Khan T, Brauer G, Boepple W. Ethnoveterinary medicines used for
254 ruminants in British Columbia Canada. *Journal of Ethnobiology and Ethnomedicine*.
255 2007; 3:11.
- 256 13. Alves RRN, Lima HN, Tavares Souto WMS. Plants used in animal healthcare in South
257 and Latin America an overview. In *Ethnoveterinary Botanical Medicine: Herbal*
258 *medicines for animal Health*. 1st edition. Edited by: Katerere RD, Luseba D. CRC
259 *Press New York USA*. 2010; 231-256.
- 260 14. Mathias E, McCorkle CM. Traditional livestock healers. *Rev Sci Tech*. 2004;
261 23(1):277-284.
- 262 15. Offiah NV, Makama S, Elisha IL, Makoshi MS, Gotep JG, Dawurung CJ, Oladipo
263 OO, Lohlum AS, and Shamaki D. Ethnobotanical survey of medicinal plants used in
264 the treatment of animal diarrhoea in Plateau State, Nigeria. *BMC Veterinary Research*.
265 2011; 7:36, 1-9.
- 266 16. Nwude N, Ibrahim MA. Plants used in traditional veterinary medical practice in
267 Nigeria. *Journal of Veterinary Pharmacology and therapeutics*. 1986; 3:261-273.
- 268 17. Kudi AC, Myint SJ: Antiviral activity of some Nigerian medicinal plant extracts.
269 *Journal of Ethnopharmacology*. 1999; 68:289-294.
- 270 18. Alawa JP, Jokthan GE, Akut K. Ethnoveterinary medical practice for ruminants in the
271 subhumid zone of northern Nigeria. *Preventive Veterinary Medicine*. 2002; 54(1):79-
272 90.
- 273 19. Rashid MH, Tanzin R, Ghosh KC, Jahan R, Khatun MA, Rahmatullah M An
274 ethnoveterinary survey of medicinal plants used to treat cattle diseases in Birishiri area,

- 275 Netrakona district, Bangladesh. *Advances in Natural and Applied Sciences*. 2010;
276 4(1):10-13.
- 277 20. UNCTAD/GATT. Markets for selected medicinal plants and their derivatives. *United*
278 *Nations Conference on Trade and Development, Headquarters, Geneva*. 1974; 1-2.
- 279 21. Farnsworth RN, Bingel AS. Problems and prospects of discovering new drugs from
280 higher plants by pharmacological screening. In Sofowora A. (ed)., *Medicinal plants*
281 *and Traditional Medicine in Africa*. 1993; 128-161.
- 282 22. Burkill HM. Useful plants of west Tropical Africa. Second ed., *Royal Botanic Gardens*
283 *Kew*. 1995; 1: 300-301.
- 284 23. Evans W C. Trease and Evans Pharmacognosy, 13th ed. *ECBS/Baillere Tindal, United*
285 *Kingdom*. 1989; 474-475.
- 286 24. Baoua M, Fayn J, and Bassiere J. Preliminary phytochemical testing of some medicinal
287 plants of Niger; *Plant Med. Phytother*. 1976;10: 251-266.
- 288 25. Alemika TOE and Oluwole FS. An investigation of the potentials of *Boswellia dalzielii*
289 and *Commiphora kerstingii* in the treatment of peptic ulcer. *West African Journal of*
290 *Pharmacology and Drug Research*. 1991; 9(10): 91-94.
- 291 26. Adelakun EA, Finbar EAV, Agina SE and Makinde A A. Antimicrobial activity of
292 *Boswellia dalzielii* stem bark; *Fitoterapia*. 1991; 72(7): 822-824.
- 293 27. Nwinyi FC, Binda L, Ajoku GA, Aniagu SO, Enwerem NM, Orisadipe A, Kubmarawa
294 D and Gamaniel KS. Evaluation of the aqueous extract of *Boswellia dalzielii* stem bark
295 for antimicrobial activities and gastrointestinal effects. *African Journal of*
296 *Biotechnology*. 2004; 3(5): 284-288.
- 297 28. Ntiejumokwu S and Alemika TOE. Antimicrobial and phytochemical investigation of
298 the stem bark of *Boswellia dalzielii*. *West African Journal of Pharmacology and Drug*
299 *Research*. 1991; 9(10): 100 - 104.
- 300 29. Alemika TOE, Onawunmi GO and Olugbade TA. Isolation and Characterization of
301 Incensole from *Boswellia dalzielii*; *Journal of Pharmacy and Bioresources*. 2004; 1(1),
302 7 - 11.
- 303 30. Nascimento GGF, Locatelli J, Freitas PC, Silva GL. Antibacterial activity of plant
304 extracts and phytochemicals on antibiotic resistant bacteria. *Brazil Journal of*
305 *Microbiology*. 2000; 31: 247-256.
- 306 31. Cowan MM. Plant products as antimicrobial agents. *Clinical Microbiology Review*.
307 1999; 12:564-582.
- 308 32. Amit R and Shailendra S. Ethnomedicinal approach in biological and chemical
309 investigation of phytochemicals as antimicrobials, (2006). In: Anago, E., Lagnika, L.,
310 Gbenou, J., Loko, F., Moudachirou, M. and Sanni, A. Antibacterial activity and
311 phytochemical study of medicinal plants. *Pakistan journal of Biological sciences*,
312 2011; 14:449-455
- 313 33. Aibinu I, Adenipekun T, Adelowotan T, Ogunsanya T and Dugbemi
314 T. Evaluation of the antimicrobial properties of different parts of
315 *Citrus aurantifolia* (lime fruit) as used locally. *Afr. J. Trad. CAM*, 2007; 4: 185-
316 195.
- 317 34. Ayoola GA, Lawore FM, Adelowotan T, Aibinu IE, Adenipekun E, Coker HAB,
318 Odugbemi TO. Chemical analysis and antimicrobial activity of the essential oil of
319 *Syzygium aromaticum* (clove). *African Journal of Microbiology Research*. 2008; 2:162-
320 166.
- 321 35. Sofowora A. *Medicinal Plants and Traditional Medicine in Africa*, 2nd edition,
322 *Spectrum Book Limited, Ibadan, Nigeria*. 1993; 134-156.

- 323 36. Cheesbrough, M. District laboratory practice in tropical countries part 2. *Antimicrobial*
324 *sensitivity testing. Published by the press syndicate of the University of Cambridge.*
325 *ISBN 0521665469.* 2000; 132-140.
- 326 37. Picman AK, Schneider EF and Gershenson J. Antifungal activities of sunflower
327 terpenoids. *Biochemical Systematics and Ecology.* 1990; 18: 325–328.
- 328 38. Evans WC. Trease and Evans' Pharmacognosy, 14th edition, *W.B Saunders Company*
329 *Ltd., U.K.* 1996; 338-542.
- 330 39. Silva GL, Lee I and Kington A.D. Special problems with the extraction of plants. In:
331 Cannell, R.J.P.(Ed.) *Methods of Biotechnology, Natural Products Isolation, Humana*
332 *Press Inc., Totowa, New Jersey, U.S.A.* 1998; 4: 343-363.
- 333 40. Hassan HS, Musa AM, Usman MA and Abdulaziz M. Preliminary Phytochemical and
334 Antispasmodic Studies of the Stem Bark of *Boswellia dalzielii*. *Nigerian Journal of*
335 *Pharmaceutical Sciences.* 2009; 8(1): 1-6.
- 336 41. Priya K and Ganjewala D. Antibacterial Activities and Phytochemical Analysis of
337 Different Plant Parts of *Nyctanthes arbor-tristis* L. *Research Journal of*
338 *Phytochemistry.* 2007; 1: 61-67.
- 339 42. Anago E, Lagnika L, Gbenou J, Loko F, Moudachirou M and Sanni A. Antibacterial
340 activity and phytochemical study of medicinal plants. *Pakistan journal of Biological*
341 *sciences.* 2011; 14:449-455.
- 342 43. Taiwo IH, Binda L and Kim YC. Antibacterial activity of of the stembark of
343 *Boswellia dalzielii*. *Journal of Ethnopharmacology.* 2006; 95:421-424.
- 344 44. Olukemi MA, Kandakai-Olukemi YT and Mawak JD. Antibacterial activity of the
345 stem bark of *Boswellia dalzielii*. *Journal of Pharmacy and Bioresources.* 2005; 2(2):
346 131-136.
- 347 45. Noumbo T, Serferbe SG, Yaouba A, Fovo JD, Keuete EK. Efficacy of three Local
348 Plant Extracts as Seed Treatment on the Germination, Infection And Vigour Index of
349 two Cotton Seed Varieties from Chad. *International Journal of Applied Biology and*
350 *Pharmaceutical Technology.* 2015; 6(2): 39-45.
- 351 46. Campbell WE. *Antimicrobial Chemotherapy in Medicinal Microbiology*, 6th edition,
352 *Bailiere-London.* 1999; 205-229.
- 353 47. **Bailiere-London.** 
- 354 48. Olukemi MA, Kandakai-Olukemi YT and Bello SS. Antibacterial activity of the stem
355 bark of *Parkia filicoidea*. *J. Pharm. Res. & Dev.* 1997; 5:64-66.