

Efficacy of upsee therapy program with an enriched stimulating active environment on improvement gross, fine motor abilities and weight bearing distribution in spastic diplegic cerebral palsy children.

ABSTRACT

Objectives: This work was carried out to investigate the efficacy of upsee therapy program with an enriched stimulating active environment on improvement gross, fine motor abilities and weight bearing distribution in spastic diplegic cerebral palsy children. **Method:** Thirty children were enrolled in this study and randomly assigned into two groups; group A received (upsee therapy program plus specific physiotherapy training) and group B (specific physiotherapy training program only). Gross motor functional classification system(GMFCS), pegboard test and weight bearing distribution stages evaluation were used to detect and follow gross, fine motor abilities and weight bearing distribution. This measurement was taken before initial treatment and after 12 weeks of treatment. The children parents in both groups A and B were instructed to complete 3 hours of the home routine program. **Results:** Data analysis was available on the 30 spastic diplegic cerebral palsied children participated in the study. The difference between pre and post-treatment results was more significant in (GMFCS) study group than the control group. Weight-bearing distribution stages evaluation and pegboard test demonstrate representative improvement in the study groups ($p=0.0001$) while insignificant in the control groups. **Conclusion:** The addition of upsee therapy program to specific physiotherapy training is recommended in improving gross, fine motor abilities and weight bearing. So this suggested approach may be used as a selective choice for improving walking, hand functions and weight bearing abilities in spastic diplegic cerebral palsy (CP) children.

Keywords: upsee therapy, upright posture, cerebral palsy, walking performance

Introduction

Cerebral palsy is considered the greatest cause of physical dysfunction in childhood¹. The child becomes immobilized and non-weight bearing which lead to impaired bone mineral density plus reduced muscle mass which lead to improper muscle pumping and decreasing of bone blood supply leading to decrease bony mass^{2,3}. There is a relationship between skeletal morphology and muscle function so to improve gross motor skills we should concentrate on activation of musculoskeletal system⁴ so upsee therapy training could be used in the modulation of musculoskeletal system⁵.

Non weight bearing state of cerebral palsy children lead to ~~coxa-valga~~ **valga thigh** (neck-shaft angle of the femur exceed than normal) with great spasticity of hip adductor and iliopsoas which direct femoral head against lateral rim of the acetabulum lead to delaying of hip joint development which leads to subluxation and dislocation of the hip-lead to pain, scoliosis and impaired upright posture^{6,7}

Upsee therapy supplies the impaired mobility **cerebral paisy** (CP) children with an enriched stimulating active environment that form the foundations and the keys for acquisitions of new skills and reacquisition of lost skills⁸. Cerebral palsy is associated with impairment of perception-cognition complex which is the base of gross and fine motor skills acquisition⁹. Upright Posture control by upsee therapy keep the head, neck, trunk in a functional position to be able to connect with the environment leading to facilitate hand functions, ADL and walking abilities¹⁰

Changing of the child passive home environment to active one place the hand ready to perform all hand functions from upright posture leading to improvement of balance control, hand functions skills, walking abilities and provide CP children with the new sensory experience of movement and participation¹¹. The dynamic compression forces of the upsee system during walking increase hip stability via increasing of acetabulum depth^{12,13} through functional weight bearing which has more impaction on joint stability while the worst instability occurred by spasticity¹⁴

Upsee therapy can provide active standing which improves the bone mineral contents leading to increased bone density as increase femoral neck mineral density¹⁵ and static standing increase lumbar bone mineral density¹⁶ in addition the cognitive skills development need interplay between higher centers, posture alignment and right enriched environment which can be provided by upsee therapy program¹⁷

Material and method

Subjects

Thirty children from both sexes with spastic diplegic cerebral palsy children were enrolled for this study, aged 5 to 10 years at the time of enrollment due to the children in this age are able to participate in (GMFCS) levels. Children are not able to walk without assistance, Children who alternatively met the inclusion standard were ruled out if they had severed tight muscles, epileptic fits, osteoporosis or brittle bone, Current fracture unless medical clearance has been given, post-operative weight-bearing restrictions, pain due to muscle strain or joint subluxation occurring during standing, previous BoNT-A injections in the L.L. in the past year or previous lower limb surgery.

Children randomized to the experimental group (A) received upsee therapy program plus specific physiotherapy training. Children randomized to the control group (B) received specific physiotherapy training program only. The individual-based upsee therapy program treatment sessions of 60 minutes were conducted five times weekly for 12 weeks in a physiotherapy treatment room after the specific physiotherapy training for one hour. In addition, children in the two groups were exposed to home routine program 3 hours daily for the 12 week treatment period.

Outcome measurements

1-Gross motor functional classification system(GMFCS)

It includes 5 levels for evaluation and follows up a gross motor abilities

I and II Walk without support

III Walk with a mobility device

IV and V cannot sit or walk without support

It has proven to give an accurate description method for evaluation of the gross motor abilities of children with CP GMFCS reflect better indication toward spastic diplegic CP children more than other forms.

GMFCS levels:

Level I the children who walk without restriction

Level II the children have restriction to walk elongated lengths and at equilibrium

Level III the children walk using a hand-held mobility tool

Level IV is carried by manual or mechanical wheelchair.

Level V there is loss of head and trunk control, requiring wide use of mechanical and physical aids¹⁸

2-Weight-bearing distribution stages evaluation

By using the weight scale we could locate the time of pressure that the child could perform on the scale to detect the level of weight bearing distribution

Stage 1: No weight bearing

Stage 2: Flickers of weight bearing

Stage 3: Weight-bearing extended for 5-30 seconds

Stage 4: Weight-bearing extended for more than 30 seconds however less than 2 minutes

Stage 5: Maintains symmetrical weight bearing through legs¹⁹

3- Evaluation of hand functions:

Via 9 whole pegboard test by determining the level of hand function grading by locating the time needed to fit all materials on its position.

Intervention

The main goal of the upsee rehabilitation program is to gain utmost functional skills, upright postural control, and hand function skills with minimal external assistance²⁰

Both groups (A and B) received a specific physiotherapy program, like the following:

Facilitation of milestones, inhibition of released primitive reflexes, Balance training program, inhibition of spasticity, facilitation of hand function, correct deformity, gait training, proprioceptive training and using of orthoses.

The experimental group (group A) received upsee therapy program adding to specific physiotherapy program as following:

Upsee therapy harness is an orthotic device. it was worn by the therapist, parents and child which is used for facilitation of standing, walking and hand functions skills²¹ it is indicated in children with severely impaired trunk and postural control²² which is developed from **cranium**-caudal direction starting by head control then upper trunk control finally lower trunk control and pelvis control^{23,24}

The upsee system consists of a waistcoat with the pelvic belt, groin support with adjustable straps to link the child from his shoulder to adult pelvic belt and sandals shared from the child and therapist. The child put on his ankle-foot orthoses before wearing upsee system⁸. The degree of weight bearing can be controlled by the therapist via changing of the supporting straps via taking the weight through therapist legs²⁵

The specific physiotherapy therapy session with upsee therapy system last for 2 hours 7 days/weeks for 3 months. Starting of weight bearing 15-20 minutes in standing and walking, increased by 5 minutes every few days until they reach 60 minutes of standing and walking²⁶ Home routine program could be performed 3h/day.

Upsee therapy program:

1- Upsee therapy from standing:

- At first, the **therapist gives** some support to the child's shoulder from front to encourage the hand function activities
- Start standing in front of a mirror then sway body to both sides
- Allow the child to perform reaching in all directions
- Shifting the weight slowly to less weight bearing side and encourage the reaching training on that side
- Step forward by the more weight bearing leg to initiate the walking
- Shifting weight from side to side
- A pegboard table placed at the level of the child's elbow or slightly above containing different types of toys, puzzles, sands, water, paint, and different occupational material
- Catching and throwing a ball, kick a balloon and ball
- When the child weight bearing improved loosen the shoulder straps slightly to increase the trunk control
- Put different magnet shapes on the door, draw, paint, and copy of a design

2-Upsee therapy from walking

- Take slowly one step forward and backward then sideways and backward
- locate one foot along with the other and by the therapist, leg induce rocking the child anterior and posterior
- Start walking with short steps and a large base of support

- Walk through obstacles, walking on different surfaces, walking downstairs then upstairs
- Walking in different directions looking for toys
- Encourage the child to locate the direction and initiate the step
- Kicking ball and balloons during walking
- Walking in the sand, garden of low and high surface
- Ask the child to point out to determine the direction by his head, hand then leg¹⁹

Result

Patients' characteristics

Table 1 display the demographic and analytic traits of all patients. There were 11 boys (36.67%) and 19 girls (63.33%) and in term of right-hand dominance reported in 22 patients (73.33%), and also 8 patients (26.67%) were left-hand dominance. There was no representative change within both groups in relation to age ($p=0.8038$), toward sex ($p=0.2712$) and in term of hand dominance ($p=0.1054$).

Table 1) patients characteristics

Variables	Study group N=15	Control group N=15	P-value
Age	7.53±1.64	7.40±1.24	0.8038
Sex N%			
Boys	4 (26.67%)	(46.67%) 7	0.2712
Girls	11 (73.33%)	8 (53.33%)	
Hand dominance N%			
Right	9(60%)	13(86.67%)	0.1054
Left	6(40%)	2(13.33%)	

Changes in gross motor abilities

Mean test scores and SD for both groups are demonstrated in table 2. The mean record of gross motor abilities level in the two groups at (pre treatment level) was insignificant ($p>0.05$) while the two groups had an expressive enhancement in gross motor abilities at post-treatment level ($p<.05$). The average improvement of gross motor abilities level had a tendency to be highly representatives improvement in the experimental group (2.33 ± 0.49 versus 3.20 ± 0.41 ., $p=0.0001$) than in the control group (3.20 ± 0.77 versus 3.47 ± 0.52 , $p=0.0406$). The percentage of improvement of gross motor abilities level was (27.19%) in the study group compared to the (7.78%) in the control group.

Table2: The average test of gross motor abilities level in both groups.

gross motor abilities levels	Study group Mean \pm SD	Control group Mean \pm SD	P-value (within groups)
Pre-treatment	3.20 ± 0.41 .	3.47 ± 0.52	0.1299
Post-treatment	2.33 ± 0.49	3.20 ± 0.77	0.0010
Improvement%	27.19%	7.78%	0.0005
P-value (within groups)	0.0001	0.0406	

Changes in weight-bearing distribution:

Mean test scores and SD for the two groups are demonstrated in table 3. The mean record of weight bearing distribution level in the two groups at (pre-treatment) was insignificant ($p>0.05$) while the two groups had an expressive enhancement in weight bearing distribution at post-treatment level ($p<.05$). The average improvement of weight bearing distribution level had a tendency to be highly representatives improvement in the study group (2.47 ± 0.52 versus 1.53 ± 0.52 , $p=0.0001$) while insignificant representatives in the control group (1.73 ± 0.70 versus 1.53 ± 0.52 , $p=0.0824$). The percentage of improvement of weight bearing distribution level was (61.44%) in the study group compared to the (13.07%) in the control group.

Table 3: The average test of weight bearing distribution level in both group

Weight-bearing distribution level	Study group Mean \pm SD	Control group Mean \pm SD	P-value (within groups)
Pre-treatment	1.53 \pm 0.52	1.53 \pm 0.52	1.0000
Post-treatment	2.47 \pm 0.52	1.73 \pm 0.70	0.0030
improvement%	61.44%	13.07%	0.0052
P-value (within groups)	0.0001	0.0824	

Changes in pegboard test

Mean test scores and SD for both groups are displayed in table 4. The mean record of pegboard score in both groups at pre-treatment was insignificant ($p > 0.05$) while the two groups had a representatives enhancement in pegboard score post-treatment ($p < .05$). The average improvement of pegboard score had a tendency to be highly representatives improvement in the study group (44.67 ± 4.42 versus 49.00 ± 5.41 $p = 0.0001$) while insignificant representatives in the control group (48.67 ± 5.81 versus

48.00 ± 5.61 , $p = 0.1643$). The percentage of improvement of pegboard score was (8.8%) in the study group compared to the (1.4%) in the control group.

Table 4: The average test of pegboard score in both groups:

pegboard score level	Study group Mean \pm SD	Control group Mean \pm SD	P-value (within groups)
Pre-treatment	49.00 \pm 5.41	48.00 \pm 5.61	0.6230
Post-treatment	44.67 \pm 4.42	48.67 \pm 5.81	0.0429
Improvement%	8.8%	1.4%	0.0001
P-value (within groups)	0.0001	0.1643	

Discussion

The spastic diplegic C.P had more liability for walking mobility because the upper limbs were less affected than L.L so they can participate better in activity²⁷. The most

important improvement time in main motor developmental stages and muscle mass in cerebral palsy children is the first 10 years because in this period the body structure, function, participation, and mobility could be influenced better and the improvement in progress. Beyond this age, the secondary motor development either improved or be stable²⁸.

The upsee therapy supply the motor impairment children with an enriched stimulating environment (by putting the children in variant circumstances and difficult situations with massed physical practice) lead to enter the child in active participation which is the key of active motor learning and gaining of motor skills²⁹.




Fig.1) Underlying mechanisms of upsee therapy³⁰.

The functional weight bearing with upsee system stimulate the reciprocal inhibition between spastic and anti-spastic muscles for improving selective motor control leading to a decrease of the degree of spasticity. A therapist could provide the lowest degree of support to the child allowing the child to increase his weight on his legs leading to sufficient inhibition of abnormal co-contraction leading to increasing of the motor control of L.L that improve walking abilities³¹.

Unilateral weight bearing and weight shift training is a vital developmental skill required for lower limb motor control not only the start of upright mobility but also for balance control abilities³². Upper trunk with upper limbs should have a degree of freedom to be coordinated during reaching and ADL³³. Posture stability at pelvis and trunk is considered the biomechanical key for enhanced gross and fine motor abilities as upright postural control, eye-hand coordination, and hand functions skills³⁴.

The requirements of posture control include the presence of postural reactions(righting-equilibrium and protective reactions), controlling on primitive and tonic reflexes, reciprocal inhibition mechanism, motor development and normal intention movement and normal muscle tone plus the normal posture curves after labor thoracic and sacral curves and with four-foot kneeling cervical curve was formed and with sitting and standing lumbar curve was formed³⁵




Fig. 2) Upsee therapy effects and preventions in cerebral palsy

Conclusion:

The addition of upsee therapy program to specific physiotherapy training is recommended in improving gross, fine motor abilities and weight bearing. So this suggested approach may be used as a selective choice for improving walking, hand functions and weight bearing abilities in spastic diplegic C.P children.

References:

- 1- Mann K. I, Cans C(2009): Cerebral palsy update. *Brain and Development*. [cited 28 July 2009];31(7):537-44.
- 2-Schonau E, Werhahn E, Schiedermaier U, Moskow E, Schiessl H, Scheidhauer K, et al. Influence of muscle strength on bone strength during childhood and Adolescence. *Horm Res* 1996;45(1):63-6.
- 3-. Rauch F, Schoenau E. Changes in bone density during childhood and adolescence: An approach based on bone's biological organization. *J Bone Miner Res* 2001; 16(4):597-604.
- 4-Frost HM. The mechanostat: A proposed pathogenic mechanism of osteoporosis and the bone mass effects of mechanical and nonmechanical agents. *Bone Miner* 1987;2(2):73-85.
- 5- Stark C, P. Smyrni, A. Stabrey O. Semler E. Schoenau (2010); Effect of a new physiotherapy concept on bone mineral density, muscle force and gross motor function in children with bilateral cerebral palsy *Musculoskelet Neuronal Interact* 10(2):151-158
- 6- Scrutton D, Baird G, Smeeton N. Hip dysplasia in bilateral cerebral palsy: incidence and natural history in children aged 18 months to 5 years. *Dev Med Child Neurol*. 2001;43(9):586-600.

7-Flynn J, Miller F. Management of hip disorders in patients with cerebral palsy. *J Am Acad Orthop Surg.* 2002;10:198-209.

8- Neill SM (2014): The benefits of upright movement for children with motor impairment: a literature review *Clinical Research Manager, Firefly*

9- Patni, k . (2018): Ayurvedic management of dyskinetic cerebral palsy. A case report, India *Int. J. Res. Ayurveda Pharm.* 9 (1), 2018

10- Bartlett DJ and Palisano RJ. (2002): perceptions of factors influencing the acquisition of motor abilities of children with cerebral palsy: implications for clinical reasoning. *Phys Ther*; 82. p. 237-248.

11. Colver A. What are we trying to do for disabled children? *Current Paediatrics.*2006; 16: 501-05

12. Abel MF, Wenger DR et al. Quantitative analysis of hip dysplasia in cerebral palsy: A study of radiographs and 3-D reformatted images. *Journal of Pediatric Orthopedics*, 14:283-289, 1994

13. Vidal J et al. The anatomy of the dysplastic hip in cerebral palsy related to prognosis and treatment. *International Orthopaedics*, 9:105-110, 1985

14. Soo B, Howard JJ, Boyd RN et al Hip displacement in cerebral palsy. *Journal of Bone Joint Surgery. Am.* 2006; 88: 121-9

15-Chad KE, Bailey DA et al The effect of a weight-bearing physical activity program on bone mineral content and estimated volumetric density in children with spastic cerebral palsy. *The Journal of Pediatrics* (1999)Vol 135 Number 1 pp115-117.

16. Carlton JM, Ward KA et al. A randomized controlled trial of standing programme on bone mineral density in non-ambulant children with CP. *Archives of Dis Child* 2004; 89(2) 131-135

17. Port RF, van Gelder T. *Mind as Motion: Exploration in the Dynamics of Cognition*. Cambridge, MA: MIT Press. (1995) Material and method

18-Palisano R, Rosenbaum P, Walter S, Russell D, Wood E and Galuppi B. Development and reliability of a system to classify gross motor function in children with cerebral palsy. *Dev Med Child Neurol*. 1997;39(4):214-23.

19-Leckey 2015 upsee therapy program: a parents guide to choosing activities for improving mobility, ability, and participation. firefly northern Ireland

20-Butler P. A preliminary report on the effectiveness of trunk targeting in achieving independent sitting balance in children with cerebral palsy. *Clin Rehabil*. 1998; 12:281–293. [PubMed: 9744664]

21- Leckey- Upsee from Firefly. <http://www.fireflyfriends.com/upsee>. Accessed May 4, 2016.

22-Flores, Megan; Manella, Kathleen; and Ardolino, Elizabeth, "Upsee Daisy! Gross Motor Outcomes after Dynamic Weight Bearing in Two Children with Truncal Hypotonia: A Case Series" (2017). *Physical Therapy*. 9.

23-Rachwani J, Santamaria V, Saavedra SL, Woollacott MH. The development of trunk control and its relation to reaching in infancy: a longitudinal study. *Front Hum Neurosci*. 2015; 9:94.doi: 10.3389/fnhum.2015.00094 [PubMed: 25759646]

- 24-Saavedra SL, van Donkelaar P, Woollacott MH. Learning about gravity: segmental assessment of upright control as infants develop independent sitting. *J Neurophysiol.* 2012; 108:2215–2229. [PubMed: 22832568]
25. Gajdosik CG & Gajdosik RL Chapter 6 Musculoskeletal Development and Adaptation in: Campbell S, Vander Linden DW & Palisano RJ (Eds) ‘Physical Therapy for Children’ 3rd Edition (2006) Pub Saunders Elsevier. ISBN 13:978-0-7216-0378-0 2006
- 26- Dalziel B, Hesketh k, Kim C, Holland Bloorview Kids Rehabilitation Hospital A teaching hospital fully affiliated with the University of Toronto www.hollandbloorview.ca
- 27- Kulak W, Sobaniec W, Smigielska-Kuzia J, Kubas B, Walecki J. A comparison of spastic diplegic and tetraplegic cerebral palsy. *Pediatr Neurol* 2005;32(5):311-7.
- 28-. Rosenbaum PL, Walter SD, Hanna SE, Palisano RJ, Russell DJ, Raina P, et al. Prognosis for gross motor function in cerebral palsy: Creation of motor development curves. *JAMA* 2002;288(11):1357-63
29. Chiarello LA & Kolobe THA Campbell S, Vander Linden DW & Palisano RJ Chapter 31 Early Intervention Services in (Eds) ‘Physical Therapy for Children’ 3rd Edition (2006) Pub Saunders Elsevier. ISBN 13:978-0-7216-0378-0 2006
- 30- Argandoña, Metal, International Classification of Functioning, Disability, and Health - Children and Youth Version. ICF-CY (World Health Organisation 2007) ISBN-13 9789241547321
31. Valvano J. Activity-focused motor interventions for children with neurological conditions. *Phys Occup Ther Pediatr.* 2004; 24:79-107
32. Adolph, K. Advances in research on infant motor development. Paper presented at APTA Combined sections meeting Tampa FL. <http://apta.org> 2003

- 33-Schneiberg S, Sveistrup H, McFadyen B, McKinley P, Levin MF. The development of coordination for reach-to-grasp movements in children. *Exp Brain Res*. 2002; 146:142–154. [PubMed: 12195516]
- 34-Cheng HYK, Lien YJ, Yu YC, et al. The effect of lower body stabilization and different writing tools on writing biomechanics in children with cerebral palsy. *Res Dev Disabil*. 2013; 34(4):1152–1159. [PubMed: 23376050]
- 35-Wandel JA (2000) Positioning and handling. In JW Solomon (Ed) *Pediatric Skills for Occupational Therapy Assistants*. London: Mosby
- 36-Pope PM (2007) *Severe and complex neurological disability: management of the physical condition*. London: Elsevier
- 37-Gibson SK, Sprod JA, Maher CA (2009) The use of standing frames for contracture management for nonmobile children with cerebral palsy. *International Journal of Rehabilitation Research*, 32(4) p316-23
- 38-Krueger, LJ & Coleman JMS (2010) Let's stand together. *Advance for Physical Therapy & Rehab Medicine* [serial online] 21(8),p28. Available: <http://physical-therapy.advanceweb.com/Archives/Article-Archives/Lets-Stand-Together.aspx> via the internet. Accessed 5 April 2011.
- 39-Paleg G (2008) Synthesised literature review on Supported Standing. Doctoral Thesis. Shortened version accessed from Easy stand website:<http://www.easystand.com/health-benefits/research-categories.cfm>. Accessed 5 April 2011.
- 40-Gericke T (2006) Postural management for children with cerebral palsy: a consensus statement. *Developmental Medicine & Child Neurology*, 48, 244.
- 41-Labandz, S (2011) Using standers to position children for success. *ADVANCE for Physical Therapy & Rehab Medicine* [serial online] 18(20),p29. Available: <http://physical-therapy.advanceweb.com/Archives/Article-Archives/Heightened-Awareness-2.aspx> via the INTERNET. Accessed 22 March 2011.
- 42-Krueger, LJ & Coleman JMS (2010) Let's stand together. *ADVANCE for Physical Therapy & Rehab Medicine* [serial online] 21(8),p28. Available: <http://physical->

therapy.advancweb.com/Archives/Article-Archives/Lets-Stand-Together.aspx via the INTERNET. Accessed 5 April 2011.

43-Smith, Y (2010) Stand for success. ADVANCE for Physical Therapy and Rehab Medicine [serial online] 21(20) p40. Available: <http://physical-therapy.advancweb.com/Archives/Article-Archives/Stand-for-Success.aspx> via the INTERNET. Accessed 5 April 2011

UNDER PEER REVIEW