

1 **Original Research Article**

2 **EFFECT OF BIOFERTILIZERS AND BIOCONTROL AGENTS IN ENHANCING GROWTH AND YIELD**  
3 **OF BRINJAL UNDER LOW COST NATURALLY VENTILATED POLYHOUSE DURING OFF SEASON**

4 **ABSTRACT**

5 An experiment was conducted to study the combined effect of biofertilizers and biocontrol agents on  
6 growth and yield of brinjal under naturally ventilated polyhouse during off season in the experimental  
7 polyhouse of the College of Agricultural Engineering and Post Harvest Technology (CAEPHT), Central  
8 Agricultural University (CAU), Ranipool, Sikkim. The experiment was planned with 3 treatments and six  
9 replications viz. T<sub>1</sub>- FYM 5 kg/m<sup>2</sup>, T<sub>2</sub>- FYM 5 kg/m<sup>2</sup> + biofertilizer (a mixture of *Azotobacter* + PSB @ 10  
10 g/kg FYM each), T<sub>3</sub>- FYM 5 kg/m<sup>2</sup> + biofertilizer + biocontrol agent (a mixture of *Pseudomonas*  
11 *fluorescens* + *Trichoderma* @ 5 g/kg FYM each). There was significant variation in vegetative growth and  
12 yield among all the treatments. The maximum plant height (45.62 cm), the number of branches/ plant  
13 (11.17) and the number of leaves/ plant (50.05), the number of fruits/ plant (38.9) and fruit yield/ plant  
14 (810 g) were observed with treatment T<sub>3</sub> which was at par with the treatment T<sub>2</sub> and were significantly  
15 higher than the treatment T<sub>1</sub> receiving FYM singly. Organic manure (FYM) inoculated with biofertilizers  
16 may, therefore, be recommended for organic brinjal cultivation-production for cultivation under naturally  
17 ventilated polyhouses in Sikkim (India) and application of biocontrol agents may be limited to areas  
18 having some history of occurrence of diseases.

19 Key words: *Brinjal*, *biofertilizers*, *biocontrol agents* and *naturally ventilated polyhouse*.

20  
21 **1.INTRODUCTION**

22 Brinjal or eggplant (*Solanum melongena* L.) is an important solanaceous vegetable crop widely  
23 grown in the subtropical and tropical regions of the world. It is of much importance as a warm weather  
24 vegetable crop of Far East being grown extensively in India, Bangladesh, Pakistan, China and the  
25 Philippines. In India, it is one of the most common, popular and principal vegetable crops grown  
26 throughout the country. Brinjal occupies 669 Thousand Ha of total area and produces 12400 Thousand  
27 MT) [1]. In the southern states with mild climatic conditions, its bearing-harvest?? period is prolonged

Comment [P1]: Add latin name of brinjal

28 whereas in the northern parts it is shortened. It is a versatile crop adapted to different agro-climatic  
29 regions and can be grown throughout the year whereas in the hilly regions, it is cultivated only in the  
30 summer season.

Comment [P2]: Where??

31 Biofertilizer is a substance which contains living microorganisms which, when applied to seed, plant  
32 root, or soil, colonizes the rhizosphere of the plant and promotes the growth by providing essential  
33 nutrients or make available primary nutrients to the host plant [2]. The use of biofertilizers is beneficial in  
34 regenerating the soil health by enriching fertility and fulfilling plant nutrient requirements by supplying the  
35 organic nutrients through microorganism and their byproducts [3]. Microorganism in biofertilizer provides  
36 three primary nutrients N, P and K through atmospheric nitrogen fixation, phosphorous solubilization, and  
37 potash mobilization which have potential to reduce the use of chemical fertilizers to the tune of 50% and  
38 increase the productivity up to 20% [4].

Comment [P3]: Add some more previous studies about effects of biofertilizers and biocontrol on the plant growth

39 The major constraint in the production of brinjal is the bacterial wilt disease caused by *Ralstonia*  
40 *solanacearum* which constitutes a serious obstacle to the cultivation of the economically important  
41 brinjal among other crops, causing total damage of plantations before as well as after bearing fruits [5].  
42 Biological control could have an important role in the management of bacterial wilt [6]. Effective  
43 management of bacterial wilt of brinjal by *Pseudomonas fluorescens* in field experiment signifies its  
44 potentiality and scope as a plant growth promoting rhizobacteria (PGPR) when formulated using effective  
45 substrate carrier and adhesive [5]. But reports on the use of a combination of biocontrol agents and  
46 biofertilizers in the quality and quantity production of brinjal are very scanty. *Trichoderma* and  
47 *Pseudomonas fluorescens* are effective against damping off, collar rot and seedling blight diseases of  
48 vegetables [7][8][9][10].

49 Sikkim being an organic state, the demand for organic vegetables is very high. Therefore, there is  
50 a need to produce qualitative and quantitative vegetables with high quality and yield through an organic  
51 mode of farming. Organic farming through the use of a combination of biofertilizers and biocontrol agents  
52 along with locally available farm manures (FYM, vermicompost, etc.), not only gives the quality organic  
53 produce but, also sustains the soil health and environment friendly practices for brinjal cultivation in the  
54 terrace farm lands of Sikkim. Keeping above points in view, present investigation has been undertaken to

55 investigate the effect of biofertilizers and biocontrol agents in enhancing growth and yield of brinjal under  
56 low cost naturally ventilated polyhouse (NVP) during the winter season.

## 57 2. METHODS AND MATERIALS

58 An experiment was conducted during October, 2012 to March, 2013 at the All India Coordinated  
59 Research Project on Plastics Engineering and Technologies (AICRP on PET) experimental field of  
60 College of Agricultural Engineering and Post-Harvest Technology, CAU, Ranipool, Sikkim to evaluate the  
61 effect of biofertilizers and biocontrol agents in enhancing growth and yield of brinjal as an offseason crop  
62 under low cost naturally ventilated polyhouse. Brinjal being a cross-pollinated crop, bee-hive with bee  
63 colony was installed in the polyhouse to enhance pollination. The soil of the experimental site was sandy  
64 loam (sand: 62%, silt: 23%, clay: 15%) with pH of 6.2.

65 Organic equivalent dose of recommended NPK (125:100:50 kg/ha) for brinjal as suggested by [11]  
66 was considered and manuring doses were calculated based on recommended doses of nitrogen (125  
67 kg/ha) for FYM. The recommended NPK dosage was found to be equivalent to 5 kg FYM per m<sup>2</sup>. The  
68 experiment was laid out in randomized block design (RBD) with 3 treatments and six replications viz. T<sub>1</sub>:  
69 FYM 5kg/m<sup>2</sup>, T<sub>2</sub>:FYM 5kg/m<sup>2</sup> + biofertilizer (a mixture of *Azotobacter* + PSB @ 10g/kg FYM each),  
70 T<sub>3</sub>:FYM 5kg/m<sup>2</sup> + biofertilizer (a mixture of *Azotobacter* + PSB @ 10g/kg FYM each) + biocontrol agent (a  
71 mixture of *Pseudomonas fluorescens* + *Trichoderma* @ 5g/kg FYM each).

72 The biological resources [*Trichoderma* (Strain UBT-18), *Pseudomonas fluorescens* (Strain VPF-1),  
73 *Azotobacter* (Strain UBAZ-1) and Phosphate solubilizing bacteria (Strain UBPS-9)] used in the  
74 experiment were collected-provided from Department of Plant Pathology, Faculty of Agriculture, UBKV.

75 The seedlings of brinjal were transplanted on raised beds of 15 cm height with row spacing of 50  
76 cm and seedling spacing of 45 cm in the low-cost NVP on October 10, 2012. The data were recorded on  
77 various growths and yield parameters viz. plant height, number of branches, number of leaves, number of  
78 fruits/plant and fruit yield/plant. The data collected for various parameters were subjected to statistical  
79 analysis using RBD One Factor SPSS-16 software.

## 80 3. Result and Discussion

### 81 3.1. Effect of Biofertilizers and Bio-control Agents on Vegetative Growth of Brinjal

82 | At the early stages of crop plant growth, the variation in vegetative growth among the treatments  
 83 | was insignificant. During the later stages (60 and 90 DAT), the treatments inoculated with biofertilizers  
 84 | alone (T<sub>2</sub>) and combination of biofertilizers + bio-control agents (T<sub>3</sub>) were observed to be varying  
 85 | significantly on vegetative growth of brinjal than the treatment (T<sub>1</sub>) receiving only FYM equivalent dose of  
 86 | recommended NPK.

87 | **Table 1: Effect of biofertilizers and biocontrol agents on the growth of brinjal**

Treatment	60 DAT*			90 DAT*			No. of fruits / plant	Fruit yield/ plant (g)
	Plant height (cm)	No. of branches/plant	No. of leaves/plant	Plant height (cm)	No. of branches/plant	No. of leaves/plant		
T <sub>1</sub>	15.96	6.00	10.26	32.76	8.74	28.94	30.80	709.20
T <sub>2</sub>	22.23	7.10	17.45	39.87	10.26	37.56	34.00	796.70
T <sub>3</sub>	25.92	7.73	18.67	45.62	11.17	50.05	38.90	810.00
LSD at 5%	3.74	NS	3.24	5.92	1.10	7.74	5.33	79.95

88 | \*DAT: Days after transplanting

89 | The maximum plant height (25.92 cm) and the number of leaves (18.67) were recorded with-in  
 90 | treatment T<sub>3</sub> at 60 DAT which shows performance at par with the treatment T<sub>2</sub> and were significantly  
 91 | higher than the treatment T<sub>1</sub> receiving FYM alone. At 90 DAT, the maximum plant height (45.62 cm), the  
 92 | number of branches (11.17) and number of leaves (50.05) was observed with-in treatment T<sub>3</sub> which  
 93 | showed performance at par with the treatment T<sub>2</sub> and were significantly higher than the treatment T<sub>1</sub>  
 94 | receiving FYM alone. Biofertilizers and bio-control agents were found to be effective in increasing  
 95 | vegetative growth parameters for organic brinjal. Higher vegetative growth in plants treated with  
 96 | biofertilizers and biocontrol agents may be attributed to improvement in plant mineral concentration

Formatted: Highlight

97 | through better nitrogen fixation caused by biofertilizer application [12], increase in phosphorus uptake by  
98 | plant caused by phosphate solubilising bacteria [13] and disease protection as well as plant growth-  
99 | promoting rhizobacteria [effects](#) caused by biocontrol agents [14]. Increase in plant height, [the](#) number of  
100 | branches/ plant and number of leaves/ plant due to [the](#) application of biofertilizers have also been  
101 | reported by [15] in tomato, [16] in gherkin. The application of biocontrol agents may have protected the  
102 | plant from disease incidence by colonizing the rhizosphere of the plant preceding to [the](#) occurrence of  
103 | any harmful disease causing pathogens as beneficial plant growth-promoting rhizobacteria and so  
104 | enhanced the growth (plant height, number of branches and number of leaves) [17]. Similar findings were  
105 | also reported by [18] and [11] for brinjal.

### 106 | **3.2. Effect of Biofertilizers and Bio-control Agents on Yield of Brinjal**

107 | The maximum number of fruits/ plant (38.90) and fruit yield/ plant (810 g) was recorded [with-in](#) the  
108 | treatment T<sub>3</sub> which showed performance at par with the treatment T<sub>2</sub> and were significantly higher than  
109 | the treatment T<sub>1</sub> receiving FYM alone. Azotobacter may have enhanced the available nitrogen in the soil  
110 | [14] and the inoculation of phosphate [solublizing](#)[solubilizing](#) microorganisms may have increased plant N  
111 | and P uptake [19], which led to [increasing](#) in yield of brinjal. Increase in [the](#) number of fruits/ plant and  
112 | fruit yield/ plant due to the application of biofertilizers have also been reported by [14] in tomato, [20] in  
113 | [safflower](#), [15] in gherkin and [21] in brinjal. Application of biocontrol agents increases the number of  
114 | fruits/ plant and fruit yield/ plant probably due to its major role as antagonistic endophytic bacteria as well  
115 | as plant growth-promoting rhizobacteria. Similar findings were also reported by [17] [11] for brinjal.

116 |

### 117 | **4. CONCLUSION**

118 | The findings [in this study](#) revealed that plant growth and yield of brinjal (local var.) cultivated within the  
119 | [low-cost](#) NVP in the [mid-hill](#) region of Sikkim have been affected significantly by combined inoculation of  
120 | biofertilizers (*Azotobacter* + PSB) and bio-control agents (*Pseudomonas fluorescens* + *Trichoderma*).  
121 | Yield in plots with [inoculation with of](#) biofertilizer alone (without bio control agent) was also found to be  
122 | at par with the corresponding yield in plots with combined inoculation of biofertilizer and bio-control  
123 | agents. Thus, it may be concluded that for obtaining optimum plant growth and yield from brinjal, the

124 treatment receiving organic manure (FYM) inoculated with biofertilizers may be recommended as there is  
125 no significant difference between the treatment of combined inoculation of biofertilizers + bio-control  
126 agents and that of biofertilizers singly. Moreover, it may be considered as cost-effective treatment, where  
127 there is no chance for the occurrence of diseases as compared to combined treatments because it  
128 involves an extra cost in the application of biocontrol agents. However, in places with some history of  
129 bacterial wilt or related infestation, biocontrol agents may be used-suggested along with biofertilizers.

### 130 REFERENCES:

- 131 1. Horticultural statistics at a glance horticulture. Statistics Division Department of  
132 Agriculture, Cooperation & Farmers Welfare Ministry of Agriculture & Farmers Welfare  
133 Government of India 2017.
- 134 2. Weyens N, Van der Lelie D, Taghavi S, Newman L, Vangronsveld J. Exploiting Plant-Microbe  
135 Partnerships to Improve Biomass Production and Remediation. Trends in Biotechnology, 2009;  
136 27(10): 591-598.
- 137 3. M, Dotaniya M, Mishra A, Dotaniya CK, Regar KL, Manju L. Role of Biofertilizers in  
138 Conservation Agriculture. 2016.10.1007/978-981-10-2558-7\_4.
- 139 4. Sabalpara AN, Panda, JR, Mahatma L. Use of beneficial microbes in agriculture. National  
140 Seminar on role of organic farming in climate resilient and sustainable agriculture. 2014; 75.
- 141 5. Chakravarty G, Kalita MC. Comparative evaluation of organic formulations of *Pseudomonas*  
142 *fluorescens* based biopesticides and their application in the management of bacterial wilt of  
143 brinjal (*Solanum melongena*L.). African Journal of Biotechnology, 2011;10 (37): 7174-7182.
- 144 6. Akiew E, Trevorrow PR, Tonells PE. Management of bacterial wilt of tobacco. In: Bacterial wilt.  
145 Hartman GL and Hayward AC (eds.). ACIAR Proceedings. Australian Centre Int. Agricultural  
146 Res. Camera, 1993 45: 270-275.
- 147 7. Prakasam V, SharmaP. Trichodermaharzianum (Th-3) a potential strain to manage the  
148 purpleblotch of onion (*Allium cepa* L.) caused by *Alternaria porriunder* North Indian plains.  
149 Journal of Agricultural Science. 2012; 10: 266-27.

- 150 8. Jadon, KS. Eco-friendly management of brinjal collar rot caused by *Sclerotium rolfsii*Sacc.  
151 Indian Phytopath. 2009; 62(3): 345-347.
- 152 9. Khalid EE. Biological control of bean damping-off caused by *Sclerotium rolfsii*. Research Gate,  
153 2013; 9: 1-11.
- 154 10. Singh R, Singh PP, Singh V. Integrated management of collar rot of *Amorphophallus*  
155 *paeoniifolius* blume caused by *Sclerotium rolfsii* Saccardo. Vegetable Science. 2006; 33(1): 45-  
156 49.
- 157 11. Harish DK, Agasimani AD, ImamsahebSJ,Patil SS. Growth and Yield Parameters in Brinjal as  
158 Influenced by Organic Nutrient Management and Plant Protection Conditions. Research  
159 Journal of Agricultural Sciences, 2011; 2(2): 221-225.
- 160 12. Samah YAE. Effect of biofertilizer on yield and berry qualities of grapevines. M. Sc. Thesis.  
161 Fac. Agric., Mansoura Univ., Egypt. (2002). (In print)
- 162 13. Singh C, Sharma, BB. Leaf nutrient composition of sweet orange as affected by combined use  
163 of bio and chemical fertilizers. South Indian Horticulture. 1993; 41: 131-134.
- 164 14. Grover MI, Nain L, Saxena A. Comparison between *Bacillus subtilis* RP24 and its antibiotic-  
165 defective mutants. World Journal of Microbiology and Biotechnology. 2009; 25:1329-1335.
- 166 15. Ramakrishnan K, Selvakumar G. Effect of biofertilizers on enhancement of growth and yield on  
167 Tomato (*Lycopersicum esculentum* Mill.). *International Journal of Research in Botany*, 2012.
- 168 16. Bindiya Y, Srihari D, Babu JD. Effect of organic manures and biofertilizers on growth, yield and  
169 nutrient uptake in gherkin (*Cucumis anguria* L.). Journal Research Angra, 2012; 40(1): 26-29.
- 170 17. Dashti\_NH, Ali NY, Cherian VM, Montasser MS. Application of plant growth-promoting  
171 rhizobacteria (PGPR) in combination with a mild strain of *Cucumber mosaic virus* (CMV)  
172 associated with viral satellite RNAs to enhance growth and protection against a virulent strain  
173 of CMV in tomato. Canadian journal of plant pathology. 2012; 34(2):177-186.
- 174 18. Ramesh R, Joshi AA, Ghanekar MP. Pseudomonads: major antagonistic endophytic  
175 bacteria to suppress bacterial wilt pathogen, *Ralstonia solanacearum* in the eggplant  
176 (*Solanum melongena* L.). World Journal of Microbiology and Biotechnology. 2009; 25(1): 47-  
177 55.

- 178 19. Sharma SP, Brar JS. Nutritional Requirements of Brinjal (*Solanum Melongena* L.) – A review.  
179 Regional Station, PAU, Bathinda-151001, India. Agricultural Review, 2008; 29(2): 79-88.
- 180 20. Mirzakhani M, Ardakani MR, Aeene BA, Rejali F, Shirani RAH. Response of spring safflower to  
181 co-inoculation with *Azotobacter chroococum* and *Glomus intraradices* under different levels of  
182 nitrogen and phosphorus. American Journal of Agricultural and Biological Sciences. 2009; 4:  
183 255-261. DOI: 10.3844/ajabssp.2009.255.261.
- 184 21. Nanthakumar S, Veeraraghavathatham D. Effect of integrated nutrient management on growth  
185 parameters and yield of brinjal (*Solanum melongena*L.) cv. PLR-1. South Indian Horticulture,  
186 2000; 48(1-6): 31-35.  
187

Formatted: Font: 10 pt

Formatted: Font: 10 pt, Not Italic

Formatted: Font: 10 pt

Formatted: Font: 10 pt, Not Italic

UNDER PEER REVIEW