

Evaluation of Allelopathic Potential of *Artemisia herba-alba* on Germination and Seedling Growth of *Raphanus sativus* and *Trigonella foenum-graecum*

Hamida M.E. Hamad

Department of Botany, Faculty of Science, Omar El-Mukhtar University, Libya

Abstract:

The allelopathic effects of the aqueous extracts of aerial parts of *Artemisia herba-alba* at different concentrations (20, 40, 80%) were evaluated on germination and seedling growth of *Raphanus sativus* (Radish) and *Trigonella foenum-graecum* (Fenugreek) after 7 days **in vitro**. *Artemisia* had strong allelopathic effects, it prevented the germination and seedling growth of Radish seeds, at highest concentrations (80%). Results obtained showed significant inhibition in germination percentage (GP), plumule length (PL) and radicle length (RL) of both Radish and Fenugreek seeds, and the degree of inhibition was concentration dependent. At 20 and 40% the germination percentage of Radish seed was reduced to (56.67%) and to (40%), while the germination of Fenugreek seeds was reduced to (80 %) and to (63.33 %) respectively. This inhibition was markedly in obvious Radish than in Fenugreek, indicating that is *Raphanus sativus* more sensitive, to the allelopathic effect of the different concentrations of the aqueous extract of the *A. herba alba* plant.

Keywords: Allelopathy; Germination; *Artemisia herba-alba*; Aqueous extract.

1. Introduction:

Allelopathy as an ecological phenomenon, has been defined as any direct or indirect effects of one plant, including micro-organisms, on another through the release of phytochemical compounds into the environment, these biochemical materials are called allelochemicals that may affect the physiological processes of the plants such as respiration, cell division, water and nutrient uptake, oxidative stress and others. Most plant species, including wild plants, crops and trees are capable of producing such molecules into the environment to inhibit the development of neighboring plants. [1,2,3]. the term allelopathy refers to any method involving secondary metabolites(allelochemicals) created by plants, or microorganisms, bacteria, viruses and fungi that influence the growth and development of agricultural and biological systems as well as positive and negative effects. Allelochemicals from plants are produced by any organ of the plant and discharged into the environment by volatilization, exudation from roots, leaching from stems and leaves or decomposition of plant material [4,5]. Allelochemicals are mainly secondary metabolites which usually associated with plant defense against herbivores and pathogens, these distinctive compounds may be linked to wide range of ecological functions [6]. **Phenolic compounds are one in every of the biggest group of secondary metabolites including flavonoids, phenolic acids tannins, saponins, cinnamic acid coumarins, terpenoids, quinones, and lignans [7].** Allelopathic effects can be stimulatory or inhibitory, depending on the identity of the active compound on the static and dynamic availability, persistence and fate of organics in the environment and on the particular target species [8]. Also, allelopathy is generally

Should be corrected "in :[1T]Comment vitro"

this description is :[2T]Comment unnecessary in the subject flow

accepted as a significant ecological factor in determining the plant growth, succession, dominance, distribution, species diversity, structure and composition of plant communities [9]. *Artemisia herba-alba* Asso. (Asteraceae), which recently classified as subgenus *Seriphidium* [10], commonly known as the desert wormwood, is a dwarf, semi shrub growing widely in Al-Gabal Al-Akhdar in Libya and in, Northern Africa (Morocco, Tunisia Algeria), the Middle East Western Asia (Arabian Peninsula) and Southwestern Europe. The plant is a perennial, strongly aromatic herb, with many basal, erect and leafy stems covered with woolly hairs [11]. It is widely used as folk medicine and in particular for common uses such as relief of coughing, intestinal disturbances, colds and muscle tensions by the local population in different countries [12]. The allelopathic consequences of the genus *Artemisia* have been broadly investigated [13,14,15]. *Artemisia herba-alba* grown in Libya was observed as dominant species in its natural micro habitat in Al-Gabal Al-akhdar area and prevents growth of another plant species those grow closely to it. Therefore, we hypothesized that this species has allelopathy effects and can be used as determination factor for the growth of some plants. Therefore, the study was aimed to evaluate the **Allelopathic Effect of *Artemisia herba-alba* (*Artemisia*) on Germination and Seedling Growth of *Raphanus sativus*(Radish) and *Trigonella foenum-graecum* (Fenugreek) .**

2. Material and Methods:

2.1. Sources of the seeds:

The seeds of Radish and Fenugreek **were obtained from local market**. And were kept in the containers which they were supplied, then stored in the laboratory at room temperature until required for sowing.

2.2. Plant Material Collection

Artemisia. herba-alba samples were collected from the south of Al-Jabal Al-Akhdar EL-Byda city – Libya (Marawa region) , in October 2018, collected only areal parts. The plant were classified and authenticated according to [16]. **In Department of Botany, Faculty of Science, Omar Al-Mukhtar University, EL- Byda, Libya.**

2.3. Preparing the aqueous extract of *Artemisia herba-alba*

A number of fresh samples from the aerial shoots of the donor species were collected from the natural habitats in the study area during the vegetative stage. The samples were air-dried then after ground in a Wiley Mill to fine uniform texture and stored in glass jars until use. Stock aqueous extract was obtained by soaking 50 g air-dried plant material in 500 ml of cold distilled water (10% w/v) at room temperature ($20 \pm 2^\circ$ C) for 24 hours with **occasional shaking**. **the** mixture was left on shaker for 24 h in room temperature at speed of 120 rpm four-folded cotton fabric was used as filter to separate rough solid particles from solution. The contents were then filtered with Whatman filter paper and then it was **centrifuged** **with** the speed of 2000 rpm for 15 minutes [17].

2.4. Preparing the aqueous extract concentrations

Three concentrations of solutions (20, 40, and 80 %) were prepared from the stock solution based on volume/volume **percent(v/v,%)** [18]. **in** addition to the distilled water as control. To prepare solutions of different concentrations, doses of *A. herba-alba* aqueous extract (20, 40 and 80 ml) were taken. Then, volume was completed to 100 ml by adding distilled water to obtain (20, 40, and 80 %).

Use of dot in parentheses is **: [3T]Comment**
incorrect

Word of the first letter **: [4T]Comment**
should be written in small letters

where were the seeds from **: [5T]Comment**
the local market? Which city?

there are shortcomings in **: [6T]Comment**
this sentence

What is the mark of shaker? **: [7T]Comment**

Word of the first letter **: [8T]Comment**
should be written in capital letters

What is the mark of **: [9T]Comment**
centrifuge?

Words and parentheses **: [10T]Comment**
must be written separately.

Word of the first letter **: [11T]Comment**
should be written in capital letters

2.5. Treatment of seeds with plant extracts:

Sixty seeds of each species(**Radish and Fenugreek**) were distributed in 12 petri dishes (5 seed in each dish) on two layers of Whatman No.1. Five ml of each prepared aqueous extract (20, 40, and 80 %) or distilled water as control (0%) were added in petri dishes (added daily). Three replicates were incubated in randomized complete block design at 20° C in an incubator. Before sowing, the seeds were surface sterilized with 2% sodium hypochlorite for 2 minutes then rinsed four times with distilled water. The sterilized seeds were soaked in aerated distilled water for 24 hours. The germination percentage (GP), plumule length (PL) and radicle length (RL) were recorded after one week at the end of the experiment [19]. A variety of parameters were used in this work to assess the effects of extracts on seed germination and seedlings development of test species. These parameters include:

1. Germination percentage (GP): was calculated according to the following equation [20].

$$\% \text{Germination} = \frac{\text{Number of germinated seeds}}{\text{Total number of seeds}} \times 100$$

2. plumule length (PL) and radicle length (RL):

Length of plumule and radicle system were measured in cm using a ruler.

Statistical Analysis:

Statistical analysis was performed using a computer run program (Minitab software). One way ANOVA followed by Tukey, s HSD test was performed to show the statistical significance among the means of the groups. Results were expressed as mean ± Standard Division (SD). P-value below 0.05 was considered to be statistically significant [21].

3.RESULTS:

3.1. The allelopathic effect of Artemisia on Radish seeds:

1. Germination percentage (GP): The germination percentage of Radish seeds was significantly affected by the increase at different concentration levels of **Artemisia aqueous extract** after seven days of germination. **Artemisia herba-alba** extracts have inhibitory effect on germination and early growth of *Raphanus sativus* and considerably suppressed the germination compare to control treatment. The total percentage of Radish seeds germination was decreased by increasing the concentration of Artemisia aqueous extract, at control (0%) GP value was about (93.33 %). While the percentage was reduced to (56.67%) and to (40 %), at concentrations 20 and 40% *Artemisia herba-alba* aqueous extract, respectively. The maximum allelopathic effect was recorded in 80% Artemisia aqueous extract concentration, which completely inhibited Radish seed germination (Table1).

2. Plumule length (PL): Findings of PL of *Raphanus sativus* imply the inhibitory effect of the allelopathic substances on seedling stage. Evidently, PL was significantly reduced due to each main effect as treatment Additionally, the value of PL was 4.2 cm at control level. Afterward, it reduced to 2.7cm at 20 %, and to 2.2 cm at 40 %, the

Words and parentheses :[12T]Comment must be written separately.

Word of the first letter :[13T]Comment should be written in capital letters

Word of the first letter :[14T]Comment should be written in capital letters

It should be written in the :[15T]Comment same way in manuscript. either Artemisia aqueous extract or Artemisia herba-alba should be used to ensure the standard

no point mark is required :[16T]Comment before parentheses

Word of the first letter :[17T]Comment should be written in capital letters

maximum allelopathic effect was recorded in 80% Artemisia aqueous extract concentration, which completely inhibited **PL** (Table1).

no point mark is required :[18T]Comment before parentheses

3. Radicle length (RL): decreased was observed among Radish RL assessment in seeds culture (table 1). The control value was 6.1cm . Elevated *A. herba-alba* aqueous extracts concentrations have possessed a significant inhibitory effect on radical growth. At 20 % *A. herba-alba* aqueous extracts concentration, it was 2.3cm. Upon applying the highest *A. herba-alba* aqueous extracts concentration (80%), it was completely inhibited **RL** (Table1).

no point mark is required :[19T]Comment before parentheses

Table1. Allelopathic effect of different concentrations of aqueous extract of *Artemisia herba-alba* on germination percentage(GP) and radicle (RL) and plumule(PL) length (cm) of *Raphanus sativus* L. (after 7 days).

Extract concentration %	Seed germination Mean \pm SD	GP %	RL(cm) Mean \pm SD	PL(cm) Mean \pm SD
0	18.67 ^a \pm 0.47	93.33	6.1 ^a \pm 0.09	4.2 ^a \pm 0.16
20	11.33 ^b \pm 0.94	56.67	2.3 ^b \pm 0.08	2.7 ^b \pm 0.14
40	8 ^c \pm 0.82	40	1.5 ^c \pm 0.05	2.2 ^c \pm 0.12
80	0 ^d	0	0 ^d	0 ^d

Data are expressed as mean \pm SD of three replicate. Within each row, means with different superscript (a, b, c or d) were significantly different at $p < 0.05$. Where means superscripts with the same letters mean that there is no significant difference ($p > 0.05$).

3.2. The allelopathic effect of Artemisia on Fenugreek seeds:

1. Germination percentage (GP): The allelopathic effects of *A. herba-alba* aqueous extracts on Fenugreek seeds were also evaluated, the total percentage of Fenugreek seed germination was decreased by increasing the concentration of Artemisia aqueous extract, at control (0%) GP value was about (98.33%) .The percentage was reduced to (80 %) at 20 % and to (63.33 %) at 40%, while recorded (46.67%) at (80% v/v) *A. herba-alba* aqueous extracts concentration. (Table 2).

2. Plumule length (PL): Evaluation of PL correlated with higher Artemisia aqueous extract concentrations has demonstrated their inhibitory influence on *Trigonella foenum-graecum* growth process (Table 2). The plumule elongation was not completely inhibited by the extract but it was less at higher concentration levels. Obviously, all allelopathic concentrations have reduced PL. at control level PL of Fenugreek was 5.9 cm. On the other hand, 20, 40 and 80 % concentrations were considered as inhibitory concentrations (the values about 4, 3.6 and 2.1cm) respectively.

3. Radicle length (RL): Compared to control, a gradual decrease in RL of Fenugreek seed was observed along gradual increase in *A. herba-alba* aqueous extracts concentrations. RL implication was significantly affected by the treatment. At control, the values of RL were 4.4 cm. higher concentrations of *A. herba-alba* aqueous extracts were notably active inhibiting radicle emergence. at 20, and 40 % concentrations, RL decreased to 3.5 and 3.2 cm respectively. Constantly, it continues reduction till it attained a value of about 1.8 cm at 80 % concentration level. (Table 2).

Table 2. Allelopathic effect of different concentrations of aqueous extract of *Artemisia herba-alba* on germination percentage (GP) and radicle (RL) and plumule (PL) length (cm) of *Trigonella foenum-graecum*. (after 7 days).

Extract concentration %	Seed germination Mean \pm SD	GP %	RL(cm) Mean \pm SD	PL(cm) Mean \pm SD
0	19.67 ^a \pm 0.47	98.33	4.4 ^a \pm 0.24	5.9 ^a \pm 0.09
20	16 ^b \pm 0.82	80	3.5 ^b \pm 0.05	4 ^b \pm 0
40	12.67 ^c \pm 1.2	63.33	3.2 ^b \pm 0.12	3.6 ^b \pm 0.21
80	9.33 ^d \pm 0.94	46.67	1.8 ^c \pm 0.16	2.1 ^c \pm 0.14

Data are expressed as mean \pm SD of three replicate. Within each row, means with different superscript (a, b, c or d) were significantly different at $p < 0.05$. Where means superscripts with the same letters mean that there is no significant difference ($p > 0.05$).

4. Discussion:

Environmental and non-environmental stresses lead to the interactions in plants. Some of environmental stresses are allelopathic compounds which secrete by some plants and cause disturbance in life cycle and activate a series of biochemical reactions [22]. The present work was carried out as a study to investigate any possible allelopathic activity *A. herba alba* aqueous extract on germination and seedling growth of *Raphanus sativus* and *Trigonella foenum-graecum*, the results showed severe toxicity at high extract concentration and moderate toxicity at low concentration. The highest germination rate of Radish and Fenugreek seeds was obtained from distilled water treatment and the lowest rate was obtained from treatments lead to lack of germination, respectively. In general the results showed that when concentration of extract increases, traits significantly decrease, this can result from the increase in amount of allelochemicals and the toxicity characteristics [23]. *A. herba alba* aqueous extract may contain some phytotoxic substances that inhibits germination and growth of Radish and Fenugreek. These results correlated with the findings that Allelochemicals presented in the aqueous extracts of different plant species commonly identified as allelopathic agents, which have inhibitory and/or lethal effects on seed germination growth and development, reduction in seedling growth and have been reported to effect on different physiological processes through their effects on enzymes responsible for phytohormone synthesis and were found to associate with inhibition of nutrients and ion absorption by affecting plasma membrane permeability [24,25]. Aqueous extract of plants may contain phenolics such as ferulic acid P-coumaric, vanillic, caffeic, chlorogenic and others [26,27]. These phenolics inhibit the germination process [28,29]. which was due to their interference with indol acetic acid metabolism, or synthesis of protein and ion uptake by the plants [26]. Therefore, *A. herba alba* might release some soluble phenolic allelochemicals to the environment [30], which has a growth inhibitory effect on new seedling of both *Raphanus sativus* and *Trigonella foenum-graecum* or other plant species. Our results are agreed with [31,32] who reported that seeds of some species can be suppressed using water extracts from *Artemisia* plants or another species and these extracts can effect on germination behavior too. The results of this study

showed that *Artemisia* extracts had deterrent effects on the germination and growth indices of *Raphanus sativus* and *Trigonella foenum-graecum*. The seed germination traits and seedling growth were decreased by increasing the extract concentration.[18] concluded that *A. herba alba* aqueous extract at different concentrations suppressed the germination of monocot (wheat) and dicot (tomato) seeds. And this suppression was possibly due to the presence of allelochemicals in this plant. There are some reports about the inhibitory effects of different species of *Artemisia* on seed germination traits of *Triticum aestivum* L., *Brassica napus*, *Sinapis arvensis* L. [33] *Amaranthus retroflexus* L, and *Convolvulus arvensis* L. [34] *Atriplex canescens*, *Agropyron elongatum* and *Agropyron desertorum* [35]. According to above researches, it can be firmly concluded that genus *Artemisia* forms the plants whose allelopathic ability is proved between different species. In this genus, a wide range of active biological compounds are produced which included artemisinin, tannin, flavonoids, sesquiterpene lactone and other secondary metabolites such as coumarin, camphor and bornyl acetate which their toxicity for some other plants is proved [36,37,38] Coumarin prevents the cell from entering the as the first group of mitochondrial mitosis. Flavonoids have been introduced absorption inhibitor that may stop ATP production in mitochondria and affect the breathing [39]. Through preventing from the cell division and cell elongation in the germination stage, flavonoids and coumarin deter germination and reduce the length of root and shoot of the seeds.

Species names should be :[20T]Comment
written in italics

5. Conclusion:

It can be concluded that *Artemisia herba alba* water extract at different concentrations suppressed the germination of *Raphanus sativus* and *Trigonella foenum-graecum* seeds. And this suppression was possibly due to the presence of allelochemicals in this plant. Based on the results of this study: *Artemisia herba alba* species have strongest allelopathic potential must be examined for their selective action on other specific plants including weeds and crops under field conditions, their allelopathic activity will be much more detailed. Analysis of possible allelochemicals in this plant is also required. The isolation and characterization of growth inhibitors, which might be responsible for the strong allelopathic potential of this species is needed. There is possibility of using these allelochemicals directly or as structural leads for the discovery and development of environment friendly herbicides to control weeds.

Artemisia herba-alba :[21T]Comment

Artemisia herba-alba :[22T]Comment
Species names should be written in italics

6. References :

- 1.Rice, E.L. Allelopathy. 2nd. edn. Orlando, Academic Press.1984 ; pp. 422.
- 2.Weir TL, Park SW, Vivanco JM. Biochemical and physiological mechanisms mediated by allelochemicals. Current Opinion in Plant Biology. 2004; 7:472-479.

3. Inderjit, Callaway RM, Vivanco JM. Can plant biochemistry contribute to understanding of invasion ecology. Trends Plant Sciences. 2006; 11:574-580.

4. J.V. Lovett and M.Y. Ryuntyu . In "Allelopathy: Basic and Applied Aspects" Edited by S.J.H. Rizvi, V. Rizvi Chapman & Hall, London. 1992 ; pp. 11-20.

5. Rizvi, S.J.H. and Rizvi, V. (Eds.). "Allelopathy: Basic and Applied Aspects". Chapman & Hall, London. 1992; pp. 480.

6. Hussain I.M and Reigosa J.M. Allelochemical stress inhibits growth, leaf water relations, PSII photochemistry, non-photochemical fluorescence quenching and heat energy dissipation in three C3 perennial species. Journal of Experimental Botany. 2011; 62(13): 4533-4545.

7. Balasundram N, Sundram K, Samman S. Phenolic compounds in plants and agri-industrial by products: antioxidant activity, occurrence and potential uses. Food Chemistry. 2006; 99: 191-203.

8. Inderjit, K. and I. Keating. Allelopathy: principles, procedures, processes, and promises for biological control . Advances in Agronomy. 1999 ; 67:141-231.

9. L.R. Scrivanti, M.P Zunino, J.A. Zygadlo. **Tagetes minuta and Schinus areira** essential oils as allelopathic agents. Biochemical Systematic and Ecology. 2003; 31:563-572.

Species names should be :[23T]Comment
written in italics

10. Torrell, M. Cerbah, M., Siljak-Yakovlev, S. And Vallès, J. Molecular cytogenetics of the genus *Artemisia* (Asteraceae Anthemideae) fluorochrome banding and fluorescence in situ hybridization. I. Subgenus *Seriphidium* and related taxa. Plant Systematic and evolution. 2003 ; 239, 141- 153.

11. Alavi, S. A. Elfateh university, Tripoli, Libya, Fl. Lib. 1983 ; 107, 180-180.

There is a deficiency in :[24T]Comment
reference writing

12. Oran, S. A. and D.M. Al-Eisawi. Check-List of **Medicinal Plants** in Jordan. Dirasat Medicine and Biology Science. 1998 ; 25(2), 84-112.

Word of the first letter :[25T]Comment
should be written in small letters

13. Barney, J.N., Hay, A.G. and Weston, L.A. () Isolation and characterization of allelopathic volatiles from mugwort (*Artemisia vulgaris*), Journal of Chemistry and Ecology. 2005 ; 31(2), 247-65.

14. Delabays, N. *Artemisia annua* L. a traditional Chinese medicinal plant for a modern weed management. Report, Swiss Federal Research Station for Plant Production, Switzerland. 2001;

15. Simon, J.E., Charles, D., Cebert, E., Grant, L., Janick, J. **And** Whipkey, A. *Artemisia annua* L.: A promising aromatic and medicinal, In: J. Janick and J.E Simon (eds.), Advances in **new crops. Timber** Press Portland, OR, 1990; p. 522-526.

Word of the first letter :[26T]Comment
should be written in small letters

Word of the first letter :[27T]Comment
should be written in capital letters

16. Jafri , S. and El-Gadi , A. Asteraceae . In flora of Libya , 1978; 107. Al-Fateh University Press, Tripoli, Libya.

17. Bajalan I., Zand M., and Rezaee S. Allelopathic effects of aqueous extract from *Salvia officinalis* L. on seed germination of barley and purslane. International Journal of Agricultural Crop Sciences ., 2013; 5(7): 802-805.

Species names should be :[28T]Comment
written in italics

18. Elshatshat S. A. Allelopathic Effects of *Artemisia Herba-Alba* Aqueous Extracts on Germination of Tomato and Wheat Seeds. Journal of Science and Its Applications. 2010; 4(1): 1-6.

Word of the first letter :[29T]Comment
should be written in small letters

19. Salhi, N., El-Darier ,S. M. and Halilat M. T. Allelopathic Effect of some Medicinal Plants on Germination of two Dominant Weeds in Algeria. Advances in Environmental Biology. 2011; 5(2): 443-446.

Word of the first letter :[30T]Comment
should be written in small letters

20. Scott, S.J., Jones R.A. and Williams W.A. Review of data analysis methods for seed germination. Crop Science. 1984 ; 24: 1192-1199.

21. Ross SM. Introduction to probability and statistics for engineers and scientists .3rd ed. USA: Elsevier. 2004;

22. Saberi, M., Tavili, A. and Shahriari, A. R.,. The influence of chemical stimulators on decrease of *Thymus kotschyianus* allelopathic effect on *Agropyron elongatum* seed germination characteristics. Watershed Management Research (Pajouhesh& Sazandegi . 2012 ; 95: 45-54. (In Persian).

23. Kohli, R.K., H.P. Singh and D.R. Batish,. Allelopathy in Agroecosystems. Journal of Crop Production. 2008; 4(2):1-41.

Word of the first letter :[31T]Comment
should be written in small letters

24. Daizy R.; B. K. Manpreet; P. S. Harminder and K. K. Ravinder . Phytotoxicity of a medicinal plant, *Anisomeles indica*, against *Phalaris minor* and its potential use as natural herbicide in wheat fields. Crop Protection. 2007; 26(7): 948-952.

Species names should be :[32T]Comment
written in italics

25. Fateh E, Sohrabi SS, Gerami F. Evaluation of the allelopathic effect of bindweed (*Convolvulus arvensis* L.) on germination and seedling growth of millet and basil. Advances in Environmental Biology. 2012; 6:940-950.

Species names should be :[33T]Comment
written in italics
Should be corrected "*Phalaris minor*"

: [34T]Comment

26. Hussain, F. and T.W. Khan,. Allelopathic effects of Pakistani weed *cynodon dactylon* L. Journal of Weed Science Research. 1988; 1:8-17

27. Habib, S.A. and A.A. Abdul Rehman,. Evaluation of some weed extract against dodder on alfalfa (*Medicago sativa*). Journal of chemical Ecology. 1988 ;14:443-452.

28. Williams, R.D. and R.E. Hoagland,. The effect of naturally occurring phenolic compounds on seed germination. Weed Science. 1982 ; 30:206-212.

29. Al-Charchafchi, F.M.R., F.M.J. Redha and W.M. Kamal. Dormancy of *Artemisia herba alba* seeds in relation to endogenous chemical constituents. Journal of Biological Science Research. Baghdad/Iraq. 1987; 2:1-12.

30. Xu, Z., D. Yu, L. Guo, M. Zhao, X. Li, D. Zhang, K. Ye and Y., Zheng., Molecular biological study on the action mechanism of rice allelochemicals against weed Ying Yong Sheng Tai Xue Bao. 2003; 14:829-833.

31. Escudero, A., Albert, M., Pita, J. M. and Pérez-García, F. Inhibitory effects of *Artemisia herba-alba* on the germination of the gypsophyte *Helianthemum squamatum*. Vegetation. 2000; 148 (1): 71-80.

32. Periotto, F., Juliano, S., C., Gualtieri, M., Lima, S., and Perez, A. Allelopathic effect of *Andira humilis* Mart. ex Benth in the germination and growth of *Lactuca sativa* L. and *Raphanus sativus* L. Acta Botanica Brasilica. 2004; 18 (3) .

33. Akramghaderi, F., Zeinali, E. and Farzaneh, S. Allelopathic effects of annual wormwood *Artemisia annua* L on seedling emergence and growth of wheat, oil seed rape, wild mustard and wild oat. Journal of Agricultural Sciences and Natural Resource. 2001; 8(3): 113-121 .

34. Tabatabaee Zade, M.S., Pajouhan, M., Soltani M., Tajamolian, M. and Shahbandari, R. Allelopathic Effects of *Artemisia aucheri* boiss **Essential Oils on Seed Germination and Early Seeding Growth of Red-root** Amaranth, (*Amaranthus retroflexus* L.) and **Field Bindweed** (*Convolvulus arvensis* L.) **Knowledge of sustainable agricultural production.** 2014; 24(3): 87- 95. (In Persian).

35. Bagheri, R. and Mohammadi, S. Allelopathic effects of *Artemisia sieberi* Besser on three important species (*Agropyron desertorum*), (*Agropyron elongatum*) and (*Atriplex canescens*) in range improvement. Iranian Journal of Range Desert Research. 2011; 17(4): 538-548. (In Persian).

36. Lydon J., Teasdale, J.R. and Chen, P.K. Allelopathic activity of annual wormwood (*Artemisia annua*) and role of artemisin. Weed Science. 1997 ; 45:807-811..

37. Klyman, D.L., Qinghaosu (artemisin): an antimalaria drug from **china**. Science. 1985; 228: 1049-1055.

38. Macro, J.A. and Babera, O., Natural products from the gents Artemisia I stud. Natural Products . 1990; 7: 201-264.

39. Maighany, F., Allelopathy: From concept to application. Parto e Veghe, Tehran Iran. 2003; 256 pp. (In Persian).

Word of the first letter :[35T]Comment
should be written in small letters

Word of the first letter :[36T]Comment
should be written in capitall letters

Word of the first letter :[37T]Comment
should be written in capitall letters

reference writing is :[38T]Comment
incorrect