

Editor's Comment:

The manuscript may be acceptable with minor revision.

Comments:

1. Line 156: "productin China" may be corrected as "product in China".
2. Lines 185-186: "We have successfully amplified and purified of CsSUBP gene in cucumber coding sequence, and constructed the recombinant prokaryotic expression vector pET30a-CsSUBP." may be corrected as "We have successfully amplified and purified the coding sequence of CsSUBP gene in cucumber, and constructed the recombinant prokaryotic expression vector pET30a-CsSUBP."
3. The presentations of references in the References Sections need to be corrected. The corrections suggested are marked in red as those shown below.

REFERENCES

1. Hasama W, Morita S, Kato T. Seed transmission of *Corynespora melonis*, causal fungus of target leaf spot, on cucumber . *Jpn J Phytopathol*. 2009; 59(2):175-9.
2. Qi YX, Zhang X, Pu JJ, et al. Morphological and molecular analysis of genetic variability within isolates of *Corynespora cassiicola* from different hosts . *Eur J Plant Pathol*. 2011; 130(1):83-95.
3. Siezen RJ, De WV, Leunissen JA, et al. Homology modelling and protein engineering strategy of subtilases, the family of subtilisin-like serine proteinases . *Protein Engi*.1991;4(7): 719.
4. Ksiazek M, Karim AY, Bryzek D, et al. Mirolase, a novel subtilisin-like serine protease from the periodontopathogen *Tannerella forsythia*. *Biol Chem*. 2015; 396(3): 261-75.
5. Duan X, Zhang Z, Wang J, et al. Characterization of a novel cotton subtilase gene *GbSBT1* in response to extracellular stimulations and its role in *Verticillium resistance*. *PLoS One*. 2016; 11(4): e0153988.
6. Meyer M, Huttenlocher F, Cedzich A, et al. The subtilisin-like protease SBT3 contributes to insect resistance in tomato. *J Exp Bot*. 2016;67(14): 4325-38.
7. Mizutani O, Shiina M, Yoshimi A, et al. Substantial decrease in cell wall α -1, 3-glucan caused by disruption of the *kexB* gene encoding a subtilisin-like processing protease in *Aspergillus oryzae*. *Biosci Biotechnol Biochem*. 2016; 80(9): 1781-91.
8. Pepe A, Frey ME, Munoz F, et al. Fibrin (ogen) olytic and antiplatelet activities of a subtilisin-like protease from *Solanum tuberosum* (StSBTc-3). *Biochimie*. 2016;125: 163-70.

9. Figueiredo A, Monteiro F, Sebastiana M. Subtilisin-like proteases in plant-pathogen recognition and immune priming: a **perspective**. *Front Plant Sci*. 2014;5(739): 739.
10. Vartapetian AB, Tuzhikov AI, Chichkova NV, et al. A plant alternative to animal caspases: subtilisin-like proteases. *Cell Death Differ*. 2011; 18(8): 1289.
11. Rantong G, Evans R, Gunawardena AH. Lace plant ethylene receptors, AmERS1a and AmERS1c, regulate ethylene-induced programmed cell death during leaf **morphogenesis**. *Plant Mol Biol*. 2015; 89(3): 215-27.
12. Coffeen WC, Wolpert TJ. Purification and characterization of serine proteases that exhibit caspase-like activity and are associated with programmed cell death in *Avena sativa*. *Plant Cell*. 2004; 16(4): 857-73.
13. Ramírez V, López A, Mauch-Mani B, et al. An extracellular subtilase switch for immune priming in *Arabidopsis*. *PLoS Pathog*. 2013;9(6): e1003445.
14. Zhang B, Pan X, Cannon CH, et al. Conservation and divergence of plant microRNA genes. *Plant J*. 2006; 46(2): 243-59.
15. Zhang B, Wang Q, Wang K, et al. Identification of cotton microRNAs and their targets. *Gene*. 2007; 397(1): 26-37.
16. Yin Z, Li C, Han X, et al. Identification of conserved microRNAs and their target genes in *tomato (Lycopersicon esculentum)*. *Gene*. 2008; 414(1): 60-6.
17. Sunkar R, Jagadeeswaran G. In silico identification of conserved microRNAs in large number of diverse plant **species**. *BMC Plant Biol*. 2008; 8(1):37.
18. Yang M. Study on the Diversity of Pathogens of Coryneform Leaf Spot in Vegetables in China. Chinese Academy of Agricultural Sciences. 2013.
19. Tanaka H, Onouchi H, Kondo M, et al. A subtilisin-like serine protease is required for epidermal surface formation in *Arabidopsis* embryos and juvenile plants. *Development*. 2001;128(23): 4681-9.
20. D' Erfurth I, Signor CL, Aubert G. A role for an endosperm-localized subtilase in the control of seed size in legumes. *New Phytol*. 2012;196(3): 738-51.
21. Othman R, Nuraziyana A. Fruit-specific expression of papaya subtilase gene. *Plant Physiol*. 2010;167(2): 131-7.
22. Roberts IN, Caputo C, Kade M, et al. Subtilisin-like serine proteases involved in N remobilization during grain filling in **wheat**. *Acta Physiol Plant*. 2011;33(5): 1997-2001.
23. Hong LI, Zhang Z. Systems understanding of plant–pathogen interactions through genome-wide protein–protein interaction networks. *Front Agr Sci Eng*. 2016;3(2):102-12. DOI: [10.15302/J-FASE-2016100](https://doi.org/10.15302/J-FASE-2016100)
24. Tornero P, Conejero V, and Vera P. Identification of a new pathogen-induced member of the subtilisin-like processing protease family from plants. *Biol Chem*. 1997; 272(22):14412-9.
25. Yamaguchi Y, Barona G, Ryan CA, et al. GmPep914, an eight-amino acid peptide isolated from soybean leaves, activates defense-related genes. *Plant Physiol*. 2011;156(2): 932-42.

26. Fan T, Bykova NV, Rampitsch C, et al. Identification and characterization of a serine protease from wheat leaves. *Eur Plant Pathol.* 2016; 146(2): 293-304.
27. Rong JJ, Diao ZY, Zhou GH. Advances in the development of the *E. coli*. *Drug Biotechnol.* 2005;12(6):416-20.

Editor's Details:

Dr. Kuo-Kau Lee
Professor,
Department of Aquaculture, National Taiwan Ocean University, Taiwan