

Original Research Article

Ageing induced hyperproduction of reactive oxygen species and dysbalance in enzymatic link of antioxidant defense system of skin and therapeutic efficacy of artichoke extract

ABSTRACT

Skin aging is multitarget persistence processing that immediately involve hyperproduction of free radicals under influence of intrinsic and extrinsic factors and deterioration in intimal antioxidant defense system. The goal of the study was the evaluation of the anti-oxidant potential of artichoke standartized extracts, 2%, as a protective strategy against skin age-associated oxidative damage caused by D-galactose (D-gal) in rats. 58 female Wistar rats included in the experimental design. D-gal-induced aging was reproduced in 36 animals of main group, and 12 rats included in control group. All animals in main group were randomized for 3 groups: I – animals with skin aging reproduced model receive saline, II – animals with skin aging rats receive artichoke extracts (with content of chloroogenic acid 2.0%) in a dose of intradermal injection 0.13 mg/kg and main III group - animals with skin aging receive 1.3 mg/kg artichoke extract twice at weeks during 4 weeks. Influence of artichoke extracts restores skin relative weight and leads to decreasing the rate of generation of superoxide anion, hydrogen peroxide and lipid peroxidation (LPx), increasing activity of superoxide dismutase (SOD), glutathione peroxidase (GSH-Px) and reverse ratio SOD/(catalase+GPx) to the production of H₂O₂ from superoxide dismutation coupling with the decrease ratio of generated O₂⁻/H₂O₂. Low-dose of intradermally microinjection of artichoke extracts, 2%, activated the enzymatic link in innate antioxidant defense system in D-gal-induced skin aging model and could be recommended for applications in cosmetics as antiaging mesotherapy.

26 *Key words: skin, aging, artichoke extract, oxidant defense system, superoxide anion, glutathione*
27 *system.*

28 **Abbreviation.** GSH - Reduced glutathione, GSSG - oxidized glutathione, GSH-Px – glutathione
29 peroxidase, lipid peroxidation MDA - Malone aldehyde, Mt –Mitochondrial, ROS - Reactive
30 oxygen species, SOD – superoxide dismutase

31

32 **1. INTRODUCTION**

33 Intrinsic skin aging process mainly includes gloomy skin, relaxation, moisture reduction,
34 thinning, is an inevitable spontaneous process and complex natural phenomenon characterized
35 aging [1-4]. More popular hypothesis that at the molecular level aging is multifactorial gradual
36 biological process associated with diminishes homeostasis, mitochondrial DNA (mtDNA)
37 damage, and progressive decline of innate defense systems of the body, and endogenous
38 antioxidant defense system and oxidative stress formation, particularly [6-7]. Free radical and
39 mitochondrial theories of aging supported by estimation of positive relation between the **sings** of
40 aging and progression of imbalance of free radical metabolism and oxidative damage affects
41 replication and transcription of mtDNA, which closely accompanied the structure and function
42 deterioration in energy supply systems of tissues and organs of the aging and age-related
43 diseases. The decline or/and disturbances of energy supply system functioning leads to increased
44 mitochondrial reactive oxygen species (ROS) generation, ROS-induced lipid peroxidation in
45 mitochondrial membranes and release of **cytochrom C**. These together with antioxidant defense
46 systems imbalance results in further greater overproduction of ROS and to a vicious cycle of
47 premature cellular senescence, skin aging and aged related diseases [4,5,8]. As a model for
48 pharmacological studies of age-dependent alterations in skin we have choice one of the most
49 widely used and demonstrated to display similar symptoms to those aging naturally D-galactose
50 (D-gal)-treated animal model [9-14]. At high levels, D-gal, an aldohexose,
51 monosaccharide sugar, is a naturally occurring substance in the body, which is completely

Formatted: Font color: Red

Formatted: Font color: Red

52 metabolized at normal concentrations and induced disruption in carbohydrate metabolism
53 pathway and causes oxidative stress via stimulation of free radical production and accumulation,
54 apoptosis and inflammation in beyond normal concentration [8-10]. In according to one of the
55 hypothesis that expressive administration of D-gal could induced damage associate with
56 mitochondrial dysfunction caused by complex I deficiency [8-10, 14] and can accelerate ageing
57 was suggested and then confirmed in experimental and clinical data. In order to evaluated the
58 molecular mechanism involved in the controlling of oxidative stress formation we firstly
59 investigated the formation of superoxide anion and hydrogen peroxide and activity of much
60 important components of enzymatic part of antioxidant defense system in D-gal induced skin
61 aging model in experimental animals. Early in clinical practice [16-19] and in experimental
62 studies [20-26], it was shown antioxidant and antitoxic activities [27-28], glycemia-lowering
63 effect [21-22, 24, 28-30], and etc. of artichoke extracts, 5%, but therapeutic properties of
64 artichoke leaves extract on the skin aging process practically have not been investigated. In this
65 study, we examined the possible mesotherapeutic potential of artichoke (*Cynara scolymus* L.
66 (Asteraceae), folium) extract, 2%, to decline the deterioration in skin oxidant defense system in
67 experimental model of skin aging.

Formatted: Font color: Red

68

69 2. MATERIALS AND METHODS.

70

71 2.1. Plant materials and Authentication

72 The fresh leaves of the artichoke *C. cardunculus* L. var. *scolymus* (L.), family Aesreraceae, were
73 collected at harvest maturity from the June to the middle of October during the 2016-17 years in
74 Mtskhethis region (Rosenthal, Georgia, latitude 41° 56' 02" N and longitude 44° 34' 36" E),
75 average minimum temperature -1°C and maximum 35°C. The plant was identified at the
76 Pharmaceutical Natural Sciences Department of Institute of Pharmacy of Sechenov First
77 Moscow State Medical University (Sechenov University)

Formatted: Font color: Red

78

79 **2.2. Preparation of plant extracts and its toxicity study**

80 The leaves the artichoke were separated, washed, cleaned, and drying in according with Eur Ph
81 monograph 01/2008:1866 corrected 6.0. Extraction of dried leaves artichoke, separation and
82 identification of volatiles artichoke was prepared in according with Eur Ph monograph
83 01/2009:2389 (content of chlorogenic acid <2,5%) as described early [31]. The studying extracts
84 of artichoke, 2%, in ampoule was characterized by the content of chloroagenic acid 1.95%
85 (related to the requirements of assessment report on *C. scolymus*, folium
86 EMA/HMPC/150209/2009), total phenolic content equal 0,31±0.04 mg gallic acid
87 equivalent/100 mg extract, total flavonoids 1.6% and total antioxidant activities determinate as
88 50% inhibition of 1,1-diphenyl-2-picrylhydrazyl (DDPH) 15.1±0.9%). The toxicity of **studding**
89 artichoke extracts under i.p. administration is very low, LD₅₀ exceeds 1g/kg body weight and no
90 rats exhibited visible signs of toxicity under 14 days of intradermal injection of extracts of
91 artichoke, 2% including absence of physiologically changes in skin and fur, eyes or mucous
92 membranes. Moderately irritating reactions induced by extracts of artichoke, observed at
93 concentration more than 10% and extracts of artichoke, 2% shows good skin compatibility in
94 patch test [31].

Formatted: Font color: Red

Formatted: Font color: Red

95

96 **2.3. Animals and experimental study design.**

97 **2.3.1. Ethical statement**

98 Animals received humane care in compliance with “Guide for the Care and Use of Laboratory
99 animals” (National Institutes of Health publication 86-23, Revised 1996) and performed with
100 approval of the local Interinstitutional (International Scientific Centre of Introduction of New
101 Biomedical Technology, Department of Medical Pharmacology and Pharmacotherapy, Tbilisi
102 State Medical University, Tbilisi) Animal Care and Use Committee. All animals secured under
103 specific pathogen free conditions according to the Federation of European Laboratory Animal

104 Science Associations guidelines in humidity- and temperature-controlled environment, with a
105 **daylit** environment for at least 1 week before the experiments. Animals were fed commercial
106 laboratory rat's food pellet and allowed drink tap water ad libitum before the experiments.

Formatted: Font color: Red

107

108 **2.2.2. Study design**

109 Experiments were performed on 58 female Wistar rats weighing 180-200 g. the rats were
110 adapted for 7 days in animal mini clinic and then randomly divided into two groups: control (22
111 animals) and main (36 animals). Animals in main group after randomization received injection
112 with D-gal (100 mg/kg/day, i.p. [31,32]), while in control group received placebo (0.9% saline,
113 0.5 ml/day, i.p.), for 8 weeks. At 21 days after injection with D-gal the 3 cm round tattoo area
114 was prefabricated on each side of rats previously disinfected hip under sterile condition and
115 general anesthesia with pentobarbital (40 mg/kg). All animals in main group were secondly
116 randomized into 3 groups in dependence to treatment (twice in week of intradermal injection
117 under general anesthesia) for 5 weeks: control III group animals treated with microinjection of
118 saline (n=12), main I group receive 0.13 mg of 2% lyophilized powder of Artichoke extracts
119 salivated in water for injection (equivalent of average intradermal dose for patients 10 mg, n=12)
120 and main II – animals receive 1.3 mg 2% lyophilized powder of Artichoke extracts (n=12).
121 After the experiments, all the rats euthanized by pentobarbital (60 mg/kg intraperitoneally).
122 Body weight and skin oedema evaluation was investigated as described below [31].

123

124 **2.3. Determination of activities of enzymatic part of endogenous antioxidant** 125 **defense system of skin of rats**

126 **Isolation** mitochondria incubated with buffer (6 mM succinate, 70 mM sucrose, 220 mM
127 mannitol, 2 mM, **Hepes**, 25 mM KH_2PO_4 , 2.5 mM MgCl_2 , 0.5 mM EDTA, 5 $\mu\text{g/ml}$ catalase, pH
128 7.4) at 37°C and immediately measured of velocity of superoxide anion generation, superoxide
129 dismutase (total), catalase, glutathione peroxidase and malone aldehyde (MDA) were described

Formatted: Font color: Red

Formatted: Font color: Red

130 [33,34]. Rate of H₂O₂ production was determinate as described below [35,36]. The activity of
131 glutathione redox system including determination of glutathione peroxidase (GSH-Px) and
132 glutathione reductase by velocity of redox NADP⁺ formation, and redox glutathione in
133 homogenate of lyophilized in liquid nitrogen skin tissue in according to [34-36]. The protein
134 concentration was determined with BSA protein assay kit.

135

136 **2.4. Statistical analysis**

137 Statistical analysis of presented data as mean ± standard deviation of mean (SD) was performed
138 using the Statistical Sciences (SPSS, version 23.1). The significance level of the differences
139 between the control and main groups assessed using Student t-test and $p < 0.05$ considered as a
140 significant.

141

142 **RESULTS**

143 The studying water artichoke (*C. cardunculus*, cultivated in Georgia, Mtskheta region) extracts,
144 2% content of chlorogenic acid and about 10% of total phenolic acids and confirmed the
145 requirements of the Assessment report on *C. scolymus*, folium EMA/HMPC/150209/2009 for
146 medicinal using artichoke preparation.

147

148 **3.1. Changes in body weight and skin oedema during D-gal-induced skin** 149 **aging and influence of artichoke extracts, 2%**

150 Prolonged 8 weeks D-gal-treated animals characterized by a unique skin appearance, with
151 wrinkling's and furrows, which indicated that developed the evident symptoms of aging. Prior to
152 euthanized, no morbidity/mortality and clinically relief differences in food intake and water
153 consumption in subgroups of main group were not observed. The relative weight of skin
154 markedly decrease in D-gal model of aging. Artichoke at the doses of 0.13 and 1,3 mg/kg

155 improved body weight of D-gal-induced aging rats (table). While the administration of artichoke
 156 extracts in normal rats for 8 weeks did not change, the body weight compared to the control
 157 group. Thus, treatment with artichoke extracts, 2% restores the water **dysbalanced** in the aging
 158 skin in both **dosage**.

Formatted: Font color: Red

Formatted: Font color: Red

159

160 Table. **Therapeutic efficacy of different doses of artichoke extracts for maintenance the**
 161 **activity of endogenous enzymatic antioxidant defense system D-galactose induced aging**
 162 **skin in experimental animals.**

Groups	Control I, n=10	Control II, n=12	D-galactose agin sking rats, n=36		
			Control III, n=12	Liophylized extract artichoke, dose, mg/kg intradermally	
				0.13, n=12	1.3, n=12
Body weight, g	187±22	312±23	245±25 ^{*##}	278±24 ^{**}	268±21 [*]
Relative weight, mg dry/100 mg wet weight	31.5±2.1	32.8±1.4	23.5± 2.3 ^{**##}	29.2± 1.8 ^x	29.7± 2.1 ^x
Velocity of O ₂ ⁻ generation	0.27±0.02	0.31±0.03	0.48± 0.06 ^{###}	0.35± 0.04 ^{*x}	0.36± 0.05 ^{*x}
H ₂ O ₂ , μmol/L · min	1.59±0.14	1.80±0.14	5.15± 0.23 ^{#####}	3.02± 0.32 ^{#####xxx}	3.17± 0.21 ^{#####xxx}
SOD, protein/min	0.33± 0.04	0.32± 0.03	0.26± 0.02 ^{*#}	0.39± 0.03 ^{#xxx}	0.32± 0.03 ^{#x§}
Catalase, H ₂ O ₂ /mg protein/min	64±9	67±8	42±4 ^{*##}	68±6	59±6 [#]
Glutathione redox	3.18±	2.90±	1.83±	2.41±	2.23±

potential, GSH/GSSG	0.38	0.29	0.23 ^{***##}	0.19 ^{**xx}	0.15 ^{***x}
Glutathione peroxidase, nMol NADP/mg protein	2.44± 0.22	2.69± 0.33	1.73± 0.23 ^{***##}	2.51± 0.20 ^x	1.97± 0.13 [#]
Glutathione reductase, µMol NADPH/g wet tissue	0.10± 0.02	0.19± 0.03 [*]	0.29± 0.04 ^{***##}	0.18± 0.04 ^{*x}	0.11± 0.03 ^{#xxx}
MDA, µmol/mg protein	0.88± 0.08	0.92± 0.10	1.48± 0.16 ^{***##}	0.96± 0.06 ^{xxx}	1.09± 0.09 ^{xx}

163 Note: * - compared with control 1, # - with control 2 group, x - with control 3 and § - between artichoke
 164 extracts treatment groups; significance of difference of comparison: one symbol – p<0.05, two – p<0.01,
 165 three - p<0.001, absence of symbol indicated that differences is not significance (p>0.05).
 166

167 3.2. D-gal-induced aging changes in skin and activity of total SOD and 168 generation of superoxide anion.

169 D-gal in dose 100 mg/kg i.p. during 8 weeks cause to significant decreased in total SOD activity
 170 in skin in comparison with control I and control II, while differences in SOD activity between
 171 control I and control II groups did not mentioned (table). At the same time, the velocity of
 172 superoxide anion generation increased by 15% in control II group when comparing the rate of
 173 O₂⁻ production in 240 days rats (table). Treatment with 2% artichoke extract from the 21 days
 174 after D-galactose induced aging in rats leads to increase SOD activity by 50% and by 23% in
 175 comparison with control III groups and this accompanied with markedly decreasing in velocity
 176 of O₂⁻ generation by 27% and 25% in low and high doses of extracts, respectively. The velocity
 177 of superoxide anion generation at the end of the treatment in both dosage of artichoke extracts
 178 did not differences from the level in placebo (control II) group.

179

Formatted: Font color: Red

Formatted: Font color: Red

180

181 3.3. D-gal-induced aging changes in skin and activity of catalase and 182 generation of hydrogen peroxide

183 There were no significant differences in catalase activity between control groups. Exposure to D-
184 galactose did not induced changes in catalase activity in skin tissue (table). However, the
185 production of H₂O₂ increased under treatment of D-gal and exceeds control II level by 186%.
186 Treatment with 2% artichoke leaf extract increased the level of catalase activity, and decrease the
187 level of H₂O₂ production by 42% in dosage of 0.13 mg and by 25% under higher doses.

188

189 3.4. D-galactose-induced aging changes in skin and activity of glutathione 190 redox system

191 Exposure to D-gal reduced the GSH content in skin tissue from 1.20±0.13 nmol/mg/protein to
192 0.74±0.13 nmol/mg/protein (p< 0.01 vs. control III). Treatment with artichoke extract at doses
193 0.13 and 1.3 mg/kg significantly recovered the GSH content up to 0.98±0.09 and 0.89±0.09
194 nmol/mg/protein (p<0.01 and p<0.05, respectively) when compared to D-gal-treated animals.
195 Simultaneously the GSH/GSSG ratio is proportionately decreased in D-gal-induced skin aging
196 model by 37%. Treatment with artichoke extracts in doses of 1.3 mg/kg restored the glutathione
197 redox and it has reached level in the same aging groups while at higher doses treatment the
198 GSH/GSSG ratio increased only by 22% (table). Due to D-gal-treatment observed significantly
199 decreasing of GSH-Px activity, without any differences in GR activity (table). Treatment with
200 artichoke extracts in dose of 0.13 mg increased the level of GSH-Px by 31% and only by 14%
201 (NS) at doses of 13 mg/kg. Ratio of activities of SOD/(Catalase + GSH-Px), which represents
202 equilibrium between formation of hydrogen peroxide from superoxide dismutation and its
203 utilization by catalase and GSH-Px equal 5.0±0.3x10⁻³ in rats at the beginning of the
204 experiments and 4.6±0.2 x10⁻³ in control II group. In D-gal model of aging skin ratio

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font color: Red

205 SOD/(Catalase + GSH-Px) increased to $6.0 \pm 0.2 \times 10^{-3}$, and decreased to 5.5 ± 0.2 and 5.2 ± 0.2
206 after artichoke extracts treatments in low and high dosage, respectively. Simultaneously, the
207 redox potential, ratio of generation $O_2^{\cdot-}/H_2O_2$ which equal in intact group 0.17 ± 0.04 decrease to
208 0.09 ± 0.01 in D-gal treated control III group and increase to 0.12 ± 0.2 ($p < 0.01$) after artichoke
209 treatment. There were no correlation between the level of ratio SOD/(Catalase + GSH-Px) and
210 MDA content in skin ($r=0.37$, NS).

211

212 3.5. D-galactose-induced aging changes in skin MDA content

213 Despite that level of MDA also determinate as a marker of lipid peroxidation in skin and other
214 tissues, MDA content, as a final product of lipid peroxidation, could not reflects the
215 disturbances in the sensitivity of lipid to oxidation [37]. In the model of D-gal-induced aging
216 levels of MDA in skin significant elevated, when compared to the control group ($p < 0.001$)
217 following 42 days of exposure to D-gal, but not in aging group without D-gal-treatment (table).
218 Interestingly, treatment of rats with artichoke at doses of 0.13 and 1.3 mg/kg significantly
219 decreased the levels of MDA in skin in both cases.

220

221 4. DISCUSSION

222 D-gal is pharmacological adaptive aging model, because D-gal primary roles in pathogenesis of
223 aging. Skin aging is a complicated multitargets **dysbalancing** progression in the epidermis and
224 dermis which documented by rising in superoxide anion production in D-gal-induced skin aging
225 model in rats. Influence of artichoke extracts restored skin relative weight and leads to an
226 increase of solubility in neutral salt, acid, and decreased pepsin solubility collagen fraction,
227 restored the hexosamine/collagen (hydroxyproline) ratio and decreased the activity of nuclear
228 transcription factor (NF- κ B). Local prolonged treatment with artichoke extracts improved
229 collagen metabolism and attenuated the progression of inflammation in D-gal-induced skin aging
230 model [29]. Early it was shown, that chronic (6-8weeks) administration of D-gal blocking of

Formatted: Font color: Red

glycometabolism (hyperproduction of advanced glycation products), **dysbalanced** and loses of antioxidant activity of tissue (decreasing the level of SOD and glutathione peroxidase activity) and increased level of MDA in dose dependent manner (50-500 mg/kg i.p. or subcutaneously) [10, 32, 38-40]. Rats in the model group exhibited the typical changes of aging skin compared with the control group, rats in the model group had significantly increased MDA content, and decreased serum SOD and GSH-Px activities ($P < 0.05$). The end product of free radicals oxidizing of unsaturated lipids of biological membranes is MDA which can influence exchange of substances between cells, and finally lead to rupture and death of cells. Extract of artichoke is rich in phenolic and flavonoids and gives a powerful antioxidant activity [14-16, 40]. Pre-clinical and clinical investigations have suggested that the artichoke leaf extract has potential lipid-lowering and hepatoprotective effects [16-19, 21,22, 24,25]. The beneficial effects of artichoke could mainly **attributed** to its antioxidant components: the main substances are mono- and dicaffeoylquinic acid (cynarin and chlorogenic acid), caffeic acid (1%) and volatile sesquiterpene and flavonoids (1%) that include the glycosides luteolin-7-beta-rutinoside (scolymoside), luteolin-7-beta-D-glucoside and luteolin-4-beta-D-glucoside [14-16, 39]. Several *in vitro* studies have shown that the antioxidant potential of artichoke extracts is dependent on radical scavenging and metal ion chelating effect of its constituents such as cynarin, chlorogenic acid and flavonoids. However, pure constituents of artichoke extracts shown to produce less inhibitory activity on free radical production than the extract itself [14,15]. Interestingly, that artichoke is favors that synthesis of coenzymes **NAD₊(NADH₂)** and **NADP⁺(NADPH₂)** and mainly of the NADP(NADPH₂) pair, which take key plays in the regulation of antioxidant/prooxidant status of the cell and its including in the antioxidant properties of artichoke extracts could be included. Preincubation of HUVEC cells or human leukocytes with the artichoke extract at concentrations of 25–100 µg/mL for 24 h abolished ROS generation induced by lipopolysaccharide and oxidation of low density lipoproteins [20, 40]. Early it was shown that artichoke (*C. scolymus*) in dosage 20, 40 80 mg/kg daily per os in D-gal (40 mg/kg

Formatted: Font color: Red

Formatted: Font color: Red

Formatted: Font color: Red

257 body weight) daily for 36 days increase activity of SOD in brain and liver, GSH-Px in brain, and
258 catalase activity in liver [32]. In present article for the first time was study influence of local
259 intradermal action of *C. scolymus* extracts on restoration the ability of endogenous antioxidant
260 defense system to prevent free radical injury development in skin of D-gal-treated (100 mg/kg
261 daily for 8 weeks, i.p.) rats. D-gal (100 mg/kg daily for 8 weeks, i.p.) skin aging in rats
262 characterized increasing in superoxide anion generation in and **hydrogene** peroxide in widely
263 applied to anti-aging pharmacology studies sub-acutely aging models of rodents induced by
264 chronic injection of D-gal [39]. States of skin in this model accompanied with decrease in the
265 activity of SOD, catalase and GSH-Px, and increased production of superoxide anion and
266 hydroperoxide. Hyperproduction of hydrogen peroxide in aging occurs in response disturbances
267 in aerobic respiration and one molecule of catalase can inactivate about 6 million hydrogen
268 peroxide molecule per min by combined them two a time. Thus, the less increased in catalase
269 activity under treatment of artichoke really could sufficient to neutralized produced hydrogen
270 peroxide under decreasing of superoxide anion generation and as a result its oxidation to H₂O₂
271 by SOD. Oxidative damage was concomitant to an imbalance in the principal antioxidant
272 cytoplasmic agent - a significant reduction in cellular GSH, which exerts antioxidant activity by
273 acting as a free-radical scavenger during the reductive detoxification of hydrogen peroxide and
274 lipid peroxide is one of the important target of skin-whitening effect of aging. Exposure to D-gal
275 reduced the GSH content in skin tissue, while artichoke extract at doses 0.13 and 1.3 mg/kg
276 significantly recovered the GSH content. Due to D-gal-treatment observed significantly
277 decreasing of GSH-Px activity, **withought** any differences in GR activity (table). Treatment with
278 artichoke extracts in dose of 0.13 mg markedly increased the level of GSH-Px by 45% and 13%
279 in dose of 13 mg/kg. The data suggest that oxidative stress reduces glutathione redox potential
280 and that prevention disturbances in GSH redox cycle activity appears to be an important
281 component of the antiaging phenomenon.

282

Formatted: Font color: Red

Formatted: Font color: Red

283 **5. CONCLUSION**

284 In conclusion the redox potential of the $O_2/2H_2O$ redox system could play a key role in the “Free
285 Radical Theory of Aging” , seems to address a key facet of intrinsic biological instability of
286 living systems throughout unavoidably formed ROS in the course of metabolism and arising due
287 to the action of various exogenous factors, damage biomolecules [1-5, 37-39]. Obtained data
288 indicate that the concomitant use of 2% artichoke extract improve reserve ability of antioxidant
289 defense system and exert antiaging action in this model of skin aging in experimental animals.
290 The increased reserve ability of intrinsic antioxidant defense system of skin after course of local
291 treatment with artichoke extracts emphasizes artichoke dry extract efficacy in cosmetic
292 formulation and its beneficial effects for anti-aging skin care.

293

294 **CONSENT**

295 Is not applicable

296

297 **ETHICAL APPROVAL**

298 Authors declared that the all procedures with animals meet the requirements of Declaration of
299 Helsinki, Finland in its seven revisions (General Assembly, October, 2013) [Declaration of
300 Helsinki History Website". Ethical Principles For Medical Research. The JAMA Network.
301 Retrieved 26 July 2015] and European Directive 2010/63/EU of the European Parliament and of
302 the Council on the protection of animals used for scientific purposes

303

304 **ACKNOWLEDGMENTS**

305 Authors are grateful to the Chief animal's clinic Kamkamudze R. and his staff for animal
306 housing and Associate Professor Bobkova Natalia V. of the Pharmaceutical Natural Sciences
307 Department Institute of Pharmacy of Sechenov First Moscow State Medical University

308 (Sechenov University) for authentication of artichoke leaves used in this study and Tsivtsivadze
 309 E, PhD as a director of “Biotechpharm GE” in part of artichoke extracts preparations.

310

311

312 **COMPETING OF INTEREST**

313 The authors declare that they have no conflict of interests regarding the publication of this paper.

314 The authors alone are responsible for the content and writing of this article.

315

316 **REFERENCES**

- 317 1. Ganceviecene R, Liakou AI, Theodoridis A, Mafrantonaki E, Zouboulis CC. Skin anti-aging
 318 strategies. *Dermato-Endocrinology*. 2012; 3:308-319. <http://doi.org/10.4161/derm.22804>.
- 319 2. Gilchrest BA, Campisi J, Chang H, Fisher G, Kulesz-Martin M. Skin aging: molecular mechanisms
 320 and tissue consequences. *J Invest Dermatol*. 2015; 135(4):950-953. doi.: 10.1038/jid.2014.546.
- 321 3. Gragnani A, MacCornic SM, Chominski V, Ribeiro de Noronha SM, Correa de Naronha SAA,
 322 Ferreira LM. Review of major theories of skin aging. *Advances in Aging Res*. 2014; 3:265-284.
 323 <http://dx.doi.org/10.4236/aar.2014.34036>.
- 324 4. [eSilva SA, Leonardi GR, Michniak-Kohn B. An overview about oxidation in clinical practice of skin](#)
 325 [aging. *An Bras Dermatol*. 2017; 92\(3\):367-374. <http://dx.doi.org/10/1590/abd1806-4841.20175481>.](#)
- 326 5. Polissak B, Dahmane RG, Godic A. Intrinsic skin aging: The role of oxidative stress. *Acta*
 327 *Dermatovenerologica* . 2012; 21:33-36. doi: 10.2478/v10162-012-0009-0
- 328 6. Sevindik M (2018). Investigation of Antioxidant/Oxidant Status and Antimicrobial Activities of
 329 *Lentinus tigrinus*. *Advances in pharmacological sciences*, 2018. Volume 2018, Article ID 1718025, 4
 330 pages <https://doi.org/10.1155/2018/1718025>
- 331 7. [Sevindik M, Akgul H, Pehlivan M, Selamoglu Z \(2017\). Determination of therapeutic potential of](#)
 332 [Mentha longifolia ssp. longifolia. *Fresen Environ Bull*. 26: 4757-4763.](#)
- 333 8. Rinnerthaler M, Bishof J, Streubel MK, Trost A, Richter K (2015). Oxidative stress in aging human
 334 skin. *Biomolecules*. 5: 545-589. <https://doi.org/10.3390/biom5020545>.

Formatted: Font color: Red

Formatted: Font color: Red

- 335 9. Chang L, Liu X, Li H, Liu J, Guo Z, Xiao K et al. D-galactose induces a mitochondrial complex I
336 deficiency in mouse muscle: potential benefits of nutrient combination in ameliorating muscle
337 impairment. *J Mol Food*. 2014; 1-8. Doi:10.1989/jmf.2013.2830
- 338 10. Chen P, Chen F, Zhou B. Antioxidative, anti-inflammatory and anti-apoptotic effects of ellagic acid in
339 liver and brain of rats treated by D-galactose *Scientific Reports*. 2018; 8:1465-
340 1474.doi:10.1038/s41598-018-19732-0
- 341 11. Ho SC, Liu JH, Wu RY. Establishment of the mimetic aging effect in mice caused by D-galactose.
342 *Biogerontology*. 2003; 4(1):15-18
- 343 12. Liu C, Hu J, Mao Z, Kang H, Liu H, Fu W, Lv Y, Zhou F. Acute kidney injury and
344 inflammatory response of sepsis following cecal ligation and puncture in d-galactose-induced
345 aging rats. *Clin Interv Aging*. 2017; 12: 593-602.
- 346 13. Liu S, Chen Z, Cai X, Sun Y, Zhao C, Liu F, Liu D. Effects of dimethylaminoethanol and
347 compound amino acid on D-galactose induced skin aging model of rat. *Scientific World J*.
348 2014; article ID 507351, 7 pages. <http://dx.doi.org/10.1155/2014/507351>.
- 349 14. Shwe T, Prachayasakul W, Chattipakorn N, Chattipakorn SC. Role of D-galactose induced brain
350 aging and its potential used for therapeutic intervention. *J Ex. Gerontology*. 2018; 101:13-36.
351 <https://doi.org/10.1016/j.exger.2017.10.029>
- 352 15. Georgievs SS, Boyadzhieva SS, Angelov G. Intensification of extraction of bioactive substances from
353 artichoke wastes. *Bulgarian Chemical Communication*. 2016; 48 (special issue E): 451-455.
- 354 16. Ben Salem M, Affes H, Ksouda K, Dhouibi R, Sahnoun Z, et al. Pharmacological Studies of
355 Artichoke Leaf Extract and Their Health Benefits. *Plant Foods Hum Nutr*. 2015; 70(4):441-53. doi:
356 10.1007/s11130-015-0503-8.
- 357 17. Joy JF, Haber SL. Clinical uses of artichoke leaf extract. *Am J Health Syst Pharm*. 2007; 64: 904-909.
358 <https://doi.org/10.2146/ajhp070013>
- 359 18. El-Deen SAS. Artichoke, the ancient hepatic friend and promising multifunctional
360 therapeutic agent. *J Pharmaciol Ther Res*. 2018; 2(1):4-5.
361 <http://www.alliedacademies.org/journal-pharmacology-therapeutic-research/>

- 362 19. Rondanelli M, Opizzi A, Faliva M, Sala P, Perna S, et al. Metabolic management in
363 overweight subjects with naive impaired fasting glycaemia by means of a highly standardized
364 extract from *Cynara scolymus*: a double-blind, placebo-controlled, randomized clinical trial.
365 *Phytother Res.* 2014;28(1):33-41. doi: 10.1002/ptr.4950.
- 366 20. Juzyszyn Z., Czerny B., Pawlik A. Drożdżik M. The effect of artichoke (*Cynara scolymus*
367 L.) extract on ROS generation in HUVEC cells. *Phytotherapy Res.* 2008; 22(9): 1159-1161
368 <https://doi.org/10.1002/ptr.2385>.
- 369 21. Gebhardt R. Antioxidative and protective properties of extracts from artichoke (*Cynara*
370 *scolymus* L.) against hydroperoxide-induced oxidative stress in cultured rat hepatocytes.
371 *Toxicol. Appl Pharmacol.* 1997; 144(2): 279-286. <https://doi.org/10.1006/taap.1997.8130>
- 372 22. Brown JE, Rice-Evans CA. Luteolin rich artichoke extract protects low-density lipoprotein from
373 oxidation *in vitro*. *Free Radical Res.* 1998; 29:247–55.
- 374 23. Perez-Garcia F, Adzet T, Canigueral S. Activity of artichoke leaf extract on reactive oxygen
375 species in human leukocytes. *Free Radical Res.* 2000; 33:661–665.
376 <https://doi.org/10.1080/10715760000301171>.
- 377 24. Küskü-Kiraz Z, Mehmetçik G, Dogru-Abbasoglu S, Uysal M. Artichoke leaf extract reduces
378 oxidative stress and lipoprotein dyshomeostasis in rats fed on high cholesterol diet.
379 *Phytotherapy Res.* 2010; 24 (4): 565–570. <https://doi.org/10.1002/ptr.2985>.
- 380 25. Ibrahim GE, El-Raey MA, Eldahshan OA, Souleman AMA. Effect of extraction on phenolic
381 content, symyarin and antiradical activities of artichoke leaves and roots. *Eur Scientific J.*
382 2013; 9(6):100-110.
- 383 26. Zapolska-Downar D, Zapolska-Dowanar A, Naruszewicz M, Siennicka A, Krasnodebska B,
384 Koldxiej B. Protective properties of artichoke (*Cynara scolymus*) against oxidative stress
385 induced in cultured endothelial cells and monocytes. *Life Sci.* 2002; 71:2897-2908.
386 [https://doi.org/10.1016/S0024-3205\(02\)02136-7](https://doi.org/10.1016/S0024-3205(02)02136-7).

- 387 27. Heidarian E, Rafieian-Kopaei M. Protective effect of artichoke (*Cynara scolymus*) leaf
388 extract against lead toxicity in rat. *Pharm Biol.* 2013; 51(9): 1104-9.
389 <https://doi.org/10.3109/13880209.2013.7779312013>
- 390 28. El Morsy E M, Kamel R. Protective effect of artichoke leaf extract against paracetamol-
391 induced hepatotoxicity in rats, *Pharmaceutical Biology.* 2015; 53(2): 167-173. doi:
392 10.3109/13880209.2014.913066
- 393 29. Fantini N, Colombo G, Giori A, Riva A, Morazzoni P, et al. Evidence of glycemia-lowering
394 effect by a *Cynarascolymus L.* extract in normal and obese rats. *Phytotherapy Res.* 2011;
395 25:463-466. <https://doi.org/10.1002/ptr.3285>
- 396 30. Safaa M, Hanaa A, Abdel F, Nahila A, Abdelaaty S. *Cynara scolymus* for relieving on
397 nonalcoholic steatohepatitis induced in rat. *Int. J. Pharmacy&Pharm. Sciences.* 2013; 5(1):
398 57-66. <http://dx.doi.org/10.1155/2016/4030476>.
- 399 31. Sukoyan G. Tsivtsivadze E, Golovach V, Kezeli T, Demina N. Anti-aging effect of *Cynara*
400 *Cardunculus L. var. Cynara Scolymus L.* extract in D-galactose-induced skin aging model in
401 rats. *Pharmacology&Pharmacy.* 2018; 9: 428-439. <https://doi.org/10.4236/pp.2018.910032>.
- 402 32. Song S, Zhao L, Wang W, Cong T, He H (2012) Using aging rats model to investigate anti-
403 oxidative ability of artichoke (*Cynara Scolymus L.*) leaf extract.
404 <https://doi.org/10.17660/ActaHortic.2012.944.15>
- 405 33. Galenko-Yaroshevskii VP, Agadzanova AV, Lapina IV, Gaironovskaya TV, Popkova LV, et
406 al. Effectiveness of combined treatment of superoxide dismutase and reamberin in skin
407 ischemia. *Bull Exp Biol Med.* 2006; 142(6):707-709. [https://doi.org/10.1007/s10517-006-](https://doi.org/10.1007/s10517-006-0457-9)
408 0457-9
- 409 34. Galenko-Yaroshevskii VP, Bagmetova EN, Fil'chukova IA, Sidel'nikov A, Lapina IV, et al.
410 Antihypoxic and antinecrotic effect of mexidol in skin ischemia. *Bull Exp Biol Med.* 2005;
411 139(2): 702-704. <https://doi.org/10.1007/s10517-005-0248-8>.

- 412 35. Karsanov NV, Sukoyan GV, Antelava NA, Varazanashvili NA. The effect of energostim on
413 myocardial damage during coronary artery occlusion. *Exper. Clin Pharmacol.* 2003; 66(5),
414 17-21. doi: <https://doi.org/10.30906/0869-2092-2003-66-5-17-21>
- 415 36. Sukoyan GV, Mumladze MR, Oboladze ED, Varazanashvili NA. *In vitro* effects of
416 gentamicin, ampicillin, and cefobid on energy supply and antioxidant protection systems of
417 venous blood erythrocytes in newborns. *Bull Exp Biol Med.* 2005; 139(6): 671-674. doi:
418 10.1007/s10517-005-0374-3.
- 419 37. Dzhanashia P, Vladitskaya OV, Kobalava MA, Sukoyan GV, Sherosia Ts, Andriadze NA.
420 Lipid hyperperoxidation in the impaired structural organization of thrombocytic membranes
421 in patients with chronic heart failure (communication 1). *Russian Medical Journal.* 2005;4:
422 35-50.
- 423 38. Lisanti MP, Martinez-Outschoorn UE, Lin Z, Pavlides S, Whitaker-Menezes D, et al.
424 Hydrogen peroxide fuels aging, inflammation, cancer metabolism and metastasis. *Cell Cycle.*
425 2011; 10(15):2440-2449. <https://doi.org/10.4161/cc.10.15.16870>
- 426 39. Mukherjee PK, Maity N, Nema NK, Sarkar BK. Bioactive compounds from natural resources
427 against skin aging. *Phytomedicine.* 2011; 19:64-73.
428 <https://doi.org/10.1016/j.phymed.2011.10.003>.
- 429 40. Piston M, Machado I, Branco CS, Cesio V, Heinzen H et al. Infusion, decoction and
430 hydroalcoholic extracts of leaves from artichoke (*Cynara cardunculus* L. subsp. *Cardunculus*)
431 are effective scavengers of physiologically relevant ROS and RNS. *Food Res. Intern.* 2014;
432 64:150-156. <http://dx.doi.org/10.1016/j.foodres.2014.05.078>
- 433
- 434