

COMPONENTS OF TREE BIOMASS IN AN INTEGRATED CROP-LIVESTOCK-FOREST SYSTEM

ABSTRACT

**Aims:** This study performed the adjustment of volumetric models, and determined the biomass of *Eucalyptus grandis* x *Eucalyptus urophylla* hybrid cultivated in a crop-livestock-forest integration system (CLF).

**Study design:** The experimental area consists of a crop-livestock-forest integration system where trees are hybrids clones of seven year old *Eucalyptus grandis* x *Eucalyptus urophylla*.

**Place and duration of study:** This work was carried out at Fazenda Santa Brígida, Ipameri, Goiás (Brazil).

**Methodology:** A forest inventory of the area was carried out in October 2015 when the tree component was fully developed. Diameter at breast height (DBH) (at 1.30 m) and total height (H) of trees were measured in the field and categorized according to 4 classes. Afterwards, 12 trees were felled, which were cubed and compartmentalized to determine the volume and biomass of their components. The volumetric models developed by Schumacher & Hall and Ogaya were applied to obtain determination coefficients.

**Results:** The average DBH was 18.28 cm and the average H was 23.47 m. The highest volumes of wood were observed in the diametric classes that presented the largest number of individuals, however in the class of higher DBH an average individual volume of 0.36 m<sup>3</sup> of wood was observed. The total biomass of *Eucalyptus* was 56.64 Mg ha<sup>-1</sup>, being 83.70% wood, 6.52% in branches, 6.37% in bark and 3.40% in leaves.

**Conclusion:** The volumetric models developed by Schumacher and Hall as well as Ogaya were found to be applicable for estimating the volume of wood in CLF systems, where both showed a determination coefficient of 0.866.

**Keywords:** compartmentalization; eucalyptus; CLFS; volume, modeling

Formatted: Font: 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

## 29 1. INTRODUCTION

30 Crop-livestock-forest (CLF) integration has been proposed as an economically  
31 viable production technology for the recovery and renovation of degraded areas in the  
32 Cerrados, a vast tropical savanna ecoregion of Brazil [1]. The main habitat types of the  
33 Cerrado include: forest savanna, wooded savanna, park savanna, and gramineous-woody  
34 savanna. Savanna wetlands and gallery forests are also included.

35 In addition to the formation or recovery of pastures, this technique CLF favors  
36 the production of grain cultivars along with the exploitation of tree biomass production, either  
37 simultaneously, sequentially, or rotationally [2]. The intensification of the production has  
38 several benefits to the producer and the environment, such as: improving the physical,  
39 chemical, and biological conditions of the soil, increasing the cycling and efficiency for the  
40 use of nutrients, reducing production costs of agriculture and livestock, opening new areas  
41 for production, as well as diversifying and stabilizing the income of the producer [3]. The  
42 included tree component biomass promotes benefits ranging from soil protection to  
43 availability of nutrients and organic matter in the soil by the deposition of leaves and tree  
44 branches [4,5].

45 *Eucalyptus* has been presented as a good option in the integrated CLF due to  
46 its rustic nature, rapid growth, great utilization, and economic value in the market, being an  
47 alternative for farmers interested in wood production [6]. The rapid growth of the eucalyptus  
48 in Brazil can be explained by the large investment of the companies in order to achieve the  
49 demand of the silvicultural products. The high productivity of stands planted by the Brazilian  
50 forest companies is recognized worldwide, due to the higher average productivity the  
51 minimum time required until harvesting, continuous investment in research on genetic  
52 improvement and silvicultural. The preservation and maintenance of native forests to  
53 preserve the biodiversity, makes planted forests even more important for preservation,  
54 because they have high productivity in a short period of time, avoid exploration of natural  
55 forests. These plantations, besides providing different products, also help with carbon

Formatted: Font: 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

56 sequestration, while also maintaining animal biodiversity [7,8].

Formatted: Font: (Default) Arial, 10 pt

57 In the integration of CLF, one of the challenges lies in the careful planning of  
58 the system in defining short, medium, and long-term actions. The competition for light  
59 between forest species and agricultural and pastoral crops requires special attention, as this  
60 directly influences the productivity of the system. However, this competition can be reduced  
61 by selecting genetic material, adapting the planting arrangement of the tree component, and  
62 silvicultural treatments, which, in addition to adding value to the wood, also allows for greater  
63 light entry into the integration system that contributes to the maintenance or increase in the  
64 productivity of the other components, as pasture and agricultural culture [9,2].

Formatted: Font: (Default) Arial, 10 pt

65 The configuration of tree component arrangements may influence plant height,  
66 diameter of breast height (DBH), and volume of wood. Clemente [10] verified that integrated  
67 systems with single and double row arrangements provided higher volumes of wood. In their  
68 study, Oliveira et al. [11] verified higher volumes of *Eucalyptus* wood in integrated systems  
69 with forages, than in monoculture.

Formatted: Font: (Default) Arial, 10 pt

70 The balanced relationship between the integrated CLF components is important  
71 for the expression of the productive potential of the species involved. In the case of tree  
72 species, especially the fast-growing ones such as *Eucalyptus*, accumulation and biomass  
73 production are influenced by age of trees, among other factors. In the juvenile phase,  
74 accumulation is higher in the canopy components, whereas a greater increase of biomass in  
75 the trunk component is perceived over time [12].

Formatted: Font: (Default) Arial, 10 pt

76 However, this work had the aim of adjusting volumetric models and determining  
77 the biomass of the *Eucalyptus grandis* x *Eucalyptus urophylla* hybrid cultivated in an  
78 integrated crop-livestock-forest (CLF) system in Ipameri / Goiás (Brazil).

Formatted: Font: (Default) Arial, 10 pt

## 80 2. MATERIALS AND METHODS

Formatted: Font: 10 pt

This study was carried out at Fazenda Santa Brígida in the municipality of  
Ipameri – Goiás (Brazil), located at 17° 39'22" south latitude and longitude west of 48°

Formatted: Font: (Default) Arial, 10 pt

12°22", and at an altitude of 800 m a.s.l. [11]. According to the classification of Köppen-Geiger [13], the climate of the region is Aw (tropical savannah with dry season in winter), and the average temperature of the region varies between 22 and 23°C. The mean annual rainfall is between 1200 and 1400 mm, having a wet period of seven months from October to April, and the remaining five months characterize the dry season.

Formatted: Font: (Default) Arial, 10 pt

According to Embrapa [14], the soil of the experimental area is classified as red latosol, being naturally acidic and with low base saturation with good drainage and sand-clay texture.

Formatted: Font: (Default) Arial, 10 pt

The experimental area consists of an integrated crop-livestock-forest system that tends towards the east-west direction. Tree planting was carried out with clones of the hybrid *Eucalyptus grandis* x *Eucalyptus urophylla* (GG 100) in 2008 in an area of approximately four hectares (ha). The trees were arranged in double rows (1 m x 1 m x 26 m), occupying 1.4 ha of the total area of the system [11].

Formatted: Font: (Default) Arial, 10 pt

81 Before establishment of the seedlings, soil acidity was corrected with the use of  
82 two tons per hectare of dolomitic limestone and one ton of gypsum. At planting, the base  
83 fertilization used was 400 kg ha<sup>-1</sup> of yoorin thermophosphate and 180 grams (g ha<sup>-1</sup>) of NPK  
84 formulation 06-30-30, supplemented with 0.4 g ha<sup>-1</sup> of zinc, 0.2 g ha<sup>-1</sup> copper, and 0.2 g ha<sup>-1</sup>  
85 boron. The half of this composition was incorporated into the bottom of the planting pit, and  
86 the remainder was distributed 20 days after planting in two lateral holes located 10 cm from  
87 the seedlings.

Formatted: Font: (Default) Arial, 10 pt

88 **15 Fifteen** months after **of** planting, **a** pruning was performed, and at 30  
89 months, the third cover fertilization was applied using NPK formulation 00 - 00 - 36 with the  
90 addition of 0.2 g ha<sup>-1</sup> of copper and 0.6 g ha<sup>-1</sup> of boron per plant, provided in continuous fillet  
91 in the crown projection.

Formatted: Font: (Default) Arial, 10 pt

## 92 93 **2.1 Determination of *Eucalyptus* biomass**

Formatted: Font: 10 pt

94 The forest inventory of the area was carried out in October 2015 when the tree

Formatted: Font: (Default) Arial, 10 pt

component was fully developed, seven years after planting. DBH (diameter at breast height (DBH) at 1.30 m above soil level) and H (total height (H) of the trees) were measured in the field with the aid of a caliper and the use of a clinometer. For the of DBH and H measurements, a systematic sampling was carried out with regular intervals on every sixth tree line in which measurements were made on the two individuals that composed it.

Based on the data obtained from the forest inventory, the trees were distributed in four classes of diameters (Table 1). Subsequently, three individuals were felled for sampling in each diameter class, considering the lower, middle, and upper limits, totaling 12 trees.

**Table 1.** Diametric distribution (cm) of *Eucalyptus* in the integrated CLF system.

Class interval	Class center	Number of sampled individuals
9.1 – 14	11.5	16
14.1 – 19	16.5	59
19.1 – 24	21.5	80
24.1 – 29	26.5	6

After the trees were felled, they were subjected to rigorous sampling, according to the method developed by Smalian and described by Finger [15]. The height points for taking diameters with and without bark were: 0.10 m, 0.30 m, 1.30 m, 2.30 m, and so on, at one meter intervals up to full height.

After measuring the diameters, the trunk was sectioned into 1 m long logs to the point where the diameter was seven centimeters (commercial diameter). From there, up to a diameter of three centimeters was considered as tip of the trees, and the remaining portions to the apex were considered branches. For the determination of the dry weight of wood, the methodology developed by Schumacher [16] was used, in which three samples were taken along the trunk. The total height of the tree was divided into three sections, and

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: Italic

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

116 | the midpoint of each third of the tree was taken to compose the sample. Each sampling point  
117 | was composed of the complete disc of the tree cylinder that had a thickness of ten  
118 | centimeters.

Formatted: Font: (Default) Arial, 10 pt

119 |         After sectioning, the logs were weighed both with and without bark to determine  
120 | the wet weight of the wood and bark. The tree canopy, in turn, was divided into two  
121 | components: leaves and branches. These components were also weighed in the field and  
122 | properly sampled to determine the dry weight in the laboratory, as well as to determine wood  
123 | biomass (WB), branch biomass (BB), and leaf biomass (LB).

Formatted: Font: (Default) Arial, 10 pt

124 |         The biomass samples were sent to the Forest Ecology Laboratory (ECOFLOR)  
125 | of the Federal University of Goiás. They were placed in a force air circulation oven at 65°C  
126 | for drying until the weight of the samples remained stable to obtain the dry mass of the  
127 | components with a precision digital scale (0.01 g).

Formatted: Font: (Default) Arial, Superscript

128 |         

Formatted: Font: (Default) Arial, 10 pt

## 129 | **2.2 Data analysis**

Formatted: Font: 10 pt

130 |         In order to relate the DBHs and biomass components of each tree, linear  
131 | regressions were performed for each component: wood, bark, branches, and leaves. For the  
132 | volumetric models, the DBH and the total height of the tree were considered the  
133 | independent variables, and the total volumes and the trunk with the bark were dependent  
134 | variables. Four volumetric models, one single-entry and three double-entry, were chosen  
135 | because they were the most used for the quantification of the production in forest stands and  
136 | have not yet been tested in integrated CLF systems. The models tested are described in  
137 | Table 2.

Formatted: Font: (Default) Arial, 10 pt

138 |

139 |

**Table 2.** Volumetric models tested.

Formatted: Font: (Default) Arial, 10 pt

Author	Type	Model

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Husch	Single entry	$V = \beta_0 + \beta_1 \text{DAP}$
Ogaya	Double entry	$V = \text{DAP}^2 (\beta_0 + \beta_1 H)$
Schumacher & Hall (log)	Double entry	$V = \beta_0 + \beta_1 \text{Ln}(\text{DAP}) + \beta_2 \text{Ln}(H)$
Spurr (log)	Double entry	$V = \beta_0 + \beta_1 \text{Ln}(\text{DAP}^2 H)$

DBH= diameter at breast height; H = total height;  $\beta_0$  = value of the height estimated when the diameter is zero;  $\beta_1$  = slope of the line, which corresponds to the value of the first derivative;  $\beta_2$  = rate of change in volume (m<sup>3</sup>) as height (m) variation occurs with constant DBH (cm);  $\beta_3$  = coefficient of the multivariate model.

The volumetric models were adjusted and evaluated by means of adjustment and precision statistics, following the importance proposed by Draper and Smith [17]: graphical analysis of the residues; estimate of the standard error in percentage (Syx%) that indicates the proximity between the estimated values and those observed and the closer to zero the model and the determination coefficient (R<sup>2</sup>) that shows how much the dependent variables are explained by the independents and, in this case, the closer to a better model.

### 3. RESULTS AND DISCUSSION

#### 3.1 Determination of eucalyptus biomass

The **integrated crop-livestock-forest (CLF)** system evaluated presents a density of 303 trees per hectare. This occupied 33.65% of the area designated to the system and an average production of 0.18 m<sup>3</sup> of wood per tree, totaling a volume of wood without bark of 54.80 m<sup>3</sup> ha in the studied system. The remaining 66.35% were destined to other economic activities within the integration, such as agricultural and forage production.

After performing the forest inventory, the diameter distribution was analyzed, and four diameter classes were obtained. It is notable that the height, density, and volume of wood without bark were higher in class III, being 30.33 m, 151 trees ha<sup>-1</sup>, and 40.06 m<sup>3</sup> ha<sup>-1</sup>,

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

152 respectively (Table 3). Through the dendrometric characteristics of this integrated CLF  
 153 system, it is possible to verify a trend in relation to the height behavior of the plants and their  
 154 DBHs, being that the DBH tends to increase as the height linear increases . This is contrary  
 155 to the expected behavior in more homogeneous forest stands where trees with higher  
 156 heights and smaller diameters are observed. More than the standard can be seen in other  
 157 eucalyptus studies, depending on the clone and the management used in the area, as well  
 158 as Miguel et al [18] observes in homogeneous plantation of *E. urograndis* in Niquelandia,  
 159 GO and Lemos-Junior et al [19] in CFL with same species in Cachoeira Dourada, GO.

160 **Table 3.** Dendrometric characteristics of *Eucalyptus grandis* x *Eucalyptus urophylla* grown in  
 161 the integrated CLF system at Fazenda Santa Brígida in the municipality of Ipameri / Goiás /  
 162 2015.

Diameters Classes (cm)	Mean height (m)	mean DBH (cm)	Density(tree. ha <sup>-1</sup> )	Volume of wood without bark (m <sup>3</sup> ha <sup>-1</sup> )
I (9-14)	12.58	10.45	30	1.25
II (14,1-19)	20.73	14.53	111	9.45
III(19,1-24)	30.33	21.57	151	40.06
IV(24,1-29)	30.22	26.55	11	4.04
Total	23.47	18.28	303	54.80

164  
 165 Generally, resource availability tends to be higher in stands with less density of  
 166 less trees, reflecting higher growth in broader plantations [20]. This fact can be observed in  
 167 this study because in spite of the densification of the trees in the planting lines, the spacing  
 168 between the eucalyptus ridges provides greater light availability in this integrated CLF  
 169 system. This causes the effect observed in the height as demonstrated by Miguel et al [18]  
 170 in homogeneous plantation of *E. urograndis* in Niquelandia, GO, DBH and wood volume as  
 171 demonstrated by Lemos-Junior et al [19] in CFL with same species in Cachoeira Dourada,

- Comment [R1]: This sentence is not cle... [1]
- Formatted ... [2]
- Formatted ... [3]
- Comment [R2]:
- Comment [R3R2]: Why there is change... [4]
- Formatted ... [6]
- Comment [R4]: Provide the error valu... [10]
- Formatted ... [5]
- Formatted ... [7]
- Formatted ... [8]
- Formatted ... [9]
- Formatted ... [11]
- Formatted ... [12]
- Formatted ... [13]
- Formatted ... [14]
- Formatted ... [15]
- Formatted ... [16]
- Formatted ... [17]
- Formatted ... [18]
- Formatted ... [19]
- Formatted ... [20]
- Formatted ... [21]
- Formatted ... [22]
- Formatted ... [23]
- Formatted ... [24]
- Formatted ... [25]
- Formatted ... [26]
- Formatted ... [27]
- Formatted ... [28]
- Formatted ... [29]
- Formatted ... [30]
- Formatted ... [31]
- Formatted ... [32]
- Formatted ... [33]
- Formatted ... [34]
- Formatted ... [35]
- Formatted ... [36]
- Formatted ... [37]
- Formatted ... [38]
- Formatted ... [39]



172 GO, that can be attributed more to the lesser effect of resource competition than to  
173 continuous plantings where the height and DBH ratio are inversely related. Different spacing  
174 and thinning regimes late in the life of the stand presented the highest values of basal area  
175 production. The choice of the best thinning regime for *Eucalyptus* clonal material will vary  
176 according to the plantation objective [21].

Formatted: Font: Italic

177 The maximum and minimum diameter found in this integrated CLF ranged from  
178 9.4 to 28.25 cm, and the highest tree density were located in classes II and III, which  
179 consequently contributed with a higher volume of wood within the CLF system (Figure 1). In  
180 its study with *Eucalyptus* clones GG100 (*E. grandis* x *E. urophylla*) of 4.5 years, Cerdeira  
181 [22] observed a diametric variation between 5.0 and 17.1 cm. Cerdeira [22] also reported  
182 that the classes of greater diameter were those that presented the greater number of  
183 individuals, a DBH variation close to that of the present study, but the central classes were  
184 those with the highest number of individuals. Thus, we highlight that CLF present higher  
185 production of trees with higher diameter class, being relevant for the production of wood with  
186 noble purpose and greater value added by planted individuals.

Formatted: Font: Italic

Formatted: Font: (Default) Arial, 10 pt

187 Although Class IV presents the highest average wood volume per tree of 0.36  
188 m<sup>3</sup>, its contribution to the system is around 10%, among the individuals of lower number  
189 present in this class. However, classes III and II were the ones that concentrated the largest  
190 number of individuals, being responsible for 86.4% of the wood produced in this area of the  
191 integrated CLF with an average volume of 0.26 and 0.08 m<sup>3</sup>, respectively, per individual  
192 within the classes.

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt




Figure 1. Volume of wood ( $\text{m}^3 \text{ha}^{-1}$ ) in different diametric classes of *Eucalyptus* trees in an integrated crop-livestock-forest system in Ipameri / Goiás.

When evaluating three 32-month-old eucalyptus trees (*E. urophylla* x *E. grandis*), Torres et al [6] found volumes of wood without bark ranging from 0.01 to 0.24  $\text{m}^3$  for DBHs between 6.79 and 20.8 cm. These results are similar to those verified in the present work in the corresponding diametric classes.

The total biomass produced by the hybrid GG 100 in this integrated CLF was 56642.76  $\text{kg ha}^{-1}$ , and the trees belonging to class III contributed the most in this production. In general, 38303.00  $\text{kg ha}^{-1}$  of biomass was quantified through individuals with DBH between 19.1 and 24 cm 24.1 and 29 cm, contributing 67.62% of the total produced (Table 2.4).

Among the components analyzed in the eucalyptus, the wood biomass contributed the most to the total biomass of the integrated CLF trees with 83.70%, followed by the branches with 6.52%. Considering the trunk biomass (wood + bark), this was 90.07% and the contribution of the canopy (leaves + branches) was 9.92% (Table.4).

Formatted: Font: Italic

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Comment [R5]: Be uniform in using this as a common name or a generic name.

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224

**Table 4.** Eucalyptus wood biomass (WB), bark biomass (KB), branch biomass (BB), and leaf biomass (LF) with seven years of integrated CLF cultivation in Ipameri / Goiás.

Class	Class I (9-14 cm)	Class II (14.1-19 cm)	Class III (19.1-24 cm)	Class IV (24.1-29 cm)	Total of components
LB (kg ha <sup>-1</sup> )	169.80	615.93	992.19	149.23	1927.15 (3.40)*
KB (kg ha <sup>-1</sup> )	393.60	1008.29	1974.81	318.41	3695.11 (6.52)
BB (kg ha <sup>-1</sup> )	139.31	859.96	2363.32	245.73	3608.32 (6.37)
WB (kg ha <sup>-1</sup> )	1540.81	9918.34	32972.68	2980.35	47412.18 (83.70)
Total Biomass	2243.52 (3.96)	12402.52 (21.90)	38303.00 (67.62)	3693.72 (6.52)	56642.76


\* Values in parentheses refer to the percentage of component contribution in relation to total biomass.

Evaluating biomass components in eucalyptus stands with different ages, [16] verified a trunk biomass around 80.3% for plantations with 8 years, a result that is consistent with the present study.

In the present work, it was verified that the order of contribution of biomasses in the different components was Wood>Branches>Bark>Leaves (Figure 2). These results were similar to those **verified reported** by Giumarães et al. [23] in Allegrete//RS in homogeneous plantation of *E. dunnii* with four years of age and those reported by Benatti [24] in Campos das Vertentes/MG using eucalyptus clones I-144 with 6.5 years of age. **Regarding biomass, it should be pointed out that the 8-year-old crops present about 80% of the biomass in the tree trunks shows the potential of planting, at this age, already for biomass for energy production, since most of the biomass of the planting may be removed and used as fuel.**

Comment [R6]: Change in font features????

Formatted	... [40]
Formatted	... [41]
Formatted	... [42]
Formatted	... [44]
Formatted	... [46]
Formatted	... [48]
Formatted	... [43]
Formatted	... [49]
Formatted	... [50]
Formatted	... [45]
Formatted	... [47]
Formatted	... [51]
Formatted	... [52]
Formatted	... [53]
Formatted	... [54]
Formatted	... [55]
Formatted	... [56]
Formatted	... [57]
Formatted	... [58]
Formatted	... [59]
Formatted	... [60]
Formatted	... [61]
Formatted	... [62]
Formatted	... [63]
Formatted	... [64]
Formatted	... [65]
Formatted	... [66]
Formatted	... [67]
Formatted	... [68]
Formatted	... [69]
Formatted	... [70]
Formatted	... [71]
Formatted	... [72]
Formatted	... [73]
Formatted	... [74]
Formatted	... [75]
Formatted	... [76]
Formatted	... [78]
Formatted	... [82]
Formatted	... [77]
Formatted	... [79]
Formatted	... [80]
Formatted	... [81]
Formatted	... [83]
Formatted	... [84]



225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240

Figure 2. Contribution of eucalyptus biomass from different components and diametric classes in the integrated CLF system in the Ipameri / Goiás municipality.

Formatted: Font: (Default) Arial, 10 pt

Different results to the present work were observed by [Utima \[25\]](#) when working with *E. benthamii* in an CLF at 12 months of age, [Caron et al. \[26\]](#) with *Eucalyptus* sp. of three years of age and [Calil et al. \[27\]](#) in eucalyptus in the agroforestry system of 18 months, where they verified the inversion in the production of leaves and bark. This explains the effect of biomass distribution during the different tree development phases, as the first one focused on leaf expansion and the second on the development of trunks and leaf area limitation [\[16, 28, 16\]](#).

Formatted: Font: (Default) Arial, 10 pt

In his study comparing different commercial clones of eucalyptus in integrated CLF systems in the municipality of Juara/MT, [9] verified that GG100 eucalyptus was the one that allocated the largest biomass in the trunk when planted in double lines corresponding to 62.6% of the total biomass of the trees at 15 months of age, Moreover, this behavior was maintained at seven years, as verified in the present work with the same clone in which this

241 | percentage reached 90.07% as predicted by [29].

Formatted: Font: (Default) Arial, 10 pt


242 |           From a commercial and structural point of view, the objective of the cultivator is  
243 | to increase the volume of the trunk and to improve the quality of the wood. Less biomass in  
244 | the branches is desirable since the primary product is the wood for commercialized [9].

Formatted: Font: (Default) Arial, 10 pt

245 |           In Figure 3, it can be verified through the regression analysis that the linear  
246 | model was adequate to explain the increase of the biomass of the different components in  
247 | relation to the DBHs. One can observe an intense relation between these, mainly for wood  
248 | and bark, and with lower intensity with the branches.

Formatted: Font: (Default) Arial, 10 pt

UNDER PEER REVIEW



249

250

251

252

Figure 3. Production of the different components of biomass of *Eucalyptus grandis* x *Eucalyptus urophylla* in relation to DBH in an integrated CLF system in the city of Ipameri / Goiás.

Comment [R7]: You have to indicate the significance level for the  $R^2$  values.

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

253 The biomass gains of wood and bark due to the increase of the DBH were  
 254 homogeneous. In other words, as the DAP increased, the biomass of these components  
 255 also increased, which is justified by the high values of the coefficient of determination  
 256 presented in the respective regressions ( $R^2$  0.9229 and  $R^2$  0.8604). While the biomass of  
 257 leaves and branches did not present significant increases with the increase of DBH ( $R^2$   
 258 0.6647 and  $R^2$  0.4672), which evidences the accumulation of biomass as a function of age.  
 259 In more developed plantations, the biomass of the leaves and branches decrease [16].

### 3.2 Adjustments of volumetric models


260 Table 5 shows the adjustments for the different models tested as a function of  
 261 height and DBH. Considering the graphical analysis of the residues, the standard error and  
 262 the determination coefficient, the double entry models of Schumacher & Hall (log) (0.866 and  
 263 21.33%) and Ogaya (0.866 and 20.78%) can be considered the most efficient to predict the  
 264 volume of wood for an integrated CLF system in this spatial arrangement (Figure 4).

266 **Table 5.** Adjustments of volumetric models attributed to the eucalyptus plantation used in  
 267 the integrated crop-livestock-forest system and their estimated coefficients ( $\beta$ ),  
 268 coefficient of determination ( $R^2$ ), and standard error (Syx %).

Models	$\beta$ 0	$\beta$ 1	$\beta$ 2	$R^2$	Syx(%)
Husch	-0.18775	0.031469	-	0.681	35.10
Ogaya	-0.27662	0.00045	0.019777	0.886	20.78
Schumacher & Hall (log)	-8.8478	0.617035	1.848882	0.886	21.33
Spurr (log)	-1.45173	0.207884	-	0.748	31.21

- Comment [R8]: It is not necessary to r [... [87]
- Formatted [... [85]
- Formatted [... [86]
- Formatted [... [88]
- Comment [R9]: See the earlier comment.
- Formatted [... [89]
- Formatted [... [90]
- Formatted [... [91]
- Formatted [... [92]
- Formatted [... [93]
- Formatted [... [94]
- Formatted [... [95]
- Formatted [... [96]
- Formatted [... [97]
- Formatted [... [98]
- Formatted [... [99]
- Formatted [... [100]
- Formatted [... [101]
- Formatted [... [102]
- Formatted [... [103]
- Formatted [... [104]
- Formatted [... [105]
- Formatted [... [106]
- Formatted [... [107]
- Formatted [... [108]
- Formatted [... [109]
- Formatted [... [110]
- Formatted [... [111]
- Formatted [... [112]
- Formatted [... [113]
- Formatted [... [114]
- Formatted [... [115]
- Formatted [... [116]
- Formatted [... [117]
- Formatted [... [118]
- Formatted [... [119]
- Formatted [... [120]
- Formatted [... [121]
- Formatted [... [122]
- Formatted [... [123]
- Formatted [... [124]

269  
 270



271

272

Figure 4. Waste residue distribution of the volumetric models tested for clones of *Eucalyptus*

273

*grandis* x *Eucalyptus urophylla* in integrated CLF system in the city of Ipameri / Goiás.

274

Formatted: Font: (Default) Arial, 10 pt



275 |           However, all models tested had a satisfactory distribution of residues, an  $R^2$   
276 | ranging from 0.681 to 0.866 and a standard error of less than 35%, which makes it possible  
277 | to use these models to estimate the volume of wood in the integrated CLF system.

Formatted: Superscript

Formatted: Font: (Default) Arial, 10 pt

278 |           Lemes Junior et al. [19] considered the Näslund and Ogaya models as the most  
279 | efficient to determine the volume of wood in the Integrated CLF system with eucalyptus of  
280 | six years of age in Cachoeira Dourada / Goiás. These presented coefficients of  
281 | determination of 99.5 and 99.1%, respectively. However, in spite of verifying a higher  
282 | coefficient of determination for the Schumacher & Hall model, Miguel [30] observed a  
283 | standard error that was considered high, another criterion used to indicate the volumetric  
284 | model was the graphical distribution of the residues. In this scenario, the Takata model was  
285 | the most suitable for estimating the volume of wood in a settlement of seven years of *E.*  
286 | *urophylla* in Niquelândia, north of Goiás.

Formatted: Font: (Default) Arial, 10 pt

          In their study with a silvipastoral system in the region of Coronel  
Pacheco/MG, Müller et al. [31] tested different volumetric models to estimate the volume of  
eucalyptus trees, and they found that the Schumacher & Hall model presented the best fit for  
those conditions, as also verified in the present work. This demonstrated that the  
Schumacher & Hall model has also been used for the integrated CLF system, since its  
statistical properties almost always result in non-biased estimates.

Formatted: Font: (Default) Arial, 10 pt

287 |           With regards to the tree component of the integrated CLF system evaluated, a  
288 | forest inventory was carried out to verify that at seven years after planting, the total biomass  
289 | produced by the *Eucalyptus grandis* x *Eucalyptus urophylla* hybrid presented 56.64 Mg ha<sup>-1</sup>,  
290 | a mean tree height of 25 m, and a chest height of 18. 222 cm. This biomass presented a  
291 | distribution with greater quantity in the wood component, followed by the branches, bark,  
292 | and leaves.

Formatted: Font: (Default) Arial, 10 pt

293 |           Although **crop-livestock-forest integration** CLF presents limitations in its  
294 | operation, this system becomes feasible from an adequate planning that meets the  
295 | production demands of the property in the short, medium, and long term. Although it is a

296 | complex system because of the need to optimize the production conditions of each  
297 | component, it is necessary to know the ecophysiology of the plants that will make up the  
298 | integration. Besides the aggregate environmental benefits, this is important to determine if  
299 | the productivity of the system is satisfactory to meet the social and economic demands and,  
300 | thus, achieve the precepts of sustainability.

Formatted: Font: (Default) Arial, 10 pt

301 |         The environmental and productive importance of the integrated CLF system can  
302 | be considered for the need to deepen the knowledge of the behavior of each component of  
303 | the integration and prompted the interest in carrying out this research. It can be concluded  
304 | that, finally, the initial objectives were reached, and it is, therefore, time for these results to  
305 | be released.

Formatted: Font: (Default) Arial, 10 pt

306 |         Another aspect to be considered is the need for continuation of this research,  
307 | both for this region of the Cerrado of Goiás and for the other regions of the country. It is  
308 | known that many agricultural systems, conducted in an inadequate way, have contributed to  
309 | the degradation of environmental quality and, due to this condition, seek to maintain  
310 | production through the opening of new arable areas. Knowledge of crop-livestock-forest  
311 | integration, as well as studies on the various possibilities of system implementation, are  
312 | important factors for the productivity of agroecosystem and reduction of negative impacts on  
313 | the Cerrado and other biomes.

Formatted: Font: (Default) Arial, 10 pt

#### 315 | **4. CONCLUSIONS**

Formatted: Font: 10 pt

316 |         The highest average volume of wood per tree was verified in the highest DBH  
317 | class; The volumetric models of Schumacher & Hall and Ogaya were efficient to estimate the  
318 | volume of wood in the integrated CLF system; The biomass of *Eucalyptus grandis* x  
319 | *Eucalyptus urophylla* was 56.64 Mg ha<sup>-1</sup>, and 90.07% was present in the components of the  
320 | trunk, while the others allocated in the canopy. Adequate cultural (debris and thinning)  
321 | treatment throughout the crop cycle has negatively influenced the development of culture.

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

322

323 | **REFERENCES**

324

325 | 1. VASCONCELOS VV, VASCONCELOS CV, VASCONCELOS DM. Phyto-Environmental  
326 | Characterization of Brazilian Savanna (Cerrado) and Brazilian Atlantic Forest, with the  
327 | Research of Stone Lines and Paleosols Geografia. Ensino & Pesquisa (UFSM), 2010; 14: 6-  
328 | 17. Portuguese.

329

330 | 2. VIANA MCM, ALVARENGA RC, MASCARENHAS MHT, MACEDO GAR, SILVA EA, SILVA  
331 | KT, RIBEIRO PCO. Consorciação de Culturas com o Eucalipto no Sistema de Integração  
332 | Lavoura-Pecuária Floresta. In: XXIX CONGRESSO NACIONAL DE MILHO E SORGO, 29,  
333 | Águas de Lindóia. SP: Associação Brasileira de Milho e Sorgo (ABMS), 2012. Portuguese.

334

335 | 3. GONTIJO NETO MM, VIANA MCM, ALVARENGA RC, SANTOS EA, SIMÃO EP,  
336 | CAMPANHA MM. Sistemas de integração Lavoura Pecuária Floresta em Minas Gerais.  
337 | Boletim de Indústria Animal, 2014; 71: 183-191. Portuguese

338

339 | 4. XAVIER DF, LÉDO FJS, PACIULLO DSC, PIRES MFÁ, BODDEY RM. Dinâmica da  
340 | serapilheira em pastagens de braquiária em sistema silvipastoril e monocultura. Pesquisa  
341 | Agropecuária Brasileira, 2011; 46: 1214-1219. Portuguese.

342

343 | 5. GATTO A, BARROS NF, NOVAIS RF, SILVA FR, LEITE FP, VILLANI EMA. Estoques de  
344 | carbono no solo e na biomassa em plantações de eucalipto. Revista Brasileira de Ciências  
345 | do Solo, 2010; 34: 1069-1079. Portuguese.

346

347 | 6. TORRES CMME, OLIVEIRA AC, PEREIRA BLC, JACOVINE LAG, OLIVEIRA NETO SN,  
348 | CARNEIRO ACO. Estimativas da produção e propriedades da madeira de eucalipto em  
349 | Sistemas Agroflorestais. Scientia Forestalis, 2016; 44: 137-148. Portuguese.

Formatted: Font: 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

7. IBA. Relatório IBÁ. Brasília. Indústria Brasileira de Árvores 2016. 6-15 . Portuguese.

8. ORTIGOZA FDSS, SENNA RJ. Caracterização do segmento de florestas plantadas de eucalipto em Mato Grosso do Sul: Revista de Política Agrícola. Brasília, 2016; 74-89, 2016. Portuguese.

9. TONINI H, MORALES MM, MENEGUCI JLP, ANTONIO DBA, WRUCK FJ. Biomassa e área foliar de clones de eucalipto para a desrama. Nativa, 2016; 4: 271-276. Portuguese.

10. CLEMENTE MA. Características agronômicas do sorgo e eucalipto em diferentes arranjos espaciais. 51 f. Dissertation (Master Degree) – Federal University of Uberlândia, 2015. Portuguese.

11. OLIVEIRA P, FREITAS RJ, KLUTHCOUSKI J, RIBEIRO AA, CORDEIRO LAM, TEIXEIRA LP, MELO RAC, SILVA A, VILELA L, BALBINO LC. Evolução de sistemas de integração lavoura-pecuária-floresta (ILPF): estudo de caso da fazenda Santa Brígida, Ipameri, Go. 2 ed. Planaltina: Embrapa Cerrados, 2015. 50 p. Portuguese.

12. SAIDELLES FLF, KÖNIG FG, SCHUMACHER MV. Avaliação da biomassa e dos nutrientes em espécies florestais de rápido crescimento. In: 1 Simpósio Brasileiro de Pós-Graduação em Engenharia Florestal. Anais...Santa Maria: Perspectivas e Tendência da Pesquisa Florestal, 2001; 1: 134-144. Portuguese.

13. CARDOSO MRD, MARCUZZO FFN, BARROS JR. Classificação Climática de Köppen-Geiger para o Estado de Goiás e o Distrito Federal. Acta Geográfica. 2014; 8: 44-55. Portuguese.

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

Formatted: Font: (Default) Arial, 10 pt

377

378 | 14. EMBRAPA. Sistema brasileiro de classificação de solos. 3. ed. Rio de Janeiro-RJ:  
379 | Embrapa Solos, 2013. 306p. Portuguese.

Formatted: Font: (Default) Arial, 10 pt

380

381 | 15. FINGER CAG. Fundamentos de biometria florestal. Santa Maria: UFSM/ CEPEF/  
382 | FATEC, 1992. 269 p. Portuguese.

Formatted: Font: (Default) Arial, 10 pt

383

384 | 16. SCHUMACHER MV, WITSCHORECK R, CALIL FN. Biomassa em povoamentos de  
385 | *Eucalyptus spp.* de pequenas propriedades rurais em Vera Cruz, RS. Ciência Florestal  
386 | (UFSM. Impresso), 2011; 21: 17-22. Portuguese.

Formatted: Font: (Default) Arial, 10 pt

387

388 | 17. DRAPER NR, SMITH H. Applied regression analysis. New York: Jonh Willey & Sons,  
389 | 407 p. 1966.

Formatted: Font: (Default) Arial, 10 pt

390

391 | 18. MIGUEL EP, LEAL FA, ONO HA, LEAL, UAS. Modelagem na predição do volume  
392 | individual em plantio de *Eucalyptus urograndis*. Revista Brasileira de Biometria, 2014; 32(4):  
393 | 584-598. Portuguese

394

395 | 19. LEMES JUNIOR JM, SILVA NETO CM, SOUZA KR, GUIMARÃES LE, OLIVEIRA FD,  
396 | GONÇALVES RA, MONTEIRO MM, LIMA NL, VENTUROLI F, CALIL FN. Volumetric models  
397 | for *Eucalyptus grandis* x *urophylla* in a crop-livestock-forest integration (CLFI) system in the  
398 | Brazilian cerrado. African Journal of Agricultural Research, 2016; 11(15): 1336-1343.

Formatted: Font: Italic

Formatted: Font: Italic

399

400 | 20. REINER DA, SILVEIRA ER, SZABO MS. O uso do eucalipto em diferentes  
401 | espaçamentos como alternativa de renda e suprimento da pequena propriedade na região  
402 | sudoeste do Paraná. Synergismus sScyentifica. 2011; 6 (1). Portuguese.

Formatted: Font: (Default) Arial, 10 pt

403

404 21. FERRAZ FILHO AC, MOLA-YUDEGO B, GONZALEZ-OLABARRIA JR, SCOLFORO  
405 JRS. (2018). Thinning regimes and initial spacing for Eucalyptus plantations in Brazil. Anais  
406 da Academia Brasileira de Ciências, 2018; 90(1): 255-265. Portuguese.

407  
408 22. CERDEIRA ALN. Modelos para quantificação do volume de diferentes sortimentos em  
409 plantio de *Eucalyptus urophylla* x *Eucalyptus grandis*. 66 f. Monography (Forest  
410 Engineering), University of Brasília, 2012. Portuguese.

Formatted: Font: (Default) Arial, 10 pt

411  
412 23. GUIMARAES CC, SCHUMACHER MV, WITSHORECK R, SOUZA HP, SANTOS JC.  
413 Biomassa e nutriente em povoamento de *Eucalyptus dunni* Maiden no pampa gaúcho.  
414 Revista Árvore. 2015; 39 (5): 873-882. Portuguese.

Formatted: Font: (Default) Arial, 10 pt

415  
416 24. BENATTI BP. Compartimentalização de biomassa e nutrientes em estruturas de plantas  
417 de eucalipto cultivadas em solos distintos. 114 f. Dissertation (Master Degree) – Federal  
418 University of Lavras. 2013; Portuguese.

Formatted: Font: (Default) Arial, 10 pt

419  
420 25. UTIMA, A. Y. Crescimento e rendimento dos componentes agrícola e arbóreo de um  
421 sistema integrado de produção agropecuária no ano de implantação em área de proteção  
422 ambiental. 78 f. Dissertation (Master Degree in Agronomy), Federal University of Paraná,  
423 2015. Portuguese.

Formatted: Font: (Default) Arial, 10 pt

424  
425 26. CARON BO, ELOY E, SOUZA VQ, SCHMIAT D, BALBINOT R, BEHILNG A,  
426 MONTEIRO GC. Quantificação da Biomassa florestal em plantios de curta rotação com  
427 diferentes espaçamentos. Comunicata Scientiae. 2015; 6 (1): 106-112. Portuguese.

Formatted: Font: (Default) Arial, 10 pt

428  
429 27. CALIL FN, VIERA M, SCHUMACHER MV, LOPES VG, WITSCHORECK R. Biomassa e  
430 nutrientes em Sistema agrossilvicultural no extremo sul do Brasil. Ecologia e Nutrição

431 Florestal. 2013; 1 (2): 80-88. Portuguese. ▲

Formatted: Font: (Default) Arial, 10 pt

432

433 28. GONÇALVES JLM, STAPE JL, BENEDETTI V, FESSEL VAG, GAVA JL. Reflexos do  
434 cultivo mínimo e intensivo do solo em sua fertilidade e na nutrição das árvores. In:  
435 GONÇALVES, J. L. M.; BENEDETTI, V. (eds.) Nutrição e fertilização florestal. Piracicaba:  
436 IPEF, 2005. 1-57. Portuguese. ▲

Formatted: Font: (Default) Arial, 10 pt

437

438 29. MUÑOZ F, RUBILAR R, ESPINOSA M, CANCINO J, HERRERA M. The effect of pruning  
439 and thinning on above ground aerial biomass off *Eucalyptus nitens* (Deane & Maiden)  
440 Maiden. Forest Ecology and Management. 2008; 255 (3): 365-373. ▲

Formatted: Font: (Default) Arial, 10 pt

441

442 30. MIGUEL, E. P. Avaliação biométrica e prognose da produção de *Eucalyptus urophylla*  
443 (s.t. blake) na região norte do estado de Goiás. 166 f. Dissertation (Mester Degree in Forest  
444 Engeneering), Federal University of Paraná. 2009. Portuguese. ▲

Formatted: Font: (Default) Arial, 10 pt

445

446 31. MÜLLER MD, FERNANDES EM, CASTRO CRT, PACIULLO DSC, ALVES FF. Estimativa  
447 de acúmulo de biomassa e carbono em sistema agrossilvipastoril na zona da mata mineira.  
448 Pesquisa Florestal Brasileira, 2009; 60: 11-17. Colombian. ▲

Formatted: Font: (Default) Arial, 10 pt

449

450

451

452

453

454

455

456 ▲

Formatted: Font: (Default) Arial, 10 pt

<b>Page 8: [1] Comment [R1]</b>	<b>Reviewer</b>	<b>09-Apr-19 9:59:00 PM</b>
This sentence is not clear. Rewrite.		
<b>Page 8: [2] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
Font: (Default) Arial, 10 pt		
<b>Page 8: [3] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
Font: (Default) Arial, 10 pt		
<b>Page 8: [4] Comment [R3R2]</b>	<b>Reviewer</b>	<b>09-Apr-19 10:00:00 PM</b>
Why there is change in font style and size??		
<b>Page 8: [5] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
Font: (Default) Arial, 10 pt		
<b>Page 8: [6] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
Font: (Default) Arial, 10 pt		
<b>Page 8: [7] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
Font: (Default) Arial, 10 pt		
<b>Page 8: [8] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
Font: (Default) Arial, 10 pt		
<b>Page 8: [9] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
Font: (Default) Arial, 10 pt		
<b>Page 8: [10] Comment [R4]</b>	<b>Reviewer</b>	<b>09-Apr-19 10:02:00 PM</b>
Provide the error values for these means.		
<b>Page 8: [11] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
Font: (Default) Arial, 10 pt		
<b>Page 8: [12] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
Font: (Default) Arial, 10 pt		
<b>Page 8: [13] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
Font: (Default) Arial, 10 pt		
<b>Page 8: [14] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
Font: (Default) Arial, 10 pt		
<b>Page 8: [15] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
Font: (Default) Arial, 10 pt		
<b>Page 8: [16] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>



Font: (Default) Arial, 10 pt

<b>Page 8: [17] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
-------------------------------	------------	-----------------------------

Font: (Default) Arial, 10 pt

<b>Page 8: [18] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
-------------------------------	------------	-----------------------------

Font: (Default) Arial, 10 pt

<b>Page 8: [19] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
-------------------------------	------------	-----------------------------

Font: (Default) Arial, 10 pt

<b>Page 8: [20] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
-------------------------------	------------	-----------------------------

Font: (Default) Arial, 10 pt

<b>Page 8: [21] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
-------------------------------	------------	-----------------------------

Font: (Default) Arial, 10 pt

<b>Page 8: [22] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
-------------------------------	------------	-----------------------------

Font: (Default) Arial, 10 pt

<b>Page 8: [23] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
-------------------------------	------------	-----------------------------

Font: (Default) Arial, 10 pt

<b>Page 8: [24] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
-------------------------------	------------	-----------------------------

Font: (Default) Arial, 10 pt

<b>Page 8: [25] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
-------------------------------	------------	-----------------------------

Font: (Default) Arial, 10 pt

<b>Page 8: [26] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
-------------------------------	------------	-----------------------------

Font: (Default) Arial, 10 pt

<b>Page 8: [27] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
-------------------------------	------------	-----------------------------

Font: (Default) Arial, 10 pt

<b>Page 8: [28] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
-------------------------------	------------	-----------------------------

Font: (Default) Arial, 10 pt

<b>Page 8: [29] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
-------------------------------	------------	-----------------------------

Font: (Default) Arial, 10 pt

<b>Page 8: [30] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
-------------------------------	------------	-----------------------------

Font: (Default) Arial, 10 pt

<b>Page 8: [31] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
-------------------------------	------------	-----------------------------

Font: (Default) Arial, 10 pt

**Page 8: [32] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 8: [33] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 8: [34] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 8: [35] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 8: [36] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 8: [37] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 8: [38] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 8: [39] Formatted** sdi 22-May-19 3:24:00 PM

Font: Italic

**Page 11: [40] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [41] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [42] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [43] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [44] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [45] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [46] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [47] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [48] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [49] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [50] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [51] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [52] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [53] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [54] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [55] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [56] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [57] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [58] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [59] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [60] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [61] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [62] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [63] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [64] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [65] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [66] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [67] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [68] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [69] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [70] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [71] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [72] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [73] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [74] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [75] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [76] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [77] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [78] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [79] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [80] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [81] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [82] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [83] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 11: [84] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [85] Formatted** sdi 22-May-19 3:24:00 PM

Superscript

**Page 15: [86] Formatted** sdi 22-May-19 3:24:00 PM

Superscript

**Page 15: [87] Comment [R8]** Reviewer 09-Apr-19 10:14:00 PM

It is not necessary to repeat the values that are already in the figures.

**Page 15: [88] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [89] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [90] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [91] Formatted** sdi 22-May-19 3:24:00 PM

Font: 10 pt

**Page 15: [92] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [93] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [94] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [95] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [96] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [97] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [98] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [99] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [100] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [101] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [102] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [103] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [104] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [105] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [106] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [107] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [108] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [109] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [110] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [111] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [112] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [113] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [114] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [115] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [116] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [117] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [118] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [119] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [120] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

**Page 15: [121] Formatted** sdi 22-May-19 3:24:00 PM

Font: (Default) Arial, 10 pt

<b>Page 15: [122] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
---------------------------------	------------	-----------------------------

Font: (Default) Arial, 10 pt

<b>Page 15: [123] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
---------------------------------	------------	-----------------------------

Font: (Default) Arial, 10 pt

<b>Page 15: [124] Formatted</b>	<b>sdi</b>	<b>22-May-19 3:24:00 PM</b>
---------------------------------	------------	-----------------------------

Font: (Default) Arial, 10 pt