

# Evaluation of untraditional Macaroni Formulated by Using Different Grain Meals and Their Mixtures

## ABSTRACT

**Aims:** wheat, Barley and millet meals are having superior nutritional qualities and health benefits; they can be used for supplementation of macaroni. Its effect on physiochemical, rheological, color parameters, cooking quality, nutritional value and sensory evaluation.

**Place and Duration of Study:** Regional Center for Food and Feed, Agriculture Research Center, Giza, Egypt.

**Methodology:** Macaroni was prepared using wheat, barley, millet and composite meals mix (1), mix (2) and mix (3). Proximate chemical composition, rheological, color parameters, cooking quality and sensory evaluation were measured of wheat, barley, millet and composite meals macaroni.

**Results:** Results show that the level of millet replacement led to increasing the fat, ash and total fiber in the products. B-glucan content in barely represented the superiority (3.90 %) as compared with other samples. Substitution of wheat, barley and millet meals (mixed) macaroni have significantly increasing in the water absorption while they have significantly decreasing the cooking time. Highest value of water absorption 54.60 % was found for wheat and lowest value 35.0 % was obtained for millet. Color characteristics indicate that an increasing proportion of millet had signed negative effect on lightness and overall acceptability. While barley addition showed significant positive effect on lightness and overall acceptability. Sensory evaluation scores indicated non significant difference among the control and barley experimental products for overall acceptance, then mixed (1) and millet were the lowest value of overall acceptance.

**Conclusion:** It can be concluded that the possibility of producing macaroni relatively higher in fiber and  $\beta$ -glucan without considerable bad effects on its cooking quality and sensory evaluation and has many benefits for health of diabetes, high cholesterol and heart diseases patients.

**Keywords:** *Wheat, barley, millet meal and mixtures; properties of macaroni.*

## 1. INTRODUCTION

At present, dietary guidelines recommend an increase in the consumption of whole grain cereal products due to their role in reducing the risk of degenerative chronic diseases. Whole grains contain all parts of the grain viz., the endosperm, germ, bran and rich in nutrients and photochemical with known health benefits [1]. Other protective compounds in whole grains include phytate, phyto-oestrogens such as lignans, plant stanols and sterols, and vitamins and minerals. Several epidemiological studies have shown that consumption of whole grain cereals is associated with reduced incidence of diabetes [2; 3], cardiovascular diseases and certain cancers [4; 5].

Traditionally, pasta products are made from wheat semolina, although more recently other cereals have been used to partially replace it [6].

Hull-less barley being a cereal grain is suitable for cereal pasta. The nutritional value of whole-grain barley to be low in fat content, higher in total dietary fiber and essential amino acid therefore has a positive health profile. Beta-glucans from barley have been found to reduce blood glucose and insulin levels with hypo-cholesterolemic effects [7]. The Food and Drug Administration (FDA) has indicated that dietary intake of 3 g / day of barley  $\beta$ -glucan helps to decrease total cholesterol in both the serum and the low-density lipoprotein [8].

Finger millet (*Eleusinecoracana*) also known as ragi is one of the important millet consumed without dehulling. It has good source of methionine, cysteine, lysine and high levels of calcium, iron, zinc, lipids then it has high concentrations of threonine and tryptophan along with less leucine than other cereals [9].

Millets have nutraceutical properties in the form of antioxidants which prevent deterioration of human health such as lowering blood pressure, risk of heart disease, prevention of cancer and cardiovascular diseases, diabetes, decreasing tumor cases etc. [10]. Millet is easily available cheap in cost and gluten-free food, which can be a substitute for celiac patients.

Bread and Pasta are the major processed cereal products that are part of the daily diets of the most people in large number of countries and especially the Mediterranean as in Egypt. While these products are low in fat and good sources of complex carbohydrates, they are usually not good sources of dietary and, in particular, soluble fiber [11].

Pasta's versatility, long shelf life in dry form, availability in numerous shapes and sizes, high digestibility, good nutrition, and relatively low cost are attractive to the consumer. It has become more popular due to its nutritional properties and being regarded as a product with low glycemic index [12]. Pasta with a mixture of durum wheat and beta-glucan enriched barley flour (BF) (60 / 40 %, w / w) and found it to have a final content of 5 %  $\beta$ -glucan. Quality parameters, cooking loss and dry matter did not vary substantially from the control, suggest in high potential for consumer acceptance [13]. The addition of millet flours to the pasta will improve the dietary fiber content [14]. Therefore the present study was aimed to evolution the macaroni formulations by wheat, barley, millet meal and their mixed and its effect on physiochemical, rheological, color parameters, cooking quality, nutritional value of macaroni and sensory analysis.

## 2. MATERIAL AND METHODS

### 2.1 Materials

Wheat (*Triticum durum*), barley (*Hordeum distychnum*) and millet (*Pennisetum Spp.*) grains cultivar was obtained from Egypt. Wheat, barley and millet which was obtained from Corp Intensification Research Department - Field Crops Research Institute - Agriculture Research Center during 2018.

#### 2.1.1 Preparation of meal grains

A ten kg of wheat, barley and millet sample used in this investigation was stored at temperature 25 °C and relative humidity less than 62 % according to the methods described in [15]. Wheat, barley and millet sample was cleaned mechanically to remove dirt, dockage, impurities and other strange grains by Carter Dockage Tester according to the methods described in [16]. The extraction rate of flour sample was adjusted to recurred rate (100 % extraction) which had milled by laboratory mill 3100 Perten according to the methods described in [17] for meal flour.

#### 2.1.2 Analysis of Raw Materials

##### 2.1.2.1 Physical properties

Cleanliness, dockage, shrunken and broken, foreign materials, total damaged kernels and total defects were separated and determined manually (hand picking). Test weight pound per bushel, Test weight P/B = (Kg/Hectoliter)  $\div$  1.278 according to [15]. A thousand kernel weights were determined by counting the kernels (wheat, barley and millet) in a 10 g sample [18]. Gluten and falling number were determined to wheat, barley, millet meals and their mixtures according to [19].

#### **2.1.2.2 Determination of color of raw materials and produced macaroni**

Colour was evaluated by a colorimeter CR-400 (Konica Minolta, Japan) in the CIE LAB colour space: Commission International de l'Eclairage (CIE) tristimulus L\* a\* b\* parameters were determined using colour meter (Colour Tec PCMTM Color Tec Associates, Inc., Clinton, NJ, USA), according to the method outlined in [18].

#### **2.1.2.3 Chemical properties**

Moisture, crude protein, ash, crude fiber, fat, mineral, vitamins and aflatoxin were determined to wheat, barley, millet meals and their mixtures according to [19] and [15]. The nitrogen free extract (NFE) was calculated by difference. Beta-glucan was determined according to [20].

#### **2.1.2.4 Rheological properties**

All mixtures of flours were tested by Alveograph, consistograph while amylograph was used to determine the maximum viscosity, temperature at the maximum viscosity and the transition point according to the methods described in [17] in Regional Center for Food and Feed, Agri. Res. Center, Cairo, Egypt). To determine the rheological properties of the different types of meal grains and their mixtures according to the methods described by [17].

### **2.2 Methods:**

#### **2.2.1 Marconi processing**

Marconi was processed into flour, using the method of fresh pasta dough according to the methods described in Regional Center for Food and Feed, Agri. Res. Center, Cairo, Egypt [17]. All macaroni was used in this formula to produce macaroni by six formulas:

1-Wheat 100 %

2-Barley 100 %

3-Millet 100 %

4-Mix1= (12.5 % barley, 12.5 % millet and 75 % wheat)

5-Mix 2= (25.0 % barley, 25.0 % millet and 50 % wheat)

6-Mix 3= (37.5 % barley, 37.5 % millet and 25 % wheat)

#### **2.2.2 Evaluation of cooking quality of produced macaroni**

Cooking quality, increase in volume, cooking loss and optimal cooking time was carried out according to the method outlined in [18].

#### **2.2.3 Sensory evaluation**

The sensory characteristics of macaroni were evaluated according to [21]. Sensory attributes like appearance, flavor, taste, colour, mouth feeling and overall acceptability for all the samples were assessed.

#### **2.2.4 Statistical analysis**

Data of three replicates were determined by Duncan's multiple range test at ( $P>0.05$ ) level was used to compare between means using SAS programs [22].

## **3. RESULTS AND DISCUSSION**

### **3.1 Proximate analysis for wheat, barley, millet meals and their mixtures**

The proximate composition of the samples, including moisture, protein, fat, ash, fiber, nitrogen free extract and total caloric values is shown in Table 1 in the present study. The results revealed that the moisture content were no significant effect for both wheat, mix 1 and mix 2 meals (10.50, 10.50 and 10.20 gm /100 gm, respectively). The average protein content of wheat meal and barley ranged between 13.4 % - 9.8 %, respectively, these agreements with work by [23]. The high fat content of meal was millet and lowest value was recorded in mix 3 (4.2 % and 1.21 %), respectively. And [24] reported 1.58 per cent fat in wheat and 35.5 per cent fat in flaxseed, per cent fat content of oat, maize, pearl millet and mungbean was (4.42, 4.74, 5.47 and 1.85 %), respectively. The ash content of meal ranged between 1.80 % to 1.06 % millet and barley respectively, [25] reported 1.53 % ash content of pearl millet. The ash content indicated a rough estimation of the mineral value of the product. The high fiber content was millet 8.5 % and the lowest was mix 3 meals 1.30 %. Our results are in conformity with [24]. Nitrogen free extracts (NFE) % ranged between 65.8

- 77.55 % for millet and mix 3, these results are lower than results by [26]. The calorific value of samples was ranged between 345 - 363.6 %. Barley had highest calorific value when compared to other treatments. Millets contain 60 - 70 % carbohydrates, 7 - 11 % proteins, 1.5 - 5 % fat, and 2 - 7 % crude fiber [10]. While  $\beta$ -glucan content in barely flour represented the superiority (3.90 mg/g) as compared with its content in millet flour (0.75 mg/g) and wheat flour (0.70 mg). These results are in agreement with [27].

**Table 1: proximate analysis for wheat, barley, millet meals and their mixtures**

Analysis	Wheat	Barley	Millet	Mix 1	Mix 2	Mix 3
Moisture content %	10.5 <sup>a</sup>	7.6 <sup>c</sup>	8.7 <sup>b</sup>	10.5 <sup>a</sup>	10.2 <sup>a</sup>	8.2 <sup>b</sup>
Protein content %	13.4 <sup>a</sup>	9.8 <sup>d</sup>	11.0 <sup>c</sup>	12.1 <sup>b</sup>	11.7 <sup>b</sup>	10.6 <sup>c</sup>
Fat content %	1.43 <sup>c</sup>	1.75 <sup>b</sup>	4.2 <sup>a</sup>	1.27 <sup>d</sup>	1.24 <sup>d</sup>	1.21 <sup>d</sup>
Ash content %	1.45 <sup>ab</sup>	1.06 <sup>c</sup>	1.8 <sup>a</sup>	1.27 <sup>bc</sup>	1.22 <sup>bc</sup>	1.14 <sup>bc</sup>
Fiber content %	1.52 <sup>c</sup>	2.64 <sup>b</sup>	8.5 <sup>a</sup>	1.35 <sup>c</sup>	1.32 <sup>c</sup>	1.30 <sup>c</sup>
Nitrogen free extracts %	71.7 <sup>c</sup>	77.2 <sup>a</sup>	65.8 <sup>d</sup>	73.91 <sup>b</sup>	73.92 <sup>b</sup>	77.55 <sup>a</sup>
Total caloric values %	353.3 <sup>b</sup>	363.6 <sup>a</sup>	345.0 <sup>c</sup>	353.9 <sup>b</sup>	355.2 <sup>b</sup>	363.2 <sup>a</sup>
$\beta$ -glucan	0.70 <sup>e</sup>	3.90 <sup>a</sup>	0.75 <sup>e</sup>	1.12 <sup>d</sup>	1.50 <sup>c</sup>	1.91 <sup>b</sup>

*a,b,...Means with the same letter in the same row are not significantly different at (P>0.05).*

### 3.2 Minerals for wheat, barley, millet meals and their mixtures

Minerals for wheat, barley, millet meals and their mixtures were presented on Table 2. It showed that millet was the low significant effect of calcium (8.0 mg) for all samples. Pearl millet accompanying grains of other types have oxalic acid which by forming a complex, which is insoluble, with calcium results in reduction of bioavailability of this mineral [28]. The concentration of calcium in pearl millet is very less and if oxalate is present then the condition will become worse. Iron value ranged between 3.19 - 2.50 mg wheat and barley respectively, millet is also a good source of other dietary minerals like manganese, phosphorus and iron [10]. The high value of Magnesium (Mg) was 126.0 mg on wheat meal and the lowest value was 79.0 mg barley meal. And the high values manganese and phosphorus was wheat meal (3.99 mg and 288 mg). The highest potassium value was wheat 363.0 mg and the lowest value was millet 195.0 mg. Selenium (Se) value in all samples ranged between 0.003 - 0.071 mg. Wheat meal sample was high in zinc value compared to all samples and low value was millet samples. Minerals are located in the germ; therefore, we may expect that they are not completely lost during the refining process. Total content of minerals is 2.3 mg per 100 g which is more in quantity in comparison too their cereals consumed commonly. It is a rich source of potassium, B-vitamin, phosphorous, copper, magnesium, zinc, iron, manganese [29].

**Table 2: Minerals for wheat, barley, millet meals and their mixtures.**

Minerals mg	Wheat	Barley	Millet	Mix 1	Mix 2	Mix 3
Calcium (Ca)	29.0 <sup>a</sup>	29.0 <sup>a</sup>	8.0 <sup>b</sup>	25.7 <sup>a</sup>	25.2 <sup>a</sup>	24.70 <sup>a</sup>
Iron (Fe)	3.19 <sup>a</sup>	2.50 <sup>a</sup>	3.0 <sup>a</sup>	2.82 <sup>a</sup>	2.77 <sup>a</sup>	2.71 <sup>a</sup>
Magnesium (Mg)	126.0 <sup>a</sup>	79.0 <sup>c</sup>	114.0 <sup>b</sup>	111.5 <sup>b</sup>	109.5 <sup>b</sup>	107.4 <sup>b</sup>
Manganese (Mn)	3.99 <sup>a</sup>	1.32 <sup>b</sup>	1.60 <sup>b</sup>	3.53 <sup>a</sup>	3.47 <sup>a</sup>	3.40 <sup>a</sup>
Phosphorus (P)	288.0 <sup>a</sup>	2.21 <sup>c</sup>	285.0 <sup>a</sup>	255.0 <sup>b</sup>	250.2 <sup>b</sup>	245.5 <sup>b</sup>
Potassium (K)	363.0 <sup>a</sup>	280.0 <sup>e</sup>	195.0 <sup>f</sup>	321.0 <sup>b</sup>	315.4 <sup>c</sup>	309.4 <sup>d</sup>
Selenium (Se)	0.071 <sup>a</sup>	0.040 <sup>a</sup>	0.003 <sup>a</sup>	0.063 <sup>a</sup>	0.062 <sup>a</sup>	0.061 <sup>a</sup>
Zinc (Zn)	2.65 <sup>a</sup>	2.13 <sup>c</sup>	1.70 <sup>d</sup>	2.35 <sup>b</sup>	2.30 <sup>b</sup>	2.26 <sup>bc</sup>

*a,b,...Means with the same letter in the same row are not significantly different at (P>0.05).*

### 3.3 Vitamins for wheat, barley, millet meals and their mixtures

Millet is an excellent source of vitamin B. In Table 3 millet was the high level of vitamin C 2.0 % in all samples. Wheat meal was the high value of vitamin E in all samples 7.00 %. Vitamin K ranged between 1.00 - 2.00 % in all samples. Matured and dried kernels do not have vitamin C but vitamin B is present in sufficient amount in aleurone layer and the germs. Decortications used for removing hull results in reduced levels of niacin, riboflavin and thiamine to an extent of 50 % in flour. In cereals, niacin is present in both bound and free form and is mainly synthesized by using tryptophan [30]. Quantity of niacin is enough even in hulled form of millet.

**Table 3: Vitamins for wheat, barley, millet meals and their mixtures**

Vitamins %	Wheat	Barley	Millet	Mix 1	Mix 2	Mix 3
Thiamine(B1)	33.0 <sup>a</sup>	17.0 <sup>c</sup>	Non <sup>d</sup>	29.2 <sup>ab</sup>	28.7 <sup>b</sup>	28.1 <sup>b</sup>
Riboflavin(B2)	10.0 <sup>d</sup>	10.0 <sup>d</sup>	24.0 <sup>a</sup>	20.3 <sup>c</sup>	20.9 <sup>bc</sup>	21.5 <sup>b</sup>
Niacin (B3)	36.0 <sup>a</sup>	31.0 <sup>bc</sup>	31.0 <sup>bc</sup>	31.8 <sup>b</sup>	31.3 <sup>bc</sup>	30.7 <sup>c</sup>
Pantothenic acid (B5)	19.0 <sup>a</sup>	6.0 <sup>c</sup>	17.0 <sup>b</sup>	16.8 <sup>b</sup>	16.5 <sup>b</sup>	16.2 <sup>b</sup>
Pyridoxine(B6)	23.0 <sup>c</sup>	20.0 <sup>d</sup>	29.0 <sup>a</sup>	20.4 <sup>d</sup>	24.7 <sup>b</sup>	25.2 <sup>b</sup>
Folic Acid (B9)	10.0 <sup>d</sup>	6.0 <sup>e</sup>	21.0 <sup>a</sup>	11.2 <sup>c</sup>	17.9 <sup>b</sup>	18.2 <sup>b</sup>
Vitamin C	Non <sup>c</sup>	Non <sup>c</sup>	2.0 <sup>a</sup>	0.5 <sup>c</sup>	1.1 <sup>b</sup>	1.7 <sup>a</sup>
Vitamin E	7.0 <sup>a</sup>	Non <sup>e</sup>	Non <sup>e</sup>	5.0 <sup>b</sup>	3.4 <sup>c</sup>	1.7 <sup>d</sup>
Vitamin K	2.0 <sup>a</sup>	2.0 <sup>a</sup>	1.0 <sup>b</sup>	1.85 <sup>a</sup>	1.93 <sup>a</sup>	1.96 <sup>a</sup>

a,b,...Means with the same letter in the same row are not significantly different at (P>0.05).

### 3.4 Mycotoxins content for wheat, barley and millet grains

Results in Table 4 show that Mycotoxin content in wheat, barley and millet grains. It can be noticed that the sample had lowest aflatoxin content before storing under detection limit (0.5 ppb) for aflatoxin, ochratoxin, zearalenone, fumonisin. More over it can be concluded that the sample wheat, barley and millet were under detection limit (0.5 ppb) of the standard Egyptian maximum (B1=10 ppb and total aflatoxin =20 ppb). Aflatoxin content was well within the safe limit 50 ml/kg recommended by [31].

**Table 4: Mycotoxins content for wheat, barley and millet grain**

Mycotoxins	Wheat	Barley	Millet
Mycotoxins	*	*	*
Ochratoxin ppb	*	*	*
Zearalenone ppb	*	*	*
Fumonisin ppb	*	*	*
Aflatoxin B1	*	*	*
Aflatoxin B2	*	*	*
Aflatoxin G1	*	*	*
Aflatoxin G2	*	*	*
Total	*	*	*

\*= Under detection limit (0.50 ppb).

### 3.5 Physical properties of wheat, barley and millet kernels cultivars

Mean values of physical properties of wheat, barley and millet were presented in Table 5. It can be concluded that the test weight for all samples which ranged from 43.1 pound per bushel for millet to 60.1 pound per bushel for wheat. Percentage of shrunken and broken of wheat was (1.10 %) while thin and sound of barley was highest percentage (2.80 % - 95.46 %). For damage kernels which consist of heat damage and total damage, especially wheat have highest total damage kernels percentage (1.5 %) while barley and millet were lowest percentage of total damage kernels (0.83%). It can be noticed that the wheat, barley and millet haven't heat damage. More over from the same table noticed that all sample are free



from insect and OK odor. Results in Table 5 showed that weight per 1000 of kernels wheat, barley and millet have highest value (60.0 gm), barley 49.50 gm while wheat has lowest value (33.50 gm). For addition the kernel colour in wheat sample is red whereas barley is white and millets green. These results are in agreement with thus obtained by the Egyptian stander no. 1601/1986 and it's modification on 23/4/2002 [32] has obligation that the dockage % (first separated from sample) not exceed 1 %, foreign material % not exceed 1 %, total damage kernels % (heat damage ,sprout damage, insect damage and mould damage kernels) not exceed than 4 %. However that difference between wheat samples, all wheat samples had graded one according to [15].

**Table 5: physical properties of wheat, barley and millet kernels cultivars**

Parameters	Wheat	Barley	Millet
Moisture Content (M.C) %	10.4 <sup>a</sup>	10.2 <sup>a</sup>	8.70 <sup>b</sup>
Test weight (T.W) p/b	60.10 <sup>a</sup>	49.0 <sup>b</sup>	43.10 <sup>c</sup>
Broken kernels & Foreign Material (BNFM) %	0.20 <sup>b</sup>	1.0 <sup>a</sup>	0.77 <sup>a</sup>
Sh.& B.N %	1.10 <sup>a</sup>	0.33 <sup>b</sup>	0.45 <sup>b</sup>
Thin	-	2.80	-
Sound	-	95.46	-
Damage Kernels Heat Damage (H.D) %	0.0	0.0	0.0
(D.K) % Total Damage (T.D) %	1.50 <sup>a</sup>	0.83 <sup>b</sup>	0.83 <sup>b</sup>
Odor	Ok	Ok	Ok
Insect	Free	Free	Free
Weigh per 1000 kernels gm	33.50 <sup>c</sup>	49.50 <sup>b</sup>	60.0 <sup>a</sup>
Hardness %	61.0 <sup>b</sup>	50.0 <sup>c</sup>	75.0 <sup>a</sup>
Colour	Red	White	Green

*a,b,...Means with the same letter in the same row are not significantly different at (P>0.05).*

*p/b= Pound per Bushel (American unit),*

### 3.6 Physicochemical properties of wheat, barley, millet meals and their mixtures

The data in Table 6 showed that the highest starch damage was in barley meal (11.20 %) while mix 3 meal was the lowest (2.75 %). In the table it could be noticed from that the wet and dry gluten of control sample was 25.60 % and 7.70 % respectively, with a gluten index of 61.41. Upon substituting wheat with 25 % (barley and millet meal), wet and dry gluten contents were 20.10 % and 5.90 %, respectively, with a gluten index of 62.80, and also, increasing the level of barley and millet meal, the gluten content (either wet or dry) and the gluten index decreased. Gluten is responsible for the elasticity and extensibility characteristics of flour dough. Wet gluten reflects protein content and is a common flour specification required by end-users in the food industry. These results agree with work by [33]. From the same table it can be concluded that the percentage of sediment ranged from 10 to 28 ml for wheat and barley meals. The same table reviewed that the falling number values were ranged from 240 to 512 sec., and wheat meal had the highest value 512.0 sec. while mix 3 meal had lower values 240.0 sec.,. It can observe that addition of barley and millet at different level to wheat meal decrease the value of falling number and developed for enzyme activity of Alfa amylase and rheological properties of dough. Generally, a falling number value of 350 seconds or longer indicates low enzyme activity and very sound wheat. As the amount of enzyme activity increases, the falling number decreases. Economic European community recommended that the falling number of flour should exceed than 230 sec [34]. Also, [35] for durum wheat has obligation that protein content of durum wheat not less than 10.5 % and ash content not exceed than 1.3 %. At the end of the Table 6 it showed that the barley had the highest value of whiteness color 32.5 % and the lowest values of yellow color 14.53 %, then the millet meal which is less whiteness 2.96 % and highest value of yellow color 23.32 %. Flour color often affects the color of the finished product and is

therefore one of many flour specifications required by end-users. Generally speaking, bright white color flour is more desirable for many products. This result is agreement with [36].

**Table 6: physicochemical properties of wheat, barley, millet meals and their mixtures**

Parameters	Wheat	Barley	Millet	Mix 1	Mix 2	Mix 3
Starch damage %	7.00 <sup>b</sup>	11.20 <sup>a</sup>	7.05 <sup>b</sup>	4.70 <sup>c</sup>	4.75 <sup>c</sup>	2.75 <sup>d</sup>
Wet %	25.6 <sup>a</sup>	Free	Free	20.1 <sup>b</sup>	Free	Free
Gluten quantity Dry %	7.7 <sup>a</sup>	Free	Free	5.90 <sup>b</sup>	Free	Free
Hydration ratio %	17.9 <sup>a</sup>	Free	Free	14.20 <sup>b</sup>	Free	Free
Index %	61.4 <sup>b</sup>	Free	Free	62.80 <sup>a</sup>	Free	Free
Protein sediment %	10.0 <sup>e</sup>	28.0 <sup>a</sup>	25.0 <sup>b</sup>	18.00 <sup>d</sup>	20.00 <sup>c</sup>	24.00 <sup>b</sup>
Falling number sec.	512.0 <sup>a</sup>	431.0 <sup>b</sup>	254.0 <sup>e</sup>	349.0 <sup>c</sup>	290.0 <sup>d</sup>	240.0 <sup>f</sup>
flour colour % White	11.5 <sup>e</sup>	32.7 <sup>a</sup>	2.96 <sup>f</sup>	16.24 <sup>d</sup>	19.58 <sup>c</sup>	20.30 <sup>b</sup>
Yellow	20.63 <sup>b</sup>	14.53 <sup>f</sup>	23.32 <sup>a</sup>	18.74 <sup>c</sup>	17.40 <sup>d</sup>	16.98 <sup>e</sup>

*a,b,...Means with the same letter in the same row are not significantly different at (P>0.05)*

*Free= free of wheat gluten.*

### 3.7 Rheological properties of wheat, barley, millet meals and their mixtures

Water absorption (WA) is a parameter indicated as the amount of water needed to develop the standard dough at the peak of the curve. Consistographe parameters of the macaroni flours resulted from different grain meals and their mixtures showed that water absorption (WA %) decreased from 54.6 % for the control sample made from wheat to 35.0 % for millet in Table 7. Higher water absorption is required for good bread characteristics which remain soft for a longer time. The gradual decrease in WA % was found to be due to decreasing the level of barley and millet from 25 to 75 %. This decrease can be attributed to lower gluten-starch network formation which is responsible for water absorption, as the ratio of wheat in blends. Those results in Paolo with those obtained by [37] who said that the presence of damaged starch tends to increase water absorption. And [38] reported that the inclusion of a higher amount of bran in the dough formulation usually resulted in increased dough water absorption due to the higher levels of pentosans present in bran. The alveograph determines the gluten strength of dough by measuring the force required to blow and break a bubble of dough. The Tenacity (P) was (156 mm H<sub>2</sub>O) for wheat to (19.0 mm H<sub>2</sub>O) for mix 2 Table 7. So that wheat flour was the high significant effect value (15 mm, 8.60 ml) of Expandability and Swelling (G) while mix 2 was the low significant effect (9.00 mm, 6.70 ml). The P/L value is high significant effect in mix 1 (14.70 %) and the low significant effect was mix 2 (6.70 ml). Baking strength (W) was the high significant effect in wheat (108.0 jol). W is the most widely used characteristic because it summaries all the others. The very different shapes of the curves from 'extreme' individuals indicate the great variation in dough strength and extensibility present in the core collection. Also, table 7 showed the transition point, maximum viscosity and temperature at maximum viscosity as measured by amylograph. The data revealed that transition point (°C) of wheat was 68.45 °C followed by mix 1 (63.91 °C). The maximum viscosity was arranged in the descending order as follows: barley (511.40 B.U.) > wheat (342.10 B.U.) which in parallel with the temperature of 94.0 °C and 92.0 °C, respectively. Our results are in agreement to work by [39] how reported that amylograph parameters indicated that hull-less barley had lower gelatinization temperature and higher maximum viscosity than the hull-barley as a result of the presence of beta-glucan with a higher ratio in hull-less barley. Also, [40] suggested that a reduction in maximum viscosity of hull-barley may be associated with a reduced enthalpy of starch gelatinization, and retention of the integrity of the starch granule.

**Table 7: Rheological properties of wheat, barley, millet meals and their mixtures**

Parameters		Wheat	Barley	Millet	Mix 1	Mix 2	Mix 3
Consist graph Water absorption %		54.60 <sup>a</sup>	47.2 <sup>d</sup>	35.0 <sup>e</sup>	52.40 <sup>b</sup>	49.5 <sup>c</sup>	47.3 <sup>d</sup>
Alveograph test	Tenacity mm H <sub>2</sub> O (P)	156.00 <sup>a</sup>	-	-	133.00 <sup>a</sup>	19.0 <sup>b</sup>	-
	Expandability mm (L)	15.00 <sup>a</sup>	-	-	10.00 <sup>ab</sup>	9.00 <sup>b</sup>	-
	Swelling ml (G)	8.60 <sup>a</sup>	-	-	7.00 <sup>b</sup>	6.70 <sup>b</sup>	-
	Baking strength Jol (W)	108.00 <sup>a</sup>	-	-	61.00 <sup>b</sup>	9.0 <sup>c</sup>	-
Configuration rate % (p/L)		10.40 <sup>b</sup>	-	-	14.78 <sup>a</sup>	1.9 <sup>c</sup>	-
Amylograph	Transition point (C°)	68.45 <sup>a</sup>	55.21 <sup>d</sup>	45.32 <sup>f</sup>	63.91 <sup>b</sup>	59.35 <sup>c</sup>	54.82 <sup>e</sup>
	Maximum viscosity (B.U.)	342.1 <sup>f</sup>	511.4 <sup>a</sup>	501.9 <sup>b</sup>	383.2 <sup>e</sup>	424.1 <sup>d</sup>	465.4 <sup>c</sup>
	Temperature at maximum viscosity (C°)	92 <sup>a</sup>	94 <sup>a</sup>	96 <sup>a</sup>	91 <sup>a</sup>	93 <sup>a</sup>	94 <sup>a</sup>

279 *a,b,...Means with the same letter in the same row are not significantly different at (P>0.05).*

280

### 281 **3.8 Chemical composition of macaroni obtained from wheat, barley, millet**

### 282 **meals and their mixtures**

283 The chemical composition of macaroni produced from the different levels of wheat, barley,  
 284 millet meals was reported in Table 8. The data revealed that no significant effect of the  
 285 moisture content for macaroni meal. Protein content decreased from 13.10 gm/100 gm for  
 286 control to 9.60 gm/100 gm for barley. These findings were in close range with those of  
 287 [33]. Fat content increased from 1.11 gm/100 gm for wheat macaroni to 3.3 gm/100 gm for  
 288 millet. The results are in agreement with work by [41]. Ash content increased from 1.00  
 289 gm/100 gm for barley macaroni to 1.5 gm/100 gm for millet macaroni. The increase in ash  
 290 content may be due to the higher ash content of millet (1.80 gm/100 gm). As for crude fiber,  
 291 the content increased from 1.30 gm/100 gm for wheat macaroni to 7.20 gm/100 gm for  
 292 millet; this may be due to the high fiber content of millet compared with wheat. Total  
 293 carbohydrates decreased from 74.9 (for barley) to 66.10 gm/100 gm (for millet). Total caloric  
 294 values increased from 337.3 to 351.5 gm/100 gm for millet and barley, respectively. These  
 295 findings were in close range with those of [33].

296

297 **Table 8: Chemical composition of macaroni obtained from wheat, barley, millet meals**  
 298 **and their mixtures**

Chemical composition	Wheat	Barley	Millet	Mix 1	Mix 2	Mix 3
Moisture content %	12.5 <sup>a</sup>	10.8 <sup>a</sup>	11.1 <sup>a</sup>	12.1 <sup>a</sup>	11.7 <sup>a</sup>	11.3 <sup>a</sup>
Protein content %	13.1 <sup>a</sup>	9.6 <sup>c</sup>	10.8 <sup>bc</sup>	12.5 <sup>ab</sup>	11.9 <sup>ab</sup>	11.3 <sup>bc</sup>
Fat content %	1.11 <sup>b</sup>	1.5 <sup>b</sup>	3.3 <sup>a</sup>	1.43 <sup>b</sup>	1.8 <sup>b</sup>	2.1 <sup>ab</sup>
Ash content %	1.22 <sup>ab</sup>	1.0 <sup>b</sup>	1.5 <sup>a</sup>	1.22 <sup>ab</sup>	1.23 <sup>ab</sup>	1.24 <sup>ab</sup>
Fiber content %	1.3 <sup>d</sup>	2.2 <sup>cd</sup>	7.2 <sup>a</sup>	2.15 <sup>cd</sup>	3.0 <sup>bc</sup>	3.85 <sup>b</sup>
Carbohydrates %	70.77 <sup>b</sup>	74.9 <sup>a</sup>	66.1 <sup>c</sup>	70.6 <sup>b</sup>	70.37 <sup>b</sup>	70.21 <sup>b</sup>
Total caloric values%	345.47 <sup>b</sup>	351.5 <sup>a</sup>	337.3 <sup>c</sup>	345.27 <sup>b</sup>	345.28 <sup>b</sup>	344.94 <sup>b</sup>

299 *a,b,...Means with the same letter in the same row are not significantly different at (P>0.05).*

300

301



### 3.9 Effect of macaroni obtained from wheat, barley, millet meals and their mixtures on the cooking quality

Cooking performance is an important factor in a consumer's judgment of macaroni quality. Table 9 revealed that macaroni cooking time decreased from control to mix 3 (10.00 to 5.5 min) with increased level of mixed. Addition of mixed meal to the macaroni resulted in lower cooking time for complete gelatinization of starch as compared to control. The data revealed that no difference significant in macaroni weight. Similar trend was found regarding volume. While [42] explained such trend be the high levels of total dietary fiber and  $\beta$ -glucan in barley and as a result increasing the water holding capacity of macaroni. The cooking loss was increased by substitution with barley 10.34 % compared to wheat macaroni 4.35 %, then after increased level of mixed. The cooking loss is an indicator of the capability of the starch-protein matrix to retain its physical integrity during cooking [43], and only values lower than 7 % are acceptable for a good quality pasta [44]. Generally, non-starch polysaccharide addition increased the cooking loss [45]. Also, [46] explained that increased amount of barley present in the semolina matrix had disrupted the protein-starch network, causing starches to leach out during the cooking, and consequently resulting in a decrease in pasta cooking quality.

**Table 9: Effect of macaroni obtained from wheat, barley, millet meals and their mixtures on the cooking quality**

	Treatments	Cooking time	Weight increase	Volume increase	Cooking loss
		(minutes)	(%)	(%)	(%)
Macaroni meal	Wheat	10.0 <sup>a</sup>	180 <sup>a</sup>	190 <sup>a</sup>	4.35 <sup>d</sup>
	Barley	7.7 <sup>b</sup>	196 <sup>a</sup>	205 <sup>a</sup>	10.3 <sup>a</sup>
	Millet	5.9 <sup>c</sup>	192 <sup>a</sup>	200 <sup>a</sup>	5.5 <sup>c</sup>
	Mix 1	6.5 <sup>bc</sup>	181 <sup>a</sup>	188 <sup>a</sup>	4.52 <sup>d</sup>
	Mix 2	6.2 <sup>bc</sup>	190 <sup>a</sup>	200 <sup>a</sup>	5.25 <sup>c</sup>
	Mix 3	5.5 <sup>c</sup>	192 <sup>a</sup>	200 <sup>a</sup>	8.15 <sup>b</sup>

*a,b,...Means with the same letter in the same row are not significantly different at ( $P>0.05$ ).*

### 3.10 The change in density as affected by cooking of macaroni made from wheat, barley, millet meals and their mixtures

To confirm the rheological data, density of macaroni was determined before and after cooking Table 10. The data revealed that no difference significant in macaroni volume before cooking, but after cooking millet was highest values 28.0 cm<sup>2</sup> and wheat was the lowest values of volume 23.0 cm<sup>2</sup>. So that revealed that no difference significant in macaroni's weight and density before cooking but millet macaroni was the highest value for weight of macaroni after cooking. Mix 2 was highest values of density after cooking. The results agree with work by [33]. This decrease may be due to the amount of water absorbed during cooking.

**Table 10: The change in density as affected by cooking of macaroni made from wheat, barley, millet and their mixtures before and after cooking**

	Treatments	Volume (cm <sup>2</sup> )		Weight (gm)		Density (gm/ cm <sup>2</sup> )	
		(Before)	(After)	(Before)	(After)	(Before)	(After)
Macaroni meal	Wheat	7.25 <sup>a</sup>	10.0 <sup>a</sup>	10.01 <sup>a</sup>	29.08 <sup>c</sup>	1.38 <sup>a</sup>	1.26 <sup>ab</sup>
	Barley	7.25 <sup>a</sup>	7.7 <sup>b</sup>	10.0 <sup>a</sup>	30.71 <sup>a</sup>	1.38 <sup>a</sup>	1.13 <sup>b</sup>
	Millet	7.26 <sup>a</sup>	5.9 <sup>c</sup>	10.05 <sup>a</sup>	31.0 <sup>a</sup>	1.38 <sup>a</sup>	1.11 <sup>b</sup>
	Mix 1	7.23 <sup>a</sup>	6.5 <sup>bc</sup>	10.03 <sup>a</sup>	29.85 <sup>b</sup>	1.38 <sup>a</sup>	1.25 <sup>ab</sup>
	Mix 2	7.21 <sup>a</sup>	6.2 <sup>bc</sup>	10.01 <sup>a</sup>	30.02 <sup>b</sup>	1.38 <sup>a</sup>	1.33 <sup>a</sup>
	Mix 3	7.24 <sup>a</sup>	5.5 <sup>c</sup>	10.0 <sup>a</sup>	30.26 <sup>b</sup>	1.38 <sup>a</sup>	1.16 <sup>b</sup>

*a,b,...Means with the same letter in the same column are not significantly different at ( $P>0.05$ ).*

### 3.11 Effect of adding different wheat, barley, millet meals and their mixtures on color parameters of macaroni product

Color plays a major role in consumer's perception and acceptability of the product. The observed color value of cooked macaroni with different combinations of the ingredients varied from L=92, a=-0.91 and b=10.41 for wheat while for millet flour the values were 52.0, -0.51 and -5.88 for L, a and b respectively, as shown in Table 11. And in this table represents change in lightness (L\*) value of macaroni millet significantly decreased the lightness (L\*) value of prepared macaroni. As the level of mixed meal (barley and millet) increased, the lightness (L\* value) and redness (a\* value) decreased, but the brightness (b\* value) increased, this increase may be due to presence of barley and millet which gives macaroni a yellow tint, as they are rich sources of carotenoids. This may be due to the brick red color of finger millet seed coat and grey color of pearl millet flour [47]. Also, [48] observed that L\* value of pasta prepared from native pearl millet was lower than the pasta prepared from depigmented pearl millet flour. This difference in color of millet flours is due to the polyphenolic pigments present in pericarp, aleuronic layer and in endosperm region [49].

**Table 11: Effect of adding different wheat, barley, millet meals and their mixtures on color parameters of macaroni product**

Treatments		Brightness "L" L*	Redness "a" a*	Yellowness "b" b*
Macaroni meal	Wheat	92.0 <sup>a</sup>	-0.91 <sup>e</sup>	10.41 <sup>a</sup>
	Barley	86.5 <sup>ab</sup>	-0.61 <sup>bc</sup>	8.97 <sup>c</sup>
	Millet	52.0 <sup>d</sup>	-0.51 <sup>b</sup>	-5.88 <sup>d</sup>
	Mix 1	86.31 <sup>ab</sup>	-0.18 <sup>a</sup>	9.66 <sup>b</sup>
	Mix 2	80.63 <sup>bc</sup>	-0.65 <sup>cd</sup>	-8.17 <sup>e</sup>
	Mix 3	74.94 <sup>c</sup>	-0.74 <sup>d</sup>	-8.92 <sup>f</sup>

a,b,c,d...Means with the same letter in the same column are not significantly different at (P>0.05).

### 3.12 Sensory evaluation of macaroni made from wheat, barley, millet meals and their mixtures

Table 12 revealed that a high significant differences in appearance at wheat and barley then mix 3 macaroni (17.14, 17.0 and 16.35 %), respectively. Meanwhile, a highly significant decrease was found as a result of millet macaroni (10.30 %). These results agree with work by [33]. Flavor showed significant decrease in millet macaroni but all produced macaroni showed that a non-significant differences in flavor. Taste showed high significant difference at wheat macaroni then barley then mix 1 (17.14, 16.28, 14.21 %), respectively. The texture of macaroni was found maximum with barley + mix 1 and lowest was found with millet. Color showed high significant difference at mix 2 and low significant difference in millet (8.78 - 5.57 %). It could be noticed that the overall quality values of tested macaroni were found to be high acceptable and scores ranged between 85.19, 84.91 % for barley and control then after that mix (1) 78.29 % but the lowest was for millet 51.88 %. Sensory evaluation is most reliable test as it allows overall characteristics of cooked macaroni. The overall acceptability of cooked macaroni within the combinations varied from 51.88 to 85.19. It was shown decreased overall acceptability by increasing the proportion of barley and pearl millet meals. This may be due to unattractive dark color of finger millet and grey to yellow color of millet which limits the wider acceptability of its food products.

**Table 12: Sensory evaluation of macaroni made from wheat, barley, millet meals and their mixtures**

Macaroni	Appearance 20 %	Flavour 20 %	Taste 20 %	Texture 10 %	Colour 10 %	Mouth feeling 20 %	Overall 100 %
Wheat	17.14 <sup>a</sup>	17.0 <sup>a</sup>	17.14 <sup>a</sup>	8.07 <sup>b</sup>	7.92 <sup>c</sup>	17.64 <sup>b</sup>	84.91 <sup>a</sup>
Barley	17.0 <sup>a</sup>	16.85 <sup>a</sup>	16.28 <sup>b</sup>	8.50 <sup>a</sup>	8.64 <sup>ab</sup>	17.92 <sup>a</sup>	85.19 <sup>a</sup>
Millet	10.28 <sup>e</sup>	10.30 <sup>b</sup>	10.13 <sup>f</sup>	5.10 <sup>d</sup>	5.57 <sup>e</sup>	10.50 <sup>d</sup>	51.88 <sup>d</sup>
Mix 1	15.78 <sup>c</sup>	14.35 <sup>a</sup>	14.21 <sup>c</sup>	8.39 <sup>a</sup>	8.60 <sup>b</sup>	16.96 <sup>c</sup>	78.29 <sup>b</sup>
Mix 2	15.28 <sup>d</sup>	14.0 <sup>a</sup>	12.84 <sup>e</sup>	8.07 <sup>b</sup>	8.78 <sup>a</sup>	17.59 <sup>b</sup>	76.56 <sup>c</sup>
Mix 3	16.35 <sup>b</sup>	14.07 <sup>a</sup>	14.07 <sup>d</sup>	7.82 <sup>c</sup>	7.28 <sup>d</sup>	16.82 <sup>c</sup>	76.36 <sup>c</sup>

*a,b,...Means with the same letter in the same colum are not significantly different at (P>0.05).*

#### 4. CONCLUSION

From the present study, it is found that of wheat, barley and millet meals fortified macaroni offer a broader spectrum for people wishing to improve the nutritional quality of their diet. Barley and millet are highly nutritious, rich in health promoting photochemical and dietary fiber. The mixed macaroni was slightly darker in appearance. Macaroni made of mixed meal grains showed lower water absorption and higher volume. The present study showed that macaroni with good nutritional and functional properties can be obtained from barley then mix 1, mix 2, and mix 3, respectively. Mixed meal grains could be effectively utilized for high quality macaroni which will increase the meal grain consumption and likely to reduce the risk of degenerative diseases.

#### REFERENCES

1. Slavin J. Whole grain and human health. *Nutr Res Rev.* 2004; 17: 99-100.
2. Liu S, Manson J E, Stampfer M J, Hu F B, Giovannucci E, Coeditz G A, et al., A prospective study of whole grain intake and risk of type 2 diabetes mellitus in US women. *Am J Public Health*, 2000; 90: 1409-1415.
3. Pereira M A, Jacobs D R, Pins J J , Raatz S K, Gross M D, Slavin J L et al., Effect of whole grains on insulin sensitivity in overweight hyperinsulinemic adults. *Am J Clin Nutr* 2002; 75: 848-855.
4. Jacobs D R, Marquart L, Slavin J and Khushi L H. Whole grain intake and cancer: an expanded review and Meta analysis. *Nutr Cancer* 1998a; 30: 85-96.
5. Jacobs D R, Mayer K A, Khushi L H and Folsom A R. Whole grain intake may reduce the risk of ischemic wart disease death in postmenopausal woman: the Iowa Women's Health Study. *Am J Clin Nutr*, 1998b; 68: 248-257.
6. Petitot, M, L Bayer, C Minier, V Micard. Fortification of pasta with split pea and faba bean flours: pasta processing and quality evaluation. *Food Research International*, 2010; 43: 634-641.
7. Rødbotten, M, Tomic, O, Holtekjølén, AK, Grini, IS, Lea, P, Granli, BS et al., Barley bread with normal and low content of salt; sensory profile and consumer preference in five European countries. *Journal of Cereal Science*, 2015; 64: 176-182.
8. FDA. Food and Drug Administration is a federal agency of the United States Department of Health and Human Services, one of the United States federal executive departments. Barley beta fiber and coronary heart disease 2006. Available at <http://www.fda.gov/ohrms/dockets/dockets/06p0393/06p-0393-cp00001-002-vol1.pdf>.
9. Vidyavati, H G, Begum, M G, Vijayakumar, J, Gokavi, S S, and Begum, S. Utilization of finger millet in preparation of Papad. *Journal of Food Science and Technology*, 2004; 41(4), 379-382.

- 424 10. Sarita, Ekta Singh. Potential of Millets: Nutrients Composition and Health Benefits. *Journal*  
425 *of Scientific and Innovative Research* 2016; 5(2): 46-50.
- 426 11. Chillo, S, J Laverse, PM, Falcone and MA Del Nobile. Quality of spaghetti in base  
427 Amaranthus whole meal flour added with quinoa, broad bean and chick pea. *Journal of*  
428 *Food Engineering*, 2008; 84: 101-107.
- 429 12. Tudorica, CM, Kuri, V and Brennan, CS. Nutritional and Physicochemical characteristics  
430 of Dietary fiber enriched pasta. *Journal of Agriculture and Food Chemistry*. 2002; 50:347-  
431 356.
- 432 13. Montalbano, A, L Tesoriere, P Diana, P Barraja, A Carbone, V Spanò, et al. Quality  
433 characteristics and in vitro digestibility study of barley flour enriched ditalini pasta. *LWT-*  
434 *Food Science and Technology*, 2016; 72: 223-228.
- 435 14. Gopalan, C, Ramasastry, BV and Balasubramanian, SC. *Nutritive value of Indian*  
436 *Foods*. National Institute of Nutrition 2000, Indian Council of Medical Research,  
437 Hyderabad, India.
- 438 15. USDA. U.S. DEPARTMENT OF AGRICULTURE, GRAIN INSPECTION HANDBOOK I.  
439 Grain Inspection, Packers and Stockyards Administration, Federal Grain Inspection  
440 Service Probe Sampling, Washington, D.C. 2013; 20090-6454.
- 441 16. USDA, U.S. Department of Agriculture, EQUIPMENT HANDBOOK. grain Inspection,  
442 Packers and Stockyards Administration, S.W. Washington, D.C. 2016.
- 443 17. AACC. American association of cereal chemists, Approved method of the AACC 10<sup>th</sup> ed.,  
444 AACC, St Paul, MN. 2000A; vol. 1.
- 445 18. AACC, American association of cereal chemists, Approved method of the AACC 10<sup>th</sup> ed.,  
446 AACC, St Paul, MN. 2000B; vol. 2.
- 447 19. AOAC. Association of Official Analytical Chemists. Official Methods of Analysis. 18<sup>th</sup> Ed.  
448 Published by A.O.A.C. W. Horwitz. North Frederick, U.S.A 2005.
- 449 20. Pérez-Vendrell A, Guash J, Frances M, Molina-Cano JL, Brufau J. Determination of  $\beta$ -  
450 (1 $\rightarrow$ 3), (1 $\rightarrow$ 4)-D-glucans in barley by reverse-phase high performance liquid  
451 chromatography. *Journal of Chromatography A*, 1995; 718, 291–297.
- 452 21. Fany, K and K Khan. Pasta containing rigids: Effect of high temperature drying on  
453 products quality. *Cereal Chemistry*, 1996; 73: 317-322.
- 454 22. SAS. SAS / Stat. User's Guide: statistics, system for windows, version 4.10 (release  
455 6.12 TS level 0020), SAS Inst., 2011; Inc. Cary, North Carolina, USA.
- 456 23. Hatcher, DW, S Lagasse, JE Dexter, B Rosnagel and M Izydorczyk. Quality  
457 Characteristics of Yellow Alkaline Noodles Enriched with Hull-less Barley Flour 1. *Cereal*  
458 *Chemistry*, 2005; 82(1): 60-69.
- 459 24. Mandge H M, Sharma S and Dar B N. Instant multigrain porridge: effect of cooking  
460 treatment on physicochemical and functional properties. *J Food Sci and*  
461 *Technol*. 2014; 51: 97-103.
- 462 25. Abdalla A A, Ahmed U M, Ahmed A R, El Tinay A H and Ibrahim K A. Physicochemical  
463 characterization of traditionally extracted pearl millet starch (Jir). *J Appl Sci Res*, 2009; 5:  
464 2016-27.
- 465 26. Hejazi, MA. Using Hull-Less Barley and Flax Seeds Flour to Produce Macaroni for  
466 Hyperglycemia Disease in Rats. *Life Science Journal*, 2014; 11(2): 354-361.
- 467 27. Dahab, D B O M. Utilization of different cereal flour mixes in the preparation of some  
468 bakery products. M.Sc., Food Sci. and Technol. 2006; Dept; Fac. Agric; Cairo Univ;  
469 Egypt.
- 470 28. Pawar VD, Machewad GM. Processing of foxtail millet for improved nutrient availability. *J*  
471 *Food Process Preserv*, 2006; 30: 269–279.
- 472 29. NIN, National Institute of Nutrition. Indian Foods Nutritional Value, 2003.
- 473 30. Sandberg, A S. Bioavailability of minerals in legumes. *Br J Nutr*. 2002; 88 (3): S281-  
474 S285.
- 475 31. FAO. Food and Agricultural Organisation, Food and Agriculture Bulletin on Food Security,  
476 (2009). [www.fao.org/docrep/x5030E/X15032E06.htm](http://www.fao.org/docrep/x5030E/X15032E06.htm).

- 477 32. ES. Egyptian Standard of wheat grains. Egyptian Organization for Standardization and  
478 Quality Control, No. 1601, 1986 and its modification No. 2/2002. Arab Republic of Egypt.  
479 33. Salem, Eman M. Quality Attributes of Pasta Substituted with Barley Meal. Egyptian  
480 Journal of Nutrition, 2005; 20(2): 25-43.  
481 34. ES. Egyptian Standard of white flour for production of bread. Egyptian Organization for  
482 Standardization and Quality Control, No. 1419, 2006. Arab Republic of Egypt.  
483 35. ES. Egyptian Standard of durum wheat. Egyptian Organization for Standardization and  
484 Quality Control, No. 1649, 2004. Arab Republic of Egypt.  
485 36. D'Appolonia, B and L Emeritus. How Flour Affects Bread Quality Department of Cereal  
486 Science, North Dakota State University, Fargo, ND 58105, Lallemand Baking Update,  
487 VOL.1 /No. 17, 1996.  
488 37. Young, TL, HM Seong, MK Cho and S Kim. Physiochemical properties of hull-less barley  
489 flours prepared with different grinding mills. Korean Journal of Food Science and  
490 Technology, 1996; 28(6): 1078-1083.  
491 38. Sanz-Penella, JM, C Collas, SM Haro. Effect of wheat bran and enzyme addition on  
492 dough functional performance and phytic acid levels in bread. Journal of Cereal Science,  
493 2008; 48: 715-721.  
494 39. Lee, MJ, KS Kwon and HG Chang. The physico-chemical properties and cooking  
495 qualities of barley isogenic lines. Journal of the Korean Society for Applied Biological  
496 Chemistry, 1997; 40(4): 301-306.  
497 40. Symons, LJ and CS Brennan. The influence of (1→3, 1→4)-Beta-D-glucan-rich fractions  
498 from barley on the physicochemical properties and in-vitro reducing sugar release of  
499 white wheat breads. Journal Food Science, 2004; 69: 463-467.  
500 41. Sawsan, Y El-Faham, Eid A Abd El-Hamid and Hussein K Ashour. Barley Flour and  
501 Durum Flour Blends in Macaroni Product. Australian Journal of Basic and Applied  
502 Sciences, 2010; 4(12): 6169-6178.  
503 42. Zahran, GAH, NM Abd El-Motaleb and OSR Shams. Chemical and biological functional  
504 aspects of pasta rich in dietary fiber and B-glucan. Egyptian Journal of Agricultural  
505 Research, 2004; 82(3): 13-25.  
506 43. Bruneel, C, B Pareyt, K Brijs and JA Delcour. The impact of the protein network on the  
507 pasting and cooking properties of dry pasta products. Food Chemistry, 2010; 120: 371-  
508 378.  
509 44. Sissons, MJ, J Abecassis, R Cubadda and B Marchylo. Methods used to assess and  
510 predict quality of durum wheat, semolina, and pasta. In M J Sissons, J Abecassis, B  
511 Marchylo & M Carcea (Eds.), Durum wheat Chemistry and technology, 2012; (pp.  
512 213e234). St Paul, MN: AACC International.  
513 45. Sandhu, GK, S Simsek and FA Manthey. Effect of xanthan gum on processing and  
514 cooking quality of nontraditional pasta. International Journal of Food Science and  
515 Technology, 2015; 50: 1922-1932.  
516 46. Makhlof, S. Feasibility study of incorporating fiber into pasta and effects on product  
517 quality attributes 2015; (Doctoral dissertation, California State Polytechnic University,  
518 Pomona).  
519 47. Gull, A, Prasad, K, and Kumar, P. Effect of millet flours and carrot pomace on cooking  
520 qualities, color and texture of developed pasta. LWT Food Science and Technology, 2015  
521 ;63(1), 470-474. <http://dx.doi.org/10.1016/j.lwt.2015.03.008>.  
522 48. Rath A, Kawatra A, and Sehgal, S. Influence of de-pigmentation of pearl millet  
523 (*Pennisetum glaucum*) on sensory attributes, nutrient composition, in vitro protein and  
524 starch digestibility of pasta. Food Chemistry, 2004; 85(2), 275-280.  
525 <http://dx.doi.org/10.1016/j.foodchem.2003.06.021>.  
526 49. McDonough, C Mand Rooney, LW. Structural characteristics of *Pennisetum americanum*  
527 (pearl millet) using scanning electron and fluorescence microscopy. Food Microstructure,  
528 1989; 81, 137-149.  
529