

1
2
3
4
5
6
7
8
9
10
11
12

Original Research Article

Spore density and arbuscular mycorrhizal colonization in sunflower

ABSTRACT

The objective of this study was to evaluate the number of spores and mycorrhizal root colonization in Cerrado soil (Red-Yellow Latosol) cultivated with different genotypes of sunflower. Sampling of rhizospheric soil occurred in three periods: sowing, flowering and sunflower harvest. The experimental design was a randomized complete block design with four replications. To proceed the evaluations different sunflower hybrids were selected: M 734, Agrobela 960 and Helio 358, in 2009, and M 734, Embrapa 122 and HLA 860 H.O. in 2010. The measured parameters were number of total spores in 50 g of soil in three periods: sowing, flowering and harvesting and the arbuscular mycorrhizal colonization was evaluated only in the final of the experiment. The mean number of spores was 4,94 g soil⁻¹ and 4,64 g soil⁻¹ in 2009 and 2010, respectively. The maximum spore production occurred during the flowering period and mycorrhizal colonization was not influenced by the genotype. The mycorrhizal colonization rate ranged from 21 to 28% in 2009 and from 28 to 48% in 2010. The number of spores varies from 153 to 342 in 2009 and 147 to 320 in 2010, and the maximum production occurs, on average, in the flowering period. Lower soil phosphorus levels favors arbuscular mycorrhizal colonization.

13 *Keywords: Helianthus annuus L.; soil; arbuscular mycorrhizal fungi, root colonization.*

1. INTRODUCTION

14
15
16 Soil quality and the viability of improvements through chemical, physical and biological
17 management are essential factors for success in agricultural production. In this context, the
18 study and the use of soil microbial population has shown the way to link sustainability to
19 efficiency.

20 The symbiotic association between plant and fungi is called mycorrhiza. Root colonization by
21 arbuscular mycorrhizal fungi (AMF) generates several improvements; the plant provides
22 photosynthates to the fungus, and this, through the branching and extension of the
23 mycelium, increases the area of nutrient absorption for the plant [1]. Thus, AMFs can be
24 used as an alternative to reduce the use of agricultural inputs, mainly fertilizers of chemical
25 synthesis.

26 The influence of arbuscular mycorrhizal fungi acts not only on soil particles aggregation but
27 also on plant growth, providing essential nutrients [2] and improving their ability to withstand
28 adverse conditions.
29

30 Studying AMF inoculation in sunflower, it was observed an increase in chapter diameter,
 31 thousand achenes weight and achenes yield. These traits parameters that were related to
 32 the better development of the plants through their association with AMFs, due to the higher
 33 absorption of nutrients such as P, K and Fe [3].

34 The sunflower cultivation (*Helianthus annuus* L.) cultivation has aroused interest, especially
 35 in Brazilian Midwest, due to its the broad adaptability to edaphoclimatic conditions, suitability
 36 for crop rotation and usage uses in the production of edible oil, biodiesel, ornamental
 37 crop, animal feed, etc., among others [4,5].

38 Considering that in the soils of the Cerrado Biome, for the optimization of the agricultural
 39 production, is necessary the use of a high amount of inputs, and that the agronomic
 40 efficiency is tied to the good indexes of soil quality, The present work aimed to evaluate
 41 three sunflower genotypes on the basis of their root mycorrhizal colonization at three
 42 different growth stages the number of spores in different times and mycorrhizal colonization
 43 in Cerrado Biome soils, under cultivation of three sunflower genotypes.

44 2. MATERIAL AND METHODS

45 The experiment was carried out at Santa Luzia Farm, in Campo Verde (MT-Brazil), latitude
 46 15°45'12"S and longitude 55°22'44"W. The soil in the experiment was a area is classified
 47 as Red-Yellow Latosol with the following properties: clayed texture, acidic pH, 50% average
 48 bases saturation of 50%, absence of aluminum and high content of organic matter content
 49 (Table 1). Soybean and corn were most crops grown under in the farm the system of
 50 minimum soil tillage practiced over more than cultivation is adopted, for more than ten
 51 years, being the most used crops soybean and corn. Over both cropping seasons
 52 considered (2009 and 2010), the species that preceding crop of sunflower was the
 53 sunflower in both years was soybean.

54 **Table 1. Chemical and physical properties of soil under sunflower cultivation in the**
 55 **2009 and 2010 harvests at Farm Santa Luzia, Campo Verde – MT, Brazil**

Year	pH CaCl ₂	P	K	Ca	Mg	Al	H	OM	CTC
		mg dm ⁻³		cmol _c dm ⁻³			gdm ⁻³		cmol _c dm ⁻³
2009	5,1	21,8	76	3,2	0,9	0	4,4	37,8	8,7
2010	4,9	8,0	80	3,3	0,7	0	5,5	39,9	9,7
Bases saturation (V%)		Sand	Silt	Clay	Saturation (%)				
		g kg ⁻¹			Ca	Mg	K	H	
2009	49,3	196	133	671	36,7	10,5	2,3	50,7	
2010	43,3	172	200	628	33,9	6,8	2,1	56,7	

56 P: phosphorous; K: potassium; Ca: calcium; Mg: magnesium; Al: aluminium; OM: organic
 57 matter; CTC: cation exchange capacity; H: hydrogen.

58 The experimental design used applied in the field was a randomized complete blocks, with
 59 four replications. Every The plots was composed of were formed by four rows of 6.0
 60 meters, with spaced in 0.8 meters of inter-row spacing, between rows, and 0.3 meters
 61 spacing within rows (19.2 m²). Two central rows (9.6 m²) were weighted at harvest to
 62 determine crop yield, between plants, considering as useful area the two central rows. NPK
 63 and boron The fertilizers were applied 30 days after sowing respectively at the following
 64 rates: used was 30-80-80 kg ha⁻¹ of NPK and 2.0 kg ha⁻¹ of boron on the sowing hill and 30

65 kg ha⁻¹ of N top-dressing, at 30 days after sowing. The 2010 cropping season was more
66 rainy compared to 2009, with respectively 974 and 442 mm of total precipitation (The rainfall
67 distribution in the region, during the experiment, is shown in Table 2).

68 **Table 2. Rainfall distribution (in mm/month) over 2009 and 2010 cropping seasons (mm**
69 **month⁻¹) in Campo Verde - MT, from February to July, in 2009 and 2010**

Year	February	March (S)	April	May (F)	June	July (H)	Total
2009	262	132	16	10	22	0,2	442,4
2010	385	206	325	55	3	2	974,0

70 S: sowing; F: flowering; H: harvest.

71

72 Three different sunflower hybrids were evaluated on the basis of their response to Fe
73 evaluate whether the response to fungi colonization was linked to genetic differences in
74 sunflower different hybrids were selected. Hybrids M 734, Agrobela 960 and Helio 358 were
75 used in 2009, whereas M 734, Embrapa 122 and HLA 860 H.O. in 2010.
76 Rhizospheric soil sampling was obtained at harvest over 0-20 cm depth and at
77 three different growth stages periods, namely sowing (first half of March), flowering (60 days
78 after sowing) and harvesting (after maturation).

79 Both the evaluated parameters evaluated were total number of spores in soil, and
80 arbuscular mycorrhizal colonization, whose root sampling occurred during crop harvest. The
81 spore extraction of spores was carried out by the wet sift methodology [6], in which the soil
82 was processed in a sieving systems (0.42 and 0.053 mm mesh) and centrifuged with water
83 at 2800 rpm for 4 min. Subsequently, the samples were re-suspended in 50% sucrose
84 solution, centrifuged and washed. The spores were counted using a stereomicroscope in
85 a petri dishes with vessels.

86 To check for the mycorrhizal colonization, the crop roots were washed, clarified with KOH
87 (10%), acidified with diluted HCl [7] and stained with trypan blue [8]. Ten segments of 1-2 cm
88 in length were selected for slide assembly. The determination of the quantification of root
89 colonization percentage was made using an optical microscope (40x) was done
90 considering the number of colonized root.

91 The Analysis of variance was calculated using software package were preceded, and
92 the significant differences between means were determined following compared by Tukey
93 test at with 5% of significance.

94 3. RESULTS AND DISCUSSION

95

96 Regarding the factor For the factor-year, there was no significant difference in the number of
97 spores of AMF (Table 3). This could be explained by a general improvement in soil fertility
98 resulting from may occurred since the studied area adopted the practice of minimum soil
99 tillage over cropping system for more than 10 years. According to Carrenho et al the authors
100 [9], the dissemination of association and mycorrhizal propagules dissemination is more
101 affected is much more affected during in the initial phases of land use, the occupation and use
102 of the soil, with later stabilization.

103

104 **Table 3. Quantification of spores of arbuscular mycorrhizal fungi in Cerrado Biome**
105 **soil, under sunflower cultivation, in two years and three periods, in Campo**
106 **Verde – MT, Brazil**

Year	Genotype	Sowing	Flowering	Harvest	Average
		n° spores 50 g soil ⁻¹			
2009	M 734	153 bB	296 aA	267 aA	247 a
	Agrobel 960	185 abB	342 aA	233 abB	
	Helio 358	262 abAB	311 aA	174 aB	
	Average	200 B	317 A	225 B	
2010	M 734	234 abAB	270 aA	147 bB	232 a
	Embrapa 122	191 abA	254 aA	216 abA	
	HLA 860 H.O.	271 aAB	320 aA	184 abB	
	Average	232 AB	281 A	182 B	
	CV (%)		11,60		

107 Means followed by different letters, uppercase in line and lowercase in column, differ from each other,
108 by the Tukey test ($P=0.05$). CV: coefficient of variation.

109 For the periods, Spore density at the flowering stage was higher in the two over both
110 cropping years of study, with on average 317 and 281 in a general average of 317 in 2009
111 and 281 in 2010, respectively (Table 3). The authors cited in the reference [10] confirm that
112 the maximum spore production can occur at the flowering and period and in the final
113 growth stages of the host crop as reported by Smith and Read [10]. They also reported that
114 possibly this increase in the amount of spores may be related to the higher production
115 of internal crop resistance structures in response to drought period of lower rainfall
116 (situation of stress).

117 According to the authors cited in the reference [2], the spore density of AMFs is generally
118 higher in agricultural systems, and variations may occur due to edaphoclimatic factors,
119 growing time, agricultural practices as well as the implanted crop.

120 The authors cited in the reference [11] studying Cerrado biome verified that the arbuscular
121 mycorrhizal fungi contribute to the growth of cultivated plants in annual cropping and pasture
122 systems and the number of spores of the native fungi varies, being the crop and the
123 cultivation system determinant for the enrichment of mycorrhizal fauna.

124 The interaction between the genetic factors and the period was significant, demonstrating
125 that the genetic material influences the sporulation process. However, the variations were
126 low indicating the stabilization of the mycorrhizal fungi sporulation.

127 In a carried study was verified that spore densities vary from 301 to 608 for maize crop,
128 whereas in soybean cultivated soil the values were between 239 and 287 [12], similar to
129 those obtained in the present work with sunflower. Mycorrhizal dynamics involving root
130 colonization and sporulation occur in different ways in different crops due to the compatibility
131 between AMF and the genetic characteristics of plants [13]. In addition, environmental,
132 climatic and edaphic factors generate changes in the symbiotic process [14].

133 In sugarcane the occurrence of AMF increase when the crop was preceded by sunflower
134 [15]. Likewise, sunflower favored the inoculum potential of AMF in the soil, and subsequent
135 corn growth [16]. Annual crops, green manures and forage species have a high degree of
136 mycorrhizal dependency, acting as a soil conditioning, multiplying the native mycorrhizal
137 community [17,14]. In this sense, sunflower is an option to benefit the soil mycorrhizal
138 population in crop rotation / succession systems.


Formatted: Highlight

139 For the mycorrhizal colonization rate, it was observed a variation from 21 to 28% in 2009
 140 and 28 to 48% in 2010 (Table 4) (Figure 1), with no difference between genotypes.
 141 According to the authors cited in the reference [18], mycorrhizal dependence can be defined
 142 as the plant's responsiveness to mycorrhization through increased growth, which may be
 143 related to the fertility and amount of phosphorus, present in the soil.

144 **Table 4. Average percentage of AMF colonization in soil under sunflower cultivation,**
 145 **in Campo Verde - MT, Brazil, in 2009 and 2010**

Year	Genotype	Mycorrhizal colonization (%)	Average
2009	M 734	28 a	24 b
	Agrobel 960	21 a	
	Helio 358	22 a	
2010	M 734	38 a	38 a
	Embrapa 122	48 a	
	HLA 860 H.O.	28 a	
CV (%)		16,24	

146 Means followed by different letters in the column differ from each other, by the Tukey test ($P = .05$).
 147 CV:coefficient of variation.



148 **Figure 1. Sunflower root colonization by arbuscular mycorrhizal fungi (AMF). Fungal**
 149 **structures: arbuscules (a) and vesicles (b).**

Formatted: Highlight

150 The authors cited in the reference [19] found colonization of AMF between 11 and 54% for
 151 arboreal species, for different crop rotation/succession systems, around 33 and 49% [20], as
 152 well as associated with crop of cassava with 31 to 71% in different localities [21] and in
 153 banana plant varying about 40 a 75% [22].

154 About the factor year, in 2010 there was a higher mycorrhizal colonization, which can be
 155 explained by the lower phosphorus content in the soil (Table 1). The effect of increase in
 156 phosphorus availability and decrease in symbiosis plant-mycorrhiza is negative [23] and
 157 emphasized in the literature [24] so, the reduction in the P content may lead to an increase
 158 in plant colonization. Evaluating different doses of P_2O_5 in the mycorrhiza colonization, it was
 159 found that doses greater than $30 \text{ mg P kg solo}^{-1}$ decrease colonization in sunflower [25].

160 Studying sunflower hybrids, it was verified that higher doses of P decreased sporulation and
 161 AMF colonization [22]. In the same crop, another study reported colonization percentage
 162 around 66 to 71% and spore density about 155 to 294, however, the soil had lower
 163 phosphorous content, if compared with the present work [26]. Contrasting the results cited in
 164 the reference [26] and this research, a lower colonization was capable to produce similar or
 165 higher quantities of spores.

176 Comparing the results cited in the reference [27] and this research, it was observed that a
177 lower colonization was able to produce similar or higher quantities of spores, so, the
178 efficiency in mycorrhiza species perpetuation was superior.

179 In general, the relationship AMF-plant can be mediated by nutrient levels, present in the soil,
180 since these fungi increase root exploration area, contributing to a greater absorption of
181 nutrients for the plant. As the increase in soil phosphorus decreases the root mycorrhizal
182 colonization and the plant dependence to mycorrhization [28], in soils with low levels of
183 phosphorus, typical of the Cerrado biome, the AMF favors sunflower cultivation [25].

184 In addition, there is evidences that mycorrhizal-sunflower ratio enables greater plant
185 resistance to heat, showing an interesting impact in Cerrado production systems, which is
186 characterized by high temperatures [29].

187 Moreover, the potential of AMFs as biofertilizer for oleaginous crops is reforced, especially
188 for soils with low fertility, since the practice allows to reach adequate levels of production,
189 with less use of synthetic fertilizers making the productive system more sustainable [26].

190 Therefore, colonization and mycorrhizal sporulation vary according to the sunflower
191 genotype and the evaluation period. On flowering period there were intense AMFs activity,
192 moment that is required to the plant a high nutritional supply for grain production.

193 4. CONCLUSION

194 The number of spores varies from 153 to 342 in 2009 and 147 to 320 in 2010, and the
195 maximum production occurs, in average, in the flowering period.

196 Mycorrhizal colonization in sunflower is not influenced by the genotype and the average
197 percentage was 24 (2009) and 38 (2010).

198 Lower soil phosphorus levels favors arbuscular mycorrhizal colonization.

199 COMPETING INTERESTS

200 We declare that no competing interests exist.
201

202 REFERENCES

203 1. Silva TFB, Santos ABS, Rozas CEO, Santos AC, Paiva LM. Influence of the density of
204 mycorrhizal fungi on the production of passion fruit (*Passiflora alata* CURTIS). The 2009; 22
205 (4): 1-6. English.

206 2. Rosseto P, Urcoviche RC, Oliviera JR, Alberton O. Spore density of mycorrhizal and
207 fungal fungi of the UNIPAR glomale germplasm bank. Archives of Veterinary Sciences and
208 Zoology of UNIPAR. 2012; 15 (1): 43-47. English. DOI: 10.25110 / arqvet.v15i1.2012.4166

209 3. Silva AJN. Soil chemical properties and growth of sunflower (*Helianthus annuus* L.) as
210 affected by the application of organic fertilizers and inoculation with arbuscular mycorrhizal
211 fungi. Revista Brasileira de Ciência do Solo. 2015; 39: 151-161. DOI:
212 10.1590/01000683rbc20150194
213
214

Formatted: Highlight

- 215 4. Grunvald AK, Carvalho CGP, Oliveira ACB, Andrade CAB. Adaptability and stability of
216 sunflower genotypes in Central Brazil. *Pesquisa Agropecuária Brasileira*. 2008; 43 (11):
217 1483-1493. English. DOI: 10.1590 / S0100-204X2008001100006
- 218 5. Souza FR, Silva IM, Pellin DMP, Bergamin AC, Silva RP. Agronomic characteristics of the
219 sunflower crop intercropped with *Brachiaria ruziziensis*. *Agronomic Science Journal*. 2015;
220 46 (1): 110-116. English. DOI: 10.1590 / S1806-66902015000100013
- 221 6. Gerdemann JW, Nicolson TH. Spores of mycorrhizal Endogone species extracted from
222 soil by wet sieving and decanting. *Transaction of the British Mycological Society*. 1963; 46
223 (2): 235-244. English. DOI: 10.1016 / S0007-1536 (63) 80079-0
224
- 225 7. Giovanetti M, Mosse B. An evaluation of techniques for measuring vesicular arbuscular
226 mycorrhizal infection in roots. *New Phytologist*. 1980;84(3):489-500. DOI: 10.1111/j.1469-
227 8137.1980.tb04556.x
228
- 229 8. Phillips JM, Hayman DS. Improved procedures for clearing roots and staining parasitic
230 and vesicular arbuscular mycorrhizal fungi for rapid assessment of infection. *Transaction of*
231 *the British Mycological Society*. 1970;55(1):158-161. DOI: 10.1016/S0007-1536(70)80110-3
232
- 233 9. Carrenho R et al. Mycorrhizal fungi in Brazilian agrosystems. In: Siqueira JO et al.
234 *Mycorrhizae: 30 years of research in Brazil*. Lavras: UFLA; 2008.
- 235 10. Smith SE, Read DJ. *Mycorrhizal symbiosis*. San Diego: Academic Press; 1997.
- 236 11. Miranda JCC, Vilela L, Miranda LN. Dynamics and contribution of mycorrhizal fungi in
237 crop rotation systems. *Pesquisa Agropecuária Brasileira*. 2005; 40 (10): 1005-1014. DOI:
238 10.1590 / S0100-204X2005001000009
- 239 12. Angelini GAR, Loss A, Pereira MG, Torres JLR, Saggin Júnior OJ. Mycorrhizal
240 colonization, spore density and diversity of mycorrhizal and fungal fungi in Cerrado soil
241 under no - tillage and conventional tillage. *Agrarian Sciences*. 2012; 33 (1): 115-130.
242 English. DOI: 10.5433 / 1679-0359.2012v33n1p115.
- 243 13. Smith SE, Gianinazzi-Pearson V. Physiological interactions between symbionts in
244 vesicular-arbuscular mycorrhizal plants. *Annual Review of Plant Physiology and Plant*
245 *Molecular Biology*. 1988; 39 (1): 221-244. DOI: 10.1146 / annurev.pp.39.060188.001253
- 246 14. Miranda JCC. *Closed: mycorrhizae: occurrence and management*. Planaltina: Embrapa
247 *Closed*; 2008.
- 248 15. Ambrosano EJ et al. Crop rotation biomass and arbuscular mycorrhizal fungi effects on
249 sugarcane yield. *Scientia Agricola*. 2010; 67 (6): 692-701. DOI: 10.1590 / S0103-
250 90162010000600011
- 251 16. Karasawa T, Kasahara Y, Takebe M. Differences in growth responses of maize to
252 preceding cropping caused by fluctuation in the population of indigenous arbuscular
253 mycorrhizal fungi. *Soil Biology and Biochemistry*. 2002; 34 (6): 851-857. DOI: 10.1016 /
254 S0038-0717 (02) 00017-2
- 255 17. Miranda JCC, Miranda LN, Vilela L, Vargas MA, Carvalho AM. Management of
256 mycorrhizal fungi through crop rotation in the agricultural systems of the Cerrado. *Technical*
257 *Communiqué* 42, Embrapa Cerrados. 2001; 42: 1-3. English. Accessed 16 Feb 2019.

- 258 18. Janos DP. Plant responsiveness to mycorrhizas differs from dependence upon
259 mycorrhizas. *Mycorrhiza*. 2007; 17 (2): 75-91. DOI: 10.1007 / s00572-006-0094-1
- 260 19. Silva LX, Figueiredo MVB, Silva GA, Goto BT, Oliveira JP, Burity HA. Fungos
261 micorrízicos arbusculares em áreas de plantio de leucena e sábia no estado de
262 Pernambuco. *Revista Árvore*. 2007; 31 (3): 427-435. Portuguese. DOI: 10.1590/S0100-
263 67622007000300008
- 264
265 20. Cordeiro MAS, Carneiro MAC, Paulino HB, Saggin Junior OJ. Colonização e densidade
266 de esporos de fungos micorrízicos em dois solos do cerrado sob diferentes sistemas de
267 manejo. *Pesquisa Agropecuária Tropical*. 2005; 35 (3): 147-153. Portuguese.
- 268 21. Balota EL, Lopes ES, Hungria M, Döbereiner J. Ocorrência de bactérias diazotróficas e
269 fungos micorrízicos arbusculares na cultura da mandioca. *Pesquisa Agropecuária Brasileira*.
270 1999; 34 (7): 1265-1276. Portuguese.
- 271 22. Oliveira NA, Oliveira LA, Figueiredo AF. Colonização micorrízica e concentração de
272 nutrientes em três cultivares de bananeiras em um latossolo amarelo da Amazônia Central.
273 *Acta Amazônica*. 2003; 33 (3): 345-352. Portuguese.
274
- 275 23. Ferreira DA, Carneiro MAC, Saggin Júnior OJ. Mycorrhizal fungi in a red latosol under
276 management and uses in the Cerrado. *Soil Science Journal*. 2012; 36 (1): 51-61. English.
277 DOI: 10.1590 / S0100-06832012000100006
- 278 24. Silva AJN, Silva RA, Santos JS, Medeiros JCS, Carvalho FG, Silva VN, Oliveira CJ,
279 Araújo AC, Silva LESF, Gomes-Júnior J. Soil chemical properties and growth of sunflower
280 (*Helianthus annuus* L.) as affected by the application of organic fertilizers and inoculation with
281 arbuscular mycorrhizal fungi. *Brazilian Journal of Soil Science*. 2015; 39 (1): 151-161.
282 English. DOI: 10.1590/01000683rbc20150194
- 283 25. Balota EL, Machineski O, Truber PV, Cerezini P, Milani KL, Scherer A, Honda C, Leite L.
284 Efeito dos fungos micorrízicos arbusculares em culturas oleaginosas. *Brazilian Congress of*
285 *Mamona, 4 & International Symposium of Oleaginose Energy*, 1. 2010;680-684. Accessed
286 22 April 2019. Available: [https://ainfo.cnptia.embrapa.br/digital/bitstream/item/18359/1/FER-
287 06.pdf](https://ainfo.cnptia.embrapa.br/digital/bitstream/item/18359/1/FER-06.pdf).
- 288 26. Sarah S, Ibrar M. Effects of Arbuscular Mycorrhizal Fungi on Spores Density and Root
289 Colonization of Four Hybrids of Sunflower (*Helianthus annuus* L.) at Different Rock
290 Phosphate Levels. *Sarhad Journal of Agriculture*. 2016; 32 (4): 258-266. DOI:
291 10.17582/journal.sja/2016.32.4.258.266
- 292 27. Alves TS, Campos LL, Borba Filho AB, Loureiro MF. Ocorrência de fungos micorrízicos
293 arbusculares na cultura de girassol no município de Campo Verde/MT. *Scientia Agraria*
294 *Paranaensis*. 2014; 13 (3): 243-250. DOI: 10.18188/1983-1471/sap.v13n3p243-250
- 295 28. Balota EL et al. Effect of mycorrhizal fungi on different doses of phosphorus in sunflower
296 and peanut. *Encyclopedia Biosphere*. 2010; 6 (11): 1-8. English.
- 297 29. Mayer Z, Duc NH, Sasvári Z, Posta K. How arbuscular mycorrhizal fungi influence the
298 defense system of sunflower during different abiotic stresses. *Acta Biologica Hungarica*.
299 2017; 68 (4): 376-387. DOI: 10.1556/018.68.2017.4.4
300

