

OPHTHALMIC PROBLEMS OF ADULTS IN RURAL COMMUNITIES OF RIVERS STATE, NIGERIA

ABSTRACT

Aim: To determine the ophthalmic problems and their possible causes among adults in rural communities in Rivers State.

Methods: A multistage population based random sampling study of adults in five clans of Etche Local Government Area of Rivers State was undertaken. Medical history was taken and comprehensive ocular examination done on each subject. Ocular examination included visual acuity, visual field, tonometry and ophthalmoscopy. Data taken were recorded and analysed using statistical software called Minitab 11. Ethical approval was obtained from relevant authorities.

Results: Out of the 600 subjects seen in this study, 276 (46.0%) were males and 324 (54.0%) females. They were all above 21 years old. 26 (2.4%) subjects had good vision, while 8 (0.8%) were blind. The ophthalmic problems identified were Presbyopia 298 (28.0%), Refractive error 247 (23.2%), Cataract 126 (11.8%), Allergic conjunctivitis 106 (9.9%), Glaucoma 94 (8.8%), Pterygium 86 (8.1%), Bacterial conjunctivitis 35 (3.3%), Corneal opacity 32 (3.0%), Chalazion 4 (0.4%), Diabetic retinopathy 3 (0.3%) and Ptosis 1 (0.1%).

Conclusion: The most common ophthalmic problems in this study which were dependent on age, gender and occupation include Presbyopia, Refractive Error, Cataract and Allergic Conjunctivitis. More females and farmers were seen in this study. The problems identified could easily be managed if well-equipped health facilities were provided by the government.

Keywords: Ophthalmic problems, adults, rural communities, Rivers State.

INTRODUCTION

Ophthalmic problems are global and constitute serious public health challenges especially among older adults [1]. According to Bethesda, the prevalence of blindness and visual impairment increases with age among all racial and ethnic groups, especially among people older than 75 years of age [2].

The World Health Organization's estimated number of people with visual impairment worldwide is 285 million, while 39 million are blind and 246 have low vision [3]. About 81% of all people who are blind or have moderate to severe visual impairment are aged 50 years and above, indicating that with an increasing population of older people, more people will be at risk of visual impairment due to chronic eye diseases [4]. About 90% of the world's visually impaired live in low income settings and 80% of all visual impairment can be prevented or cured and over 90% of the world blind are in Sub Saharan African and Asia and especially among the persons in the rural communities [5]. Lawallen and Courtright reported the major causes of blindness in Africa as cataract, trachoma and glaucoma [6]. Blindness prevalence rates vary globally but evidence based study suggests that approximately 1% of Africans are blind and majority of the blindness in that region are preventable or curable [6].

The Nigeria National blindness and visual impairment survey that was carried out in the year 2009 [7] showed that the major causes of blindness and visual impairment among adults in Nigeria were uncorrected refractive error, cataract and glaucoma. The survey also stated that increasing age was associated with increasing prevalence of all blinding conditions. According to the survey, 4.25 million adults aged 40 years and above have moderate to severe visual impairment or blindness. The prevalence of blindness in Nigeria is 0.78%, attributed to poor technology, minimal eye care services, malnutrition and poverty [7,8].

The commonest causes of blindness worldwide are cataract, glaucoma, trachoma, onchocerciasis and refractive errors [9]. Most of these blinding diseases are preventable and easily treatable, but the majority of the victims in Africa and Asia are either poor, ignorant, or do not have eye -care services available to them [10]. Etche indigenes are predominantly farmers and farmers according to Momoh and Abadom are

usually exposed to certain occupational hazards that predispose them to ocular diseases and injuries [11]. Visual impairment obviously compromises people's quality of life because it makes them unable to read, watch television, drive cars, operate machines or attend to themselves. Most times, it isolates older people from friends and family which may lead to depression.

Ejimadu and Pedro-Egbe[12] in their study on prevalence and causes of Blindness in Ikwerre Local Government Area of Rivers State, revealed that the three top causes of blindness in that community were cataract, Glaucoma, Optic Atrophy, Corneal Opacity, Phthisis Bulbi, Absent Globe, Chorioretinitis and Maculopathy. They further concluded that most of these blinding eye diseases were avoidable; therefore more emphasis on eye care should focus on prevention through public enlightenment and regular eye screening, with participation of the government. Also the prohibition of harmful traditional practices, discouragement of self-medication, provision of basic eye care delivery and increasing cataract surgery will reduce prevalence of blindness.

METHODOLOGY

A multistage, population based, random sampling study, of adults in five clans of Etche Local Government Area of Rivers State was undertaken.

Medical history was recorded and comprehensive ocular examination done on each of the 600 subjects (276 males and 324 females) who were at least 21 years, at the community health centre, after obtaining consent from them. Ocular examination included visual acuity, visual field, tonometry and ophthalmoscopy.

Instruments used during the research were Pen torch for examination of the external structures of the eyes, Keeler ophthalmoscopes for fundus examination, Snellen's charts both literate and illiterate charts for visual acuity assessment, Reichert AT 555 Auto non-contact tonometer for measurement of the intra-ocular pressure and trial lens cases used for subjective refraction. Subjects requiring visual field analysis were referred to the tertiary hospital in the state.

Data obtained were analysed using statistical software called Minitab 11 where the raw data obtained were classified into different groups and categories based on their common characteristics. The data were

logically represented, where raw data were summarized and displayed in a compact form that is statistical tables.

An ethical approval to carry out the study was obtained from Rivers State Ministry of Health through the office of Planning, Research and Statistics. Afterwards a second approval was obtained from Rivers State Ethical Committee following due applications.

Inclusion criteria was adults in Etche local Government Area who were 21 years and above and was randomly selected at the sampling stage. It also involved those that signed the consent forms and were ready to participate.

RESULTS

Table 1 shows the demographical characteristics of the respondents. Out of the 600 subjects seen in this study 276 (46.0%) were males and 324 (54.0%) were females. Their ages ranged from 21 years and above. The highest age group was 41-50 with 174 (29.0%) subjects, followed by age group of 31-40 years 161 (26.8%) while the lowest age group was >60years with frequency of 38 (6.3%).

The second segment of the table shows the occupational distribution of the subjects. Majority were farmers; 276 (46.0%) while others were civil servants 152 (25.3%), and traders 102 (17.0%), few students 46 (7.6%), Retirees 15 (2.5%) and unemployed 9 (1.5%).

Table 2 summarizes the distribution of ophthalmic conditions of subjects. The most predominant oculo-visual condition was presbyopia (28.0%), followed by refractive error (23.1%) and cataract (11.8%). The least common conditions were chalazion (0.4%), diabetic retinopathy (0.3%) and ptosis (0.1%).

Table 3 shows the distribution of common ophthalmic problems with respect to occupation. Farmers (41.3%) presented more with cataract than other occupations. Civil Servants had more errors refractive (51.8%) than other groups. The highest prevalence of presbyopia occurred amongst Civil Servants (49.0%).

Table 4 shows gender related ocular conditions seen in the subjects. Females presented more with allergic conjunctivitis (7.1%), pterygium (4.1%) and cataract (6.0%) than males. While the males presented more with refractive error (11.9%) and presbyopia (14.8%) than females.

107 **Table 1: DEMOGRAPHIC CHARACTERISTICS OF THE RESPONDENTS**

108	AGE (YEARS)	MALE (%)	FEMALE (%)	FREQUENCY (%)
109	21-30	57 (9.5)	60 (10.0)	117 (19.5)
110	31-40	79 (13.1)	82 (13.6)	161 (26.8)
111	41-50	72 (12.0)	102 (17.0)	174 (29.0)
112	51-60	46 (7.6)	64 (10.6)	110 (18.3)
113	>60	22 (3.6)	16 (2.6)	38 (6.3)
114	TOTAL	276 (46.0)	324 (54.0)	600 (100)
115	OCCUPATION			
116	Civil Servants	81 (13.5)	71 (11.8)	152 (25.3)
117	Traders	52 (8.6)	50 (8.3)	102 (17.0)
118	Farmers	115 (19.1)	161 (26.8)	276 (46.0)
119	Students	17 (2.8)	29 (4.8)	46 (7.6)
120	Retirees	9 (1.5)	6 (1.0)	15 (2.5)
121	Unemployed	2 (0.3)	7 (1.1)	9 (1.5)
122	TOTAL	276 (46.0)	324 (54.0)	600 (100)

123

124 **TABLE 2: DISTRIBUTION OF OPHTHALMIC CONDITIONS OF SUBJECTS.**

125	OCULO/VISUAL STATUS	FREQUENCY (N)	(%)
126	Presbyopia	298	28.0
127	Refractive Error	247	23.1
128	Cataract	126	11.8
129	Allergic Conjunctivitis	106	9.9
130	Glaucoma	94	8.8

131	Pterygium	86	8.0
132	Bacterial Conjunctivitis	35	3.3
133	Corneal Opacity	32	3.0
134	Good Vision	26	2.4
135	Blindness	8	0.8
136	Chalazion	4	0.4
137	Diabetic Retinopathy	3	0.3
138	Ptosis	1	0.1
139	Total	1066	100

140

141 **Table 3: DISTRIBUTION OF COMMON OPHTHALMIC PROBLEMS WITH RESPECT TO**
142 **OCCUPATION.**

OCCUPATION	OCULO-VISUAL CONDITIONS (NO (%))				
	CATARACT	REFRACTIVE ERROR	PRESBYOPIA	ALLERGIC CONJUNCTIVITIS	PTERYGIUM
CIVIL SERVANT	19 (15.1%)	128(51.8%)	146(49.0%)	15(14.2%)	10(11.6%)
TRADERS	45 (35.7%)	22(8.9%)	50(16.8%)	18(17.0%)	16(18.6%)
FARMERS	52 (41.3%)	57 (23.1%)	69(23.2%)	55(51.9%)	53(61.6%)
STUDENTS	0 (0%)	31(12.6%)	10(3.3%)	12(11.3%)	3(3.5%)
RETIREEES	8 (6.3%)	6(2.4%)	15(5.0%)	3(2.8%)	3(3.5%)
UNEMPLOYED	2 (1.6%)	3(1.2%)	8(2.7%)	3(2.8%)	1(1.2%)

TOTAL	126(100%)	247 (100%)	298(100%)	106(100%)	86(100%)
-------	-----------	------------	-----------	-----------	----------

143

144 **Table 4: GENDER –RELATED OHTHALMIC CONDITIONS IN SUBJECTS**

145	Ocular conditions	Gender		No (%) Prevalence
		Male	Female	Total
147	Presbyopia	158 (14.8%)	140 (13.1%)	298 (28.0%)
148	Refractive error	127 (11.9%)	120 (11.3%)	247 (23.1%)
149	Cataract	62 (5.8%)	64 (6.0%)	126 (11.8%)
150	Allergic	30 (2.8%)	76 (7.1%)	106 (9.9%)
151	Conjunctivitis			
152	Glaucoma	46 (4.3%)	48 (4.5%)	94 (8.8%)
153	Pterygium	42 (3.9%)	44 (4.1%)	86 (8.0%)
154	Bacterial			
155	Conjunctivitis	17 (1.6 %)	18 (1.7%)	35 (3.3%)
156				
157	Corneal Opacity	21 (2.0%)	11 (1.0%)	32 (3.0%)
158	Good Vision	14 (1.3%)	12 (1.1 %)	26 (2.4%)
159	Blindness	5 (0.5%)	3 (0.3%)	8 (0.8%)
160	Chalazion	3 (0.3%)	1 (0.1%)	4 (0.4 %)
161	Diabetic			
162	Retinopathy	3 (0.3%)	0 (0%)	3 (0.3%)
163				
164	Ptois	0 (0%)	1 (0.1%)	1 (0.1%)

Total	528 (49.5%)	538 (50.5%)	1066 (100%)
--------------	--------------------	--------------------	--------------------

165

UNDER PEER REVIEW

DISCUSSION

The ophthalmic problems found among adults in Etche LGA were Presbyopia 298 (28.0%), Refractive Error 247 (23.1%), Cataract 126 (11.8%), Allergic Conjunctivitis 106 (9.9%), Glaucoma 94 (8.8%), Pterygium 86 (8.1%), Bacterial Conjunctivitis 35 (3.3%), Corneal Opacity 3 (3.0), Chalazion 4 (0.4%), Diabetic Retinopathy 3 (0.3%) and Ptosis 1 (0.1%). 26 (2.4%) subjects had good vision while 8 (0.8%) were blind. These findings were similar to studies by WHO [4] that listed the common ocular diseases worldwide as cataract, glaucoma, conjunctivitis, corneal ulcers, uveitis, refractive errors, pterygium, trachoma, onchocerciasis, xerophthalmia and ocular malignancies.

Presbyopia was the most common type of ophthalmic problems found in our study, accounting for 28.0% of all cases seen. This was similar to findings by Nwosu [14]

The second most prevalent ophthalmic problem was Refractive Error 247(23.1%). According to WHO,[4] uncorrected refractive errors were the most common causes of visual impairment accounting for 43% of cases and representing an important causes of blindness [15]. The prevalence of refractive errors in this study was higher in males than females, which may be linked to the fact that majority of the males were educated, in school or were civil servants which may be a contributory factor to the diagnosis of refractive error or presbyopia. This was in agreement with a study which stated that the prevalence of refractive errors vary with race, age, gender and geographical region and that environmental factors like level of education, occupation, near work load, time of outdoors as a child are also associated with aetiology of refractive error[16]. Uncorrected refractive error was the main cause of Low vision and second commonest cause of blindness in a study which also revealed that uncorrected refractive error can hamper performance at school, reduce employability and productivity, and generally impair quality of life [15].

Uncorrected refractive error which was the commonest cause of ophthalmic problems in this study has been reported as the commonest cause of ocular morbidity in another study [17]. It was the commonest cause of mild and moderate visual impairment in the Nigerian national blindness and visual impairment survey, accounting for 77.9% and 57.1% respectively [7,18].

We recorded cataract as the third most common type of ophthalmic problem, accounting for 11.8% of all cases which was similar to that of 16.7% reported from a study in Benin, Nigeria [13]. In a study on causes of visual impairment and blindness in Kwara State of Nigeria [19], Cataract was responsible for more than half the cases of ocular morbidity and was the commonest cause of visual disability. The high rate of cataract cases in the study was basically unknown, but may be attributed to their constant exposure to ultraviolet rays, firewood smoke, trauma, age group of the study area and poorly controlled diabetes, since a lot of the cataract patients reported to be diabetic.

The fourth commonest ocular problem reported in this study was Allergic Conjunctivitis with an incidence of 9.9%. This was similar to other studies that reported Allergic Conjunctivitis as the third leading cause of ocular morbidity with prevalence of less than 20% [14, 20].

The high occurrence of allergic conjunctivitis in this study may be associated with higher pollen content of the farming environment since they were basically farmers, this was related to a study by Momoh and Abadon¹¹ where high rate of allergic conjunctivitis found in farming environment was linked to higher pollen content of farming environment and also they postulated that allergic conjunctivitis may be prevalent in a dusty environment.

Glaucoma is one of the common ocular diseases found in this study accounting for 8.8%. This was similar to a study where Glaucoma was seen in 11.9% of patients [17] and it had been reported to be the second most common cause of blindness or visual impairment worldwide [5]. It is the leading cause of irreversible blindness in West Africa and it had been estimated that 20% of people older than age 40 in West Africa may be at risk from the disease [17].

Pterygium is another prevalent ocular disease in this population with an incidence of 8.1%. This is consistent with Momoh and Abadom [11] where incidence of pterygium was common among farmers but

in contrast in another study [20] that showed Pterygium as the second common eye disorder among the welders in their study with a prevalence of 17.5 %

Corneal Opacity accounted for 3.0% in this study. This may be attributed to the fact that the majority of the subjects were predominantly farmers and most of the subjects reported applying traditional medicine in the eyes. Majority of the corneal opacity occurred as a result of trauma and traditional medical practices. About 321(68.0%) respondents have never had any form of ocular trauma, while 151(32.0%) respondents reported of having at least one episode of ocular trauma but only 32(3.0%) subjects had corneal Opacity, this may have connection with the majority indigene of the study area being predominately farmers. This is in line with the global estimates that showed that there are about total of 1.6 million ocular trauma cases of blindness and about 2.3 million ocular trauma from agricultural labour, also victims have less access to eye care services than their urban counterparts, it is likely that rural people may have a greater burden of vision impairment or blindness caused by trauma[14].

In contrast with those of Wokoma and Ichenwo [10] in rural community in Rivers State, Nigeria where a lower occurrence of corneal opacities was reported (0.9%). The subjects being basically farmers had a high occurrence of trauma-related visual problems (corneal opacity) which may be attributed to the fact that they came directly in contact with occupational hazards such as dust, projectiles of organic agricultural materials such as twigs, seeds and falling objects.

Surprisingly bacterial conjunctivitis (3.3%) showed to be an uncommon ocular problem in this study. This was similar to the study by Momoh and Abadom¹¹ with incidence of 1.3%. Other rare ocular diseases found in this study include chalazion 0.4%, diabetic retinopathy 0.3%, ptosis 0.1% and blindness 0.8%.

The distribution of blindness in this study showed that six subjects (75.0%) had mono-ocular blindness while two subjects (25.0%) were bilaterally blind. The three causes of blindness in the subjects were Glaucoma (25%), Cataract (50%) and Corneal Opacity / Trauma (25%). The incidence of blindness (0.8%) may suggest poor or no availability of eye care services in the locality.

This study revealed significant relationship between the subjects' occupations and their common oculo-visual problems. The majority of the subjects were mainly farmers 276 (46.0%), civil servants 152(25.3%) and traders 102 (17.0%). Civil Servants 128 (51.8%) and Students 31(12.6%) had the highest prevalence on Refractive error/Presbyopia respectively. This may be attributed to their visual task being higher than those in other occupations. This was similar to a study by Njebuome, Onyebuchi, and Igbe²¹ that showed the pattern of oculo-visual problems among public / civil servants in Abuja as follows: refractive error 88.7%, Cataract 1.1%, Pterygium 2.3%, Disc cupping 3.4%, Chalazion 1.1% and Conjunctivitis 3.4%, where the ages of the subjects ranged from 25 years to 60 years. The study showed refractive error as a leading cause of visual impairment among civil servants in Abuja. Farmers were found to have the highest prevalence of Allergic Conjunctivitis (51.9%) and Pterygium (61.6%). This may also be attributed to the nature of their occupation that was basically outdoor activities that expose them to dust and ultra violet rays.

More so, the common ocular diseases prevalent among adults in our study were dependent on gender. The adult females had the highest prevalence on Cataract 64 (50.8%), Allergic Conjunctivitis 76 (71.7%) and pterygium 44 (51.2%) while Refractive Error 127 (51.4%) /Presbyopia 158 (53.0%) were more prevalent in males. This may be associated with the fact that majority of their females were more exposed to farm related activities while the males mostly did official works hence, had higher near visual tasks. This was in contrast to similar studies in the same environment and in southern Nigeria where there were a higher proportion of males to females and the male had a higher prevalence of Pterygium and allergic conjunctivitis in the study by Edema and Okojie¹³. But this finding was similar to a study by Nwosu¹⁴ on rural young adults in Anambra state whose predominant occupation was farming, in which there were more females than males in the study and they had higher prevalence of allergic conjunctivitis than males. Nwosu (1998) postulated that it was probably due to the rural- urban drift of more males than females. It was also similar to a study by Wokoma¹⁰ in a rural community in Rivers State where the proportion of

female participants was higher than that of male and they also presented with higher rate of allergic conjunctivitis. The absence of any form of eye care service in this community, no doubts contributed to the relatively high prevalence of visual impairment. Eye diseases that would have been detected earlier and intervention given, continued to persist and deteriorated, eventually progressing to blindness. None of the General hospitals in our study area had any form of eye service. The available state owned hospitals that had eye sections were at Port Harcourt, Okrika, Ahoada and Bori. Unfortunately, the distance from our study area to these facilities, the logistics and costs involved hindered majority from accessing quality eye services. The greater majority remained in the community with their problem until they became blind. The observation in this rural community is not peculiar to Etche as similar observations have been reported in other rural communities in Nigeria [10,22]. The causes of blindness in this study were preventable and treatable if detected early.

CONCLUSION

The most common ophthalmic problems among adults in this study were Presbyopia, Refractive Error, Cataract, Allergic Conjunctivitis and Pterygium and they accounted for more than two-third of the ocular problems and were dependent on age, gender and occupation.

The lack of regular health education, inaccessibility of health facilities and the nature of their occupation may have been a contributing factor to the ocular diseases found in this study.

Also most of the subjects were predominantly farmers or were combining their occupations with farming and this may have exposed them to trauma, foreign body, dust or ultra violet rays that may have posed ocular problems. Regular screening, eye check and treatment of common eye diseases were highly recommended. The need to wear protective eye devices such as goggles could reduce exposure to ultraviolet radiation and offer protection against ocular injury.

The state Government should make eye care services available.

REFERENCES

1. Saadine JB, Venkat KM, Vinicor F. Vision Loss: A Public Health Problem. *Ophthalmology* 10(2):253-254
2. Bethesda MD. The vision problems in US: Prevalence of Adult vision Impairment and Age-Related Eye Diseases in America. Prevent Blindness America National Eye Institute of Health, 1-11.
3. World Health Organization. Available data on Blindness update 1989, 14; 1-23.
4. World Health Organization. Visual Impairment and Blindness. Fact Sheet N°282. Updated August 2014. http://www.who.int/mediacentre/fact_sheets/fs282/en/
5. World Health Organization. State of the World's sight: Vision 2020: the right to sight: 1999-2005. WHO library cataloguing in publication Data, 1-8.
6. Lawallen S, Courtright P. Blindness in Africa; Present Situation and future needs. *British Journal of Ophthalmology*, 85:897-903.
7. Abdul MM, Sivasubramaniam S, Murthy GVS, Gilbert C, Abubakar T, Ezelum C, Rabui MM. Causes of Blindness and Visual Impairment in Nigeria: The Nigeria National Blindness and Visual Impairment Survey. *Investigative Ophthalmology and Visual Science*, 50: 4114-4120.
8. Chukwuka IO, Pedro-Egbe CN, Onua AA. Ocular Problems Among Public Service Retirees in Southern Nigeria Metropolitan. *Nigerian Journal Of Ophthalmology*, 24:16-19.
9. Roodhooft MJ. Leading Causes of Blindness Worldwide. *Bulletin of the Belgian Society of Ophthalmology*, 283:19-25.
10. Wokoma FS, Ichenwo T. Pattern of Eye disorder in Ogbodo: A rural Community in Rivers State. *The Nigerian Health Journal*, 11(1):1
11. Momoh RO, Abadom EG. Pattern and Prevalence of Eye Diseases among Farmers in Agricultural Industry in Southern Nigeria. *Journal of Biomedical Science*, 14(2):164-166.
12. Ejimadu CS, Pedro-Egbe CN. Prevalence and Causes of Blindness in Ikwere Local Government Area of Rivers State, Nigeria. *The Nigeria Health Journal*, 9:1-4.
13. Edema OT, Okojie OH. Pattern of Eye Diseases cases in Benin city, Nigeria. *Journal of Medical Practice*, 4:86-90
14. Nwosu SNN. Ocular problems of young adults in rural Nigeria. *International Ophthalmology*, 22: 259-263.
15. Pascolini D, Marriotti SP. Global Estimate of Visual Impairment in 2010. *British Journal of Ophthalmology*, 96, 614-618.

- 317 16.Pan CW, Wong TY, Lavanya R, Wu RY. Zheng YF, Lin XY, Mitchell P, Aung T, Saw. SM.
318 Prevalence and risk factors for refractive errors in Indians: the Singapore Indian Eye Study. Invest
319 Ophthalmology Visual Science, 16,52(6): 3166-73.
- 320 17.Akinsola FB, Majekodunmi AA, Obowu CB, Ekanem EE. Pattern of Eye Diseases in Adults 16 years
321 and above in Alimoso Local Government Areas of Lagos State. Nigerian Postgraduate Medical Journal
322 2:56–61.
- 323 18.Ukponmwan CU. Pattern of Ocular Morbidity in Nigeria. Asian Pacific Journal of Tropical Diseases,
324 3(2):164-166.
- 325 19.Mahmoud AO, Olatunji FO, Buari SB. Survey of Blindness and
326 Ocular Morbidity in Kwara State, Nigeria. Nigerian Journal Surgical Science, 15:26–30.
- 327 20.Adeoye AO, Omotoye OJ. Eye disease in Wesley Guild Hospital, Ilesa, Nigeria. African Journal of
328 Medicine and Medical Sciences, 36 (4): 377-380.
- 329 21.Njepuome N, Onyebuchi U, Onwusoro M, Igbe, M. Visual Impairment Among Public Servants in
330 Nigeria. The internet Journal of Ophthalmology and Visual Science, 9:1.
- 331 22.Adegbehingbe BO, Majengbassan T O. Ocular Health Status of rural dwellers in South-Western
332 Nigeria. Australian Journal of Rural Health, 15: 269-272.