



23 available N and P. The results suggest that cowpea responds differently to Zinc sulphate  
24 application depending on its rates.

25 Keywords: N-fixed, Nodulation, Zinc fertilizer, NPK uptake and yield

## 26 INTRODUCTION

27 Proper nutrition of plants with micronutrients depends on various factors, such as the rate of  
28 absorption of nutrients by the plants, distribution of nutrients to functional sites and nutrient  
29 mobility within the plant. Interactions occur between the micronutrients and some nutrients  
30 [1, 2, 3]. The amount of nitrogen fixed is usually high in soils with low mineral N but with  
31 sufficient water and enough of other nutrients capable of supporting plant growth [4].  
32 Another factor is the differential response of plants to one nutrient in combination with  
33 varying levels of a second element applied simultaneously i.e. the two elements combine to  
34 produce an added effect not due to each of them acting alone [1, 2]. Such interactions may  
35 take place in the soil and within the plant [3]. However, the amount of nutrients uptake is  
36 strongly dependent on nutritional and environmental factors.

37 Cowpea is especially important for dry savannah of West Africa between latitudes 7 and  
38 14°N [5] and second after groundnut as the most important legume of Ghana in terms of  
39 space under cultivation (156,000 ha) and quantity produced and consumed annually (143,000  
40 Mg) making Ghana among the largest cowpea producer in Africa [6]. Cowpea is a protein-  
41 rich component of an otherwise protein-poor diet [7]. Many researchers have observed that  
42 Zn have a positive relationship with the nitrogen metabolism pathway of plants, its deficiency  
43 cause a reduction in protein synthesis into the plants. [8] identified the positive relationship  
44 between the flowering and fruiting process and Zn. As micronutrient, Zinc has received much  
45 recent attention [9] because it is present in all body tissues and fluids [10].

46 The native rhizobia are often low in numbers or ineffective and are therefore not able to fix  
47 enough nitrogen to meet the nitrogen demand of plants. The study was undertaken to examine  
48 the dynamics mineral contents in grain and haulm tissues and nutritional benefits following  
49 by zinc fertilizer application. The nodule parameter was also under investigation.

## 50 MATERIALS AND METHODS

51  The study was conducted at the Plantation Section of the Department of Crop and Soil  
52 Sciences, Faculty of Agriculture, KNUST, in the cropping seasons of 2016. The site is  
53 located at 06° 45' N and 01° 31' W in the rainforest belt of Ghana. The site was located at  
54 06° 45' N and 01° 31' W in the rainforest belt of Ghana. The total nitrogen content was low  
55 with a mean value of 0.06%, available P content was low with value of 6.4 mg/kg, soil Zn  
56 content was moderately low, found to be 1.290 mg/kg. Three early maturing cowpea  
57 varieties (Asontem, Agyenkwa and Zamzam) were grown in both experiments and Zn   
58 fertilizer rates of 0, 5 and 10 kg/ha were applied into the varieties. The Zn fertilizer was  
59 applied as foliar application in both experiments. The split plot design, arranged in RCBD  
60 was used for both cropping seasons. All recommended cultural practices were done in  
61 schedule. Cowpea varieties were obtained from the Crops Research (CSIR) at Fumesua,  
62 Kumasi/Ghana. Zinc sulfate heptahydrate was applied at 3 weeks (40%) and 5 weeks (60%)  
63 after sowing. The application was done early morning before 9:00 am, using a sprayer. The  
64 plots were demarcated three days after harrowing and seeds were sown by hand using manual  
65 labour. Seeds were sown at a spacing of 60 cm x 20 cm with a rate of two seeds per hill at the  
66 depth of 3-5 cm. Urea and triple superphosphate (TSP) fertilizers were applied as band  
67 placement by making a furrow of 5-7 cm deep and covering with 2 cm of soil. As starter  
68 nitrogen, Urea was applied at the rate of 20 kg N/ha uniformly to all plots at two weeks after  
69 sowing (WAS). Triple super phosphate (TSP) was also applied two weeks after sowing

70 (WAS) to the cowpea plant at the rate of 40 kg P<sub>2</sub>O<sub>5</sub>/ha. Standard agronomic and plant  
71 protection treatments were used uniformly across the plots for the duration of the experiment.  
72 Grass hoppers (*Empoasca kerri Pruth*), Thrips (*Caliothrips indicus Bagnall*) and Aphids  
73 (*Aphis craccivora Koch*) were pests, respectively at vegetative stage and flowering to the end  
74 of pod filling. Lambda master 2.5 % E.C. [Active ingredients (Lambda-Cyhalothrin, 9.8 %)]  
75 was the pesticides used for pests' control.

76 A random sample of five plants from each plot and a random sample of five pods from each  
77 of the five plants were selected to measure. Plant height, Stem girth and number of leaves  
78 were measured at 30, 45 and 60 days after sowing (DAS) and mean for each plot was  
79 calculated. Nodules were sampled at 30 and 45 days after sowing. Plants samples were  
80 uprooted gently washed with water and the total nodules counted and the mean calculated for  
81 each plot. The ground was sufficiently soaked with water 48 hours before sampling to each  
82 uprooting of plants. To determine nodule effectiveness, nodules were cut open using a razor  
83 blade and hand lens. Nodules with pink or reddish colour were considered effective and  
84 fixing nitrogen, while those with green or colourless appearance were recorded as ineffective  
85 nodules. Nodules per plot were kept in labelled envelopes and sent to the laboratory to oven-  
86 dry at 70°C for 48 hours. Average dry weight of nodules per plant was computed and  
87 expressed in grams. For mineral content analysis, random samples of five plants were  
88 uprooted gently from each plot at harvest and the root system was removed. The above  
89 ground parts were put in labelled envelopes and oven dry at 70° C for 72 hours and milled and  
90 one hundred gram samples of each of the plant part (seeds and haulms) were taken to  
91 determine nitrogen, phosphorus and potassium content. The nitrogen content was determined  
92 using the Kjeldahl method [11]. The protein content of seed was determined on the basis of  
93 total nitrogen content [12]. N-fixed was obtained using the N-difference method. The total


94 nitrogen content of the maize (reference crop) was subtracted from that of the cowpea [13]. In  
95 this study Omankwa maize variety was the reference crop. Phosphorus (P), the content was  
96 measured on the Spectronic 20 spectrophotometer to give absorbance measurements at a  
97 wavelength of 420 nm. The observed absorbance was used to determine the P content from  
98 the standard curve [14, 15] and Potassium (K) was obtained using the flame photometer.  
99 From the standard curve, the concentration of K was calculated using the particular  
100 absorbance observed for the sample. NPK uptake were done by multiplying the grain and  
101 haulm yield in kilograms per hectare by each analysed parameters separately, nitrogen,  
102 phosphorus and potassium, in the grain and haulm then divided by 100 percent. Zn content  
103 was determined using Perkins model 403 atomic absorption spectrophotometer after  
104 digestion. The file for the type of analysis and hollow cathode lamps were selected with  
105 appropriate wavelengths of 213.9 nm [16]. The grain and straw yields were recorded  
106 separately. Total Zn uptakes by grain and tissue were computed by multiplying Zn content  
107 and their respective dry weights/ha. Data collected were subjected to analysis of variance  
108 (ANOVA) using GenStat statistical package version 15<sup>th</sup>. The LSD test was used to compare  
109 treatment means at 5% probability.

## 110 RESULTS


### 111 1. **Effects of cowpea varieties** on growth




112 Fig 1 illustrates the effect of different cowpea varieties on plant height, stem girth and  
113 number of leaves over the period of the experiment. The significant effect at 5% level of  
114 probability of cowpea varieties used was recorded over all sampling period of the study. The  
115 tallest plant was obtained by Asontem variety and the lowest by Zamzam. However, cowpea  
116 varieties did not show any significant ( $P > 0.05$ ) effect on stem girth and number of leaves.  
117 Additionally, variety by Zinc rates was not significantly different on all days of sampling.



118



119




120


121 **Figure 1. Effects of cowpea varieties on Plant height (a and b), stem girth (c and d) and**  
 122 **number of leaves (e and f) under Zinc foliar application in 2016 cropping seasons**

123 **2. Effects of Zinc rates on growth**


124 Fig 2 illustrates the effect of Zn fertilizer application on plant height, stem girth and number  
 125 of leaves over the period of the experiment. Analysis of variance showed no significant effect  
 126 of Zn fertilizer on plant height and leaf production. However, plots with Zn application had  
 127 the tallest plants compare to the control. Branch production was significantly affected by Zn  
 128 rates in both seasons. At 30 DAS, the 5 kg/ha treatment effect was significantly higher than  
 129 other treatment effects. Treatment effect at 45 DAS was similar. At 60 DAS, the control  
 130 treatment effect was significantly lower ( $P < 0.05$ ) than all Zinc treatments. **Variety by zinc**  
 131 **rates interaction** was not significant at 5% level of probability on all sampling days.



132




133




134


d



135



136



137



138 **Figure 2. Effects of Zinc rates on plant height (a and b), stem girth (c and d) and**  
 139 **number of leaves (e and f) in 2016 cropping seasons**

140 **3. Nodulation parameters**

141 Results on number of nodules per plant, effective nodules per plant and nodule dry weight per  
 142 plant as influenced by cowpea varieties and Zinc fertilizer application in the two sampling  
 143 periods in both experiments are presented in Tables 1 and 2. Treatment differences for all  
 144 parameters on all days at both seasons were not significant ( $P > 0.05$ ). The interaction effect  
 145 was also not significant at 5% probability. Nodule number was nearly successively decreased  
 146 over time at all treatments and is not correlated with the Zinc fertilizer applied. No interaction  
 147 effect was significant for all parameters at all sampling periods.

148 **Table 1. Effect of cowpea varieties in changes of nodule number (nodules per plant),**  
 149 **nodule dry weight (g per plant) and effective nodules (%)**

Varieties	Time (Days after sowing)			
	Major season		Minor season	
	30	45	30	45
	<b>Nodule number (nodules per plant)</b>			
Agyenkwa	5	3	7	6
Asontem	8	5	10	6
Zamzam	5	4	8	6
LSD (0.05)	NS	NS	NS	NS
CV (%)	21.8	31.8	14.2	22.4
	<b>Nodule dry weight (g per plant)</b>			
Agyenkwa	0.06	0.09	0.22	0.16
Asontem	0.04	0.08	0.19	0.14
Zamzam	0.05	0.12	0.18	0.13
LSD (0.05)	NS	NS	NS	NS
CV (%)	18.8	21.0	10.5	20.3
	<b>Effective nodules (%)</b>			
Agyenkwa	82.54	37.82	84.48	47.55
Asontem	74.44	32.64	76.56	42.11
Zamzam	76.94	65.13	79.72	68.27
LSD (0.05)	NS	NS	NS	NS
CV (%)	4.9	5.5	2.4	13.3
<b>Variety x Zn rate</b>	NS	NS	NS	NS

150

151 **Table 2. Changes in nodule number (nodules per plant), nodule dry weight (g per plant)**  
 152 **and effective nodules (%) of cowpea growing under Zinc foliar application**

Rates	Time (Days after sowing)			
	Major season		Minor season	
	30	45	30	45
	<b>Nodule number (nodules per plant)</b>			
0	6	4	8	6
5	5	4	8	5
10	7	4	9	6
LSD (0.05)	NS	NS	NS	NS
CV (%)	11.3	13.8	5.0	11.9

<b>Nodule dry weight (g per plant)</b>				
0	0.05	0.11	0.18	0.15
5	0.05	0.09	0.20	0.15
10	0.06	0.08	0.21	0.14
LSD (0.05)	NS	NS	NS	NS
CV (%)	9.8	22.8	7.1	28.5
<b>Effective nodules (%)</b>				
0	78.56	43.33	81.97	49.25
5	74.26	53.11	76.48	48.88
10	81.11	39.16	82.31	59.80
LSD (0.05)	NS	NS	NS	NS
CV (%)	3.6	8.8	4.4	8.4
<b>Variety x Zn rate</b>	NS	NS	NS	NS

#### 153 4. Yield and harvest index

154 The cowpea grain yield was significant ( $P < 0.05$ ) under Zinc fertilizer application in all the  
 155 two seasons (Table 3). Cowpea grain yield recorded on the application of Zinc fertilizer at 5  
 156 kg/ha increased at 28 % for Agyenkwa > Zamzam (20 %) > Asontem (19%) compare to the  
 157 control in major season and the minor season the results followed the same trend. The Zinc  
 158 levels are increased the cowpea grain yield in the order: 5 kg Zn/ha > 10 kg Zn/ha > 0 kg  
 159 Zn/ha during all cropping seasons (2016 major and minor seasons). The cowpea grain yield  
 160 decline over increasing the Zinc rate beyond 5 kg/ha. There was about 6 % and 10 % yield  
 161 reduction in the main season obtained with Agyenkwa and Asontem respectively. 100 seed  
 162 weights were different at 5% level of probability. One hundred seeds weight was higher with  
 163 Zamzam following by Agyenkwa and at the end Asontem with the lowest one. The shoot dry  
 164 weight was significant ( $p < 0.05$ ) affect by Zinc fertilizer application. Similarly, cowpea  
 165 varieties did significant ( $P < 0.05$ ) affect the cowpea biomass yield in all sampling periods  
 166 and the interaction follows the same trend.

167

168 **Table 3. Effects of Zn rates on harvest index, haulm and grain yield of cowpea**

Treatments	Major season			Minor season		
	100 seeds weight ← g →	Haulm ← Kg/ha →	Grain yield	100 seeds weight ← g →	Haulm ← Kg/ha →	Grain yield
<u>Varieties</u>						
Agyenkwa	16.25	1382.74	1142.23	16.20	1311.58	1620
Asontem	13.63	1596.68	1082.15	13.26	1650.58	1326
Zamzam	17.18	1470.64	1423.62	17.07	1630.33	1707
LSD (0.05)	1.29	342.94	268.69	1.25	317.90	125.2
CV (%)	4.7	13.40	12.80	4.7	6.40	6.1
<u>Zn levels (kg.ha<sup>-1</sup>)</u>						
0	15.34	1451.48	1087.45	14.93	1440.58	1493
5	15.58	1440.24	1283.94	15.61	1587.67	1600
10	16.14	1558.34	1276.60	16.00	1564.25	1560
LSD (0.05)	NS	225.13	120.87	0.79	218.492	79.4
CV (%)	5.5	17.20	16.00	4.7	10.40	4.7

169

170 **5. N-fixed and crude protein**

171 The results of cowpea N-fixed and crude protein at all sampling periods are presented in  
 172 Table 4. In this study Zinc fertilizer application interacted to significantly ( $p < 0.05$ ) affect  
 173 cowpea N-fixed and crude protein. Zinc levels increased the cowpea N-fixed and crude  
 174 protein in the order: 5 kg Zn/ha > 10 kg Zn/ha > 0 kg Zn/ha. N-fixed and crude protein  
 175 interaction differed significantly ( $p < 0.05$ ) among some the treatment interactions. Zamzam  
 176 variety interacted markedly to produce the highest value of N-fixed in cowpea haulm and  
 177 grain and Agyenkwa presented the lowest one. For the cowpea crude protein, the result is  
 178 presented in the following order: Asontem > Agyenkwa > Zamzam.

179

180

181

182 **Table 4. Effects of Zn rates on cowpea N-fixed and crude protein**

Treatment	Major season			Minor season		
	Grain N-fixed ← Kg.ha <sup>-1</sup> →	Tissue N-fixed →	Crude protein (%)	Grain N-fixed ← Kg.ha <sup>-1</sup> →	Tissue N-fixed →	Crude protein (%)
<u>Varieties</u>						
Agyenkwa	22.40	26.72	25.80	39.61	15.70	24.84
Asontem	26.40	27.64	29.44	32.23	16.59	26.86
Zamzam	31.80	30.82	24.87	39.14	20.95	23.44
LSD (0.05)	NS	12.32	0.91	4.81	NS	1.46
CV (%)	16.60	13.50	2.00	10.40	3.20	3.40
<u>Zn levels (kg.ha<sup>-1</sup>)</u>						
0	21.80	26.12	26.88	33.59	14.12	24.57
5	30.90	27.26	27.28	38.39	20.09	25.56
10	27.90	31.79	25.95	39.00	19.03	25.01
LSD (0.05)	5.26	7.11	0.59	4.81	5.58	1.15
CV (%)	17.30	14.30	1.40	4.00	2.30	4.50
<b>Variety x Zn rate</b>	*	*	*	*	*	*

183

184 **6. Effects of Zinc rates on NPK content**

185 The results of grain nutrients analysis showed no significant varietal effects for content of  
 186 nitrogen and Phosphorus (Table 5). Additionally, Potassium content in the haulms was not  
 187 different among varieties. However, for Potassium content in seed, the Zn treatments effects  
 188 were similar, but greater either effect was greater than the control treatment effect in both  
 189 seasons.

190 **Table 5. NPK content of cowpea as affected by varieties**

Rates	Nutrient uptakes (kg/ha)			
	Major season		Minor season	
	Haulm	Grain	Haulm	Grain
	N			
Agyenkwa	40.92	47.21	29.90	64.41
Asontem	41.84	51.18	30.80	57.03
Zamzam	45.46	56.61	35.20	63.94

LSD (0.05)	NS	NS	NS	NS
CV (%)	8.4	6.9	9.40	6.2

**P**

Agyenkwa	3.23	5.46	3.53	12.19
Asontem	3.39	5.09	3.80	14.33
Zamzam	2.75	5.68	3.77	12.33
LSD (0.05)	NS	NS	NS	NS
CV (%)	7.5	6.5	7.1	7.70

**K**

Agyenkwa	14.32	12.01	13.95	20.92
Asontem	13.17	13.61	13.58	16.84
Zamzam	10.79	13.83	11.72	20.43
LSD (0.05)	NS	1.27	NS	2.89
CV (%)	12.1	7.20	7.70	4.30

191

192 For the Zn treatments, N content of grain was affected by Zn application in both seasons.

193 Haulm N content was also significantly affected by Zn fertilizer application. In all these

194 cases, treatments differences between the Zinc treatments were similar, but either effect was

195 greater than the control treatment (Table 6). Haulm P content were significantly affected by

196 Zn fertilizer application (Table 6), with the exception haulm Zinc content in the major season,

197 where the control treatment effect was similar to the 5 kg/ha Zn treatment. In all cases, the Zn

198 treatment effects were similar, and either effect was significantly higher than the control

199 treatment effect. Haulm K content was not affected by Zn fertilizer in both seasons (Table 6).

200 However, grain K content significantly affected by the Zn fertilizer in both seasons. In both

201 seasons, grain K content in the control treatment was lower than the Zn treatment effects.

202 **Table 6. NPK content of cowpea as affected by Zinc fertilizer**

Rates	Nutrient uptakes (kg/ha)			
	Major season		Minor season	
	Haulm	Grain	Haulm	Grain
0	40.32	46.63	28.30	58.39

5	41.46	55.69	34.30	63.19
10	45.99	52.68	33.20	63.80
LSD (0.05)	NS	5.26	5.59	2.29
CV (%)	8.90	8.20	9.40	2.40

**P**



0	2.94	4.70	3.22	12.01
5	2.89	6.09	3.88	13.61
10	3.54	5.44	4.01	13.83
LSD (0.05)	0.55	0.70	0.65	1.27
CV (%)	9.20	13.50	9.60	7.20

**K**



0	12.45	12.01	12.42	17.84
5	13.38	13.61	14.38	20.27
10	12.45	13.83	12.45	20.08
LSD (0.05)	NS	1.27	NS	1.67
CV (%)	14.30	7.20	17.30	8.10

203 **7. Interrelationship between Zinc and NPK uptake in plant grain**



204 The linear regression showed the positive relationship between grain Zn uptake and NPK  
 205 content for three sampling periods during the experiment in the major and minor seasons  
 206 (Fig. 3). The argument on the enhanced NPK uptake y Zn content was ably supported by the  
 207 significant positive relationship observed in the present study between NK and Zn uptake  
 208 (0.9929\*\*\* with N and 0.9096\*\* with K) in the major cropping season. The minor cropping  
 209 season also follows the same trend with 0.9942\*\* and 0.9389\*\* with N and K respectively.  
 210 And with P the relationship was weak but positive (0.3839 in major season and 0.7289 in  
 211 minor season).



212



213



214

215 **Figure 3. Relationship between Zn uptake and the macronutrients (NPK) in 2016**  
216 **cropping seasons**

## 217 **DISCUSSION**

218 Plant height was affected by both variety and Zn rates. Among the Zn treatments, Plant  
219 height was greatest in the Zn plots applied in both cropping seasons. [17] reported that added  
220 Zn significantly increased plant height by increasing internodes distances. [18] stated that  
221 grain yield was positively correlated with leaf weight, stem weight, plant height and number  
222 of branching per plant. Zinc fertilizer application did not, however, have any significant  
223 effect on the number of leaves and stem girth in all the days examined. It was reported that  
224 application of zinc had positive effects on growth parameters [19]. Contrarily, foliar  
225 application of micronutrients increased the diameter of plant over the control treatment [20].  
226 So, these findings conclude that the entire cowpea varieties gave equal stem diameter at all  
227 treatments of zinc application.

228 The application of the Zn fertilizer did not affect nodulation, indicating that some of  
229 inoculation factors were limiting such as soil pH, initial phosphorus and others  
230 micronutrients. And also, [21] reported that the nodule initiation may depend on the relative   
231 concentrations of plant-specific signals and host species appears to be a significant factor  
232 determining the maximum number of nodules generated. Effective nodule is essential for a  
233 functioning Legumes-Rhizobium symbiosis and Zinc, chloride and cobalt have no effect on  
234 nodulation but are required for the growth of the host legume [22]. Two hosts may have the  
235 same sensitivity to bacterial signal molecules, but might differ in their ability to elicit  
236 synthesis of required nodulation signals in the bacteria [23]. Cowpea root exudates have also  
237 been reported to contain substances that enhance nodule initiation [24, 25, 26]. However,

238 lower efficiency of cowpea cannot be readily explained in terms of reduced numbers of  
239 bacteria in contact with the root [21]. Varieties most susceptible to infection and capable of  
240 producing effective nodules should have greater potential to fix more atmospheric N.  
241 However, this assumption often depends on other factors such as the environment and crop  
242 management [27]. Indeed [28] reported that the ability to form nodules is not enough to  
243 obtain an effective nitrogen fixation symbiosis. Nodule number was nearly successively  
244 decreased over time at all treatments and is not correlated with the Zinc fertilizer applied. The  
245 amount of nodule dry biomass was drastically reduced with the mineral Zinc fertilizer,  
246 whereas the amount of nodule biomass was not affected in the control group, probably  
247 because the soil had satisfactory levels of available N and P. Nodule number correlated  
248 negatively with nodule dry weight [29]. The interaction effect was also not significant at 5%  
249 probability.

250 The present results were supported by [30] who reported that foliar application of  
251 micronutrients help in improving yield. In both seasons, foliar spray of Zn fertilizer had effect  
252 on hundred grain weights. In all these parameters, the control treatment effect was lower than  
253 Zn treatments, whereas among the Zn treatments. [31] reported that following Zn fertilization  
254 increased hundred seed weight. Also, [32] reported that yield and its components in lentil are  
255 improved by foliar application of micronutrients. Crop yields and quality are reduced by Zn  
256 inadequate in soil; therefore, Zn utilization is essential to obtain high yield and quality in crops as  
257 showed the results (Table 3). These results are in close conformity with those of [33, 34, 35].  
258 This was because of the fact that better and higher availability of Zinc, resulting better  
259 nutritional environment, higher dry matter accumulation and its associated effect on growth  
260 attributes increased haulm and grain yield. It is also evident from table 3 that all the Zinc  
261 treated plots increased the grain yield over the control, as there was a consistent increase in

262 cowpea grain yield up to 10 kg Zn/ha. This suggests that, the application of Zn significantly  
263 affect cowpea yield. Similar results were reported as in [36] and [37]. In this connection, [38]  
264 reported that the foliar application of Zn affected yield and its components of soybean. Also,  
265 [39] reported that the highest yield of common bean was obtained by Zinc foliar application.  
266 [40] believe that more production of chlorophyll and IAA can cause delay in plant oldness  
267 and prolong the period of photosynthesis. This incident improves the production of  
268 carbohydrates and their transportation to the growing seeds.

269 The Zn deficiency symptoms can be prevented by the application of Zn fertilizers. The actual  
270 causal relationship and mechanisms are still not fully understood [3]. As shown in Tables 4  
271 and 5 the mean percentage total nitrogen, phosphorus and potassium uptakes in the harvested  
272 leaves were quantitatively higher under zinc fertilizer application and increased with  
273 incremental zinc rates. These results corroborate the findings of [41] and [42] who reported  
274 that zinc is an essential micronutrient for plant growth and plays an important role in the  
275 catalytic part of several enzymes its deficiency will result in stunted growth and nutrient  
276 uptakes. And also, [43] reported that zinc exerts a great influence on basic plant life  
277 processes, such as (i) nitrogen metabolism – uptake of nitrogen and protein quality; (ii)  
278 photosynthesis - chlorophyll synthesis and carbon anhydrase activity. Also many researchers  
279 have observed that Zn is closely related to the nitrogen metabolism pathway of plants, thus  
280 causing a reduction in protein synthesis for Zn deficient plants. Zinc deficiency significantly  
281 affects the root system including root development [44].

## 282 CONCLUSION

283 Zn fertilizer significantly affected NPK content and grain yield of cowpea varieties used. The  
284 increment of Zn content in the grain had a positive relationship with NK, which will

285 definitely enhance nutrition of both human and animals. At all sampling periods, nodule  
286 number per plant was not affected by Zinc rates and nodule number was nearly successively  
287 decreased over time at all treatments and is not correlated with the Zinc fertilizer applied. The  
288 Zinc fertilizer significantly enhanced N-Fixed and Crude protein in both cropping season's  
289 trial investigating effect of Zinc rates on growth, nodulation and mineral content of cowpea in  
290 the semi-deciduous forest zone of Ghana. This implies the Zn rates used can be applied to  
291 any of the varieties used.

## 292 REFERENCES

- 293 1. Olsen SR. Micronutrient Interactions. *JM Mortved, JJ Goirdano, WL Lindsay (eds)*  
294 *Micronutr Agric Soil Sci Soc Am.* 1972;6(August):243-264.
- 295 2. Imtiaz M., Alloway BJ., Shah KH., Siddiqui SH., Memon MY., Alsam M. and Khan  
296 P. Zinc Nutrition of Wheat: II: Interaction of Zinc with other Trace Elements. *Asian J*  
297 *Plant Sci.* 2003;2:156-160.
- 298 3. Alloway BJ. Zinc in Soils and Crop Nutrition. Second edition, published by IZA and  
299 IFA, Brussels, Belgium and Paris, France. 2008;139.
- 300 4. Unkovich, M. and Baldock, J. Measurement of asymbiotic N<sub>2</sub> fixation in Australian  
301 agriculture. *Soil Biol Biochem.* 2008;40:2912 – 2921.
- 302 5. Craufurd PQ., Summerfield RJ., Ellis RH. and Roberts EH.. Photoperiod, temperature,  
303 and the growth and development of cowpea. *Adv cowpea Res Int Inst Trop Agric*  
304 *Ibadan, Niger Japan Int Res Cent Agric Sci Tsukuba, Ibaraki, Japan.* 1996;II:75-86.
- 305 6. Gates M. Bulletin of Tropical Legumes. *Area.* 2011;(October):5-8.
- 306 7. Hall AE. Breeding for adaptation to drought and heat in cowpea. *Eur J Agron.*

- 307           2004;21(4):447-454.
- 308    8.    Epstein E. and Bloom AJ. Mineral Nutrition of Plants: Principles and Perspectives.  
309           *Sinauer Assoc;*. 2005.
- 310    9.    Latham MC. Human Nutrition in the Developing World. Food and N. Rome/Italy;  
311           1997.
- 312    10.   FAO and WHO. Vitamin and Mineral Requirements in Human Nutrition Second  
313           Edition.; 1998.
- 314    11.   AOAC. Official Methods of Analysis. 17th ed. Gaithersburg, Maryland, USA, AOAC  
315           International; 2000.
- 316    12.   Jones DB. Factors for Converting Percentages of Nitrogen in Foods and Feeds into  
317           Percentages of Proteins. Circular No. 183. Washington, DC, United States Department  
318           of Agriculture.; 1941.
- 319    13.   Varvel GE. and Peterson TA. Nitrogen fertilizer recovery by corn in monoculture and  
320           rotation systems. *Agron J.* 1990;82:935-938.
- 321    14.   Motsa, M.R. and Roy RN. Guide to Laboratory Establishment for Plant Nutrient  
322           Analysis. FAO. Rome, Italy; 2008.
- 323    15.   Moss P. Limits of interference by Fe, Mn, Al and phosphate in the EDTA  
324           determination of Calcium in the presence of Mg using calcon-red as indicator. *J Sci, F*  
325           *Agric.* 1961;12:30-34.
- 326    16.   Okalebo JR., Gathua KW. and Woomer PL. Laboratory Methods of Soil and Plant  
327           Analysis: *A Working Manual*. Vol 1. Tropical S. Nairobi/Kenya: Soil Science Society  
328           of East Africa; 1993.

- 329 17. Kaya C., Higgs D. and Burton A. Phosphorus acid phosphates enzyme activity in  
330 leaves in leaves of tomato cultivars in relation to Zn supply. *Commun Soil Sci Plant*  
331 *Anal.* 2000;31:3239-3248.
- 332 18. Jatasra, D.S. and Dahiya B. Relative importance of forage yield components in  
333 cowpeas under dry land conditions. *Indian J Agric Resour.* 1988;22(1):1-5.
- 334 19. Malakouti MJ. The Effect of Micronutrients in Ensuring Efficient Use of  
335 Macronutrients. *Turkish J Agric For.* 2008;32:215-220.
- 336 20. Shuman LM. Fractionation Method for Soil Micronutrients. *Soil Sci.* 1985;140:11-22.
- 337 21. Bhuvanewari, T.V., Lesniak, A.P., and Wolfgang D. Efficiency of Nodule Initiation  
338 in Cowpea and Soybean. *Plant Physiol.* 1988;86:1210-1215.
- 339 22. FAO. Legume Inoculants and Their Use. Rome; 1984.
- 340 23. Peters, N.K, Frost, J.W and Long S. A plant flavone, luteolin induces expression of  
341 *Rhizobium meliloti* nodulation genes. *Sci Agric.* 1986;234:977-980.
- 342 24. Bhagwat A.A and Thomas J. Legume-Rhizobium interactions: cowpea root exudate  
343 elicits faster nodulation response by *Rhizobium* species. *Appl Environ Microbiol.*  
344 1982;43:800-805.
- 345 25. Bhagwat, A.A and Thomas J. Legume-Rhizobium interactions: role of cowpea root  
346 exudate in polysaccharide synthesis and infectivity of *Rhizobium* species. *Arch*  
347 *Microbiol.* 1983;136:102-105.
- 348 26. Bhagwat, A.A and Thomas J. Legume-Rhizobium interactions: host induced  
349 alterations in capsular polysaccharides and infectivity of cowpea rhizobia. *Arch*  
350 *Microbiol Microbiol.* 1984;140:260-264.

- 351 27. Kellman AW. Rhizobium inoculation cultivar and management effects on the growth,  
352 development and yield of common bean (*Phaseolus vulgaris* L.). 2008.
- 353 28. Giller KE. Nitrogen Fixation in Tropical Cropping System. Wallingfor. UK; 2001.
- 354 29. Sarkodie-Addo J. Evaluation of Bradyrhizobium japonicum isolates from Ontario 74  
355 soybean fields. 1991.
- 356 30. Arif M., Chohan MA., Ali S., Gul R. and Khan S. Response of wheat to foliar  
357 application of nutrients. *J Agric Biol Sci.* 2006;1:30-34.
- 358 31. Yilmaz A., Ekiz H., Torun B., Gulekin I., Karanlink S., Bagci S.A. and Cakmak I.  
359 Effect of different zinc application methods on grain yield and zinc concentration in  
360 wheat cultivars grown on zincdeficient calcareous soils. *J Plant Nutr.* 1997;20:461-  
361 471.
- 362 32. Zeidan M.S., Hozayn M., and Abd El-Salam MEE. Yield and quality of lentil as  
363 affected by micronutrient deficiencies in sandy soils. *J Appl Sci Res.* 2006;2:1342-  
364 1345.
- 365 33. Sharma SK. and Jat ML. Effect of sulphur on growth and yield of cowpea [*Vigna*  
366 *unguiculata* (L.) Walp]. *Ann Agric Res New Ser.* 2003;24:215-216.
- 367 34. Yadav SS. Growth and yield of cowpea [*Vigna unguiculata* (L.) Walp] as influenced  
368 by phosphorus and sulphur fertilization. *Haryana J Agron.* 2004;20:10-12.
- 369 35. Tripathi H.C., Pathak R.K., Kumar A. and Dimsec S. Effect of sulphur and zinc on  
370 yield attributes, yield and nutrient uptake in chickpea. *Ann Plant Soil Res.* 2011;13:134-  
371 136.
- 372 36. Moswatsi MS. Response of cowpea to variable rates and methods of zinc. Thesis,

- 373 University of Limpopo. 2015; 77.
- 374 37. Oseni T. Growth and Zinc Uptake of Sorghum and Cowpea in Response to Phosphorus  
375 and Zinc Fertilization. *World J Agric Sci.* 2009;5(6):670-674.
- 376 38. Banks L. Effect of timing of foliar zinc fertilizer on yield component of soybean. *Aust*  
377 *J Exp Agric Anim Husb.* 2004;22:226-231.
- 378 39. Seifi NM., Yarnia M. and Rahimzade KF. Effect of zinc and manganese and their  
379 application method on yield and yield components of common bean (*Phaseolus*  
380 *vulgaris* L. cv. Khomein). *Middle-East J Sci Res.* 2011;8:859-865.
- 381 40. Hemantaranjan, A. and Grag OK. Iron and zinc fertilization with reference to the grain  
382 quality of triticum Aestivum L. *J Plant Nutr.* 1988;11:1439-1450.
- 383 41. Fageria NK. Dry matter yield and nutrient uptake by lowland rice at different growth  
384 stages. *J Plant Nutr.* 2004;27:947-958.
- 385 42. Sunitha, K Padma, S N Vasandha, S and Anitha S. Microbial Inoculants- A Boon to  
386 Zinc Deficient Constraints in Plants : A Review. *Int J Sci Res Publ.* 2014;4(6):4-7.
- 387 43. Potarzycki, J. and Grzebisz W. Effect of zinc foliar application on grain yield of maize  
388 and its yielding components. *Plant, Soil Environ.* 2009;55:519-527.
- 389 44. Fageria N. Influence of micronutrients on dry matter yield and interaction with other  
390 nutrients in annual crops. *Pesq Agropec Bras.* 2002;37:1765-1772.