

Evaluation of the nutritional status and acceptability of powdered reconstituted Kunu-zaki; an index of increasing shelf life

ABSTRACT

Increasing the shelf life of Kunu zaki via conversion to powder form at the same time retaining its nutritional and sensory acceptability was evaluated on different cereal grains often used in the production of Kunu (Maize, sorghum, and millet). The grains were bought from Ibadan Market, sorted carefully after which equal mass (250g) of the sorted grains were soaked for seven two (72) hours, drained and was milled with the inclusion of other ingredients as spice. The mixture was sieved to remove the chaff and the liquid portion was allowed to Sediment after which it the upper layer containing water was decanted and the sediment layer was further processed to powdered form via drying. Sensory evaluation of the freshly prepared Kunu and powdered reconstituted prepared kunu was evaluated and compared. Similarly, the proximate, mineral composition and an assessment to verify any storage challenges such as insect infestation were investigated. The result obtained show that the shelf life of Kunu-zaki was elongated from three days to more than six month as a result of the conversion to powdered form. The sensory evaluation of freshly prepared and powdered reconstituted Kunu-zaki showed that there is no clear cut significant difference in the overall acceptability between freshly prepared kunu and powdered reconstituted Kunu. Sorghum showed a better appearance, color and aroma both in the freshly prepared Kunu and powdered reconstituted Kunu above the other grains used. The proximate analysis carried out showed that the processed powdered form of Kunu had a significant increase in protein, lipid and carbohydrate with a drastic reduction in the moisture content when compared with the freshly prepared Kunu. Also, Fe, Cu, Zn, Co, Mg are the predominant minerals measured in mg/100g. It can be concluded that powdered reconstituted Kunu-zaki showed no significant different in the major nutritional, sensory evaluated properties and proven to have an elongated shelf life for more than 6months with no insect infestation during storage when compared to freshly prepared Kunu.

Keywords: Kunu Zaki, Sensory properties, Shelf life, Nutritional evaluation, Insect infestation, storage

INTRODUCTION

One of the mostly consumed non alcoholic beverages especially in the Northern part of Nigeria is Kunu Zaki^{1,2}, which tend to deteriorate with time owing to its short shelf life³ and has thus, become a concern and difficult to produced in large quantity knowing fully that the non alcoholic beverage is on high demand due to its quenching properties⁴, a source of energy and soothing taste⁵. The wastage incurred when produced in large quantity has dramatically reduced the income of the manufacturers. Different grains in the production of

Kunu Zaki and inclusion of several spices has only been advantageous to the nutritional enhancement not to its shelf life. Different types of grains in conjunction with spices such as ginger, black pepper, garlic, red pepper, and clover are commonly used as raw materials. Examples of grains being used include Millet, sorghum, maize, groundnut amongst others⁶. However, the choice of any of these grains is dependent upon its relative abundance in any location. Consequently, the compound name of Kunu is determined by the grain is made of. For instance, *Kunun-gero* is the name for Kunu made from millet, *Kunun-dawa* for the one made from sorghum, *Kunun-masara* for the one from maize, *Kunun-acha* from hungry rice or acha, *Kunun-shinkafa* from rice, *Kunun-gyada* from groundnut, *Kunun-tsamiya* from tamarind, and *Kunun-kanwa* from potassium hydroxide⁷. However, different culture in a way to increase the shelf life of Kunnu involved the use of several processing procedure which only was successful in only the sensory properties. The major area of concern is the short shelf life of this drink³. Reasons for this demerit include the unhygienic local technology being employed, the high moisture level of the product, microorganisms present in the drink that helps in the fermentation process, rodents and insect pests present in the vicinity during preparation amongst others^{8,9}. This study is aimed at addressing this challenge. Hence, Kunu made from three different sources (i.e. maize, millet, and wheat) were converted into a powdered form in order to reduce their moisture contents as to increase the shelf life. These powders were further investigated in term of shelf life, nutritional value and sensory evaluation. The goal is to ascertain a lasting solution to the spoilage and short shelf life of Kunu and also to establish the best grain in terms of nutritional, shelf life, titratable acid, sensory evaluated properties when compared to the conventional mode of production.

Methodology

Production of powdered Kunu

The grains were bought from Ibadan Market, sorted carefully after which equal mass (250g) of the sorted grains where soaked for seven two (72) hours, drained and was milled with the inclusion of other ingredients as spice. The mixture was sieved to remove the chaff and the liquid portion was allowed to Sediment after which the upper layer containing water was decanted and the sediment layer was further processed to powdered form via drying. The powdered sample can further be reconstituted to Kunu-zaki

71 **Mineral Analysis**

72 9ml of concentrated HCl was added into 1.5g of the sample, followed by 3ml of Conc. HNO₃
73 and heated on a hot plate slowly at first, until frothing ceases. Heating was continued until
74 HNO₃ evaporated and white fumes observed. It will be allowed to cool and filtered. This was
75 diluted and made up to 100ml with distilled water. The following elements were determined
76 using atomic absorption spectrophotometer. The elements are calcium, chromium, manganese
77 copper, zinc, iron, magnesium and nickel, potassium, Cobalt, Molybdenum, Selenium,
78 vanadium.

79 **Proximate composition**

80 Estimations were made of nitrogen (as an index of crude protein), water, fat, ash, and crude
81 fiber. When the total was subtracted from 100%, the difference was termed carbohydrate by
82 difference. Determination of the moisture content, ash, and crude fat followed the method of
83 AOAC (2005)¹⁰. Crude fiber determination followed the method of Pearson (1981)¹¹.
84 Estimation of nitrogen content was by the Kjeldahl method multiplied by 6.25, the nitrogen-
85 protein factor to convert to crude protein.

86 **Crude fiber**

87 Two grams of the sample was transferred into a 1 L conical flask. One hundred milliliters of
88 sulfuric acid (0.255 mol/L) was heated to boiling and then introduced into the conical flask
89 containing the sample. The contents were then boiled for 30 min, ensuring that the level of
90 the acid was maintained by the addition of distilled water. After 30 min, the contents were
91 then filtered through a muslin cloth held in a funnel. The residue was rinsed thoroughly until
92 its washing was no longer acidic to litmus. The residue was then transferred into a conical
93 flask. One hundred milliliters of sodium hydroxide (0.313 mol/L) was then brought to boil
94 and then introduced into the conical flask containing the sample. The contents were then
95 boiled for 30 min, ensuring that the level of the acid was maintained by the addition of
96 distilled water. After 30 min, the contents were then filtered through a muslin cloth held in a
97 funnel. The residue was rinsed thoroughly until its washing was no longer alkali. The residue
98 was then introduced into an already dried crucible and ashed at 600°C ± 200°C.

99 **Sensory evaluation**

The 9-point hedonic scale assessment and the paired comparison tests were used as described by Larmond 1977¹². A total number of 30 staff of the Nigerian stored products research institute and industrial training student of the university of Ibadan were selected based on their familiarity with Kunu zaki, The panelists scored coded drinks in terms of degree of liking to taste, color, and aroma. The 9-point hedonic scale used by the panelists for the evaluation ranged from 1 to 9 representing “extremely dislike” to “extremely like”. The coded samples were served in clean, transparent cups at room temperature 25°C. Water was given to each panelist for oral rinsing in between tasting of the samples.

Result

The proximate analysis carried out on the powdered Kunu showed a similar nutritional status with the freshly prepared non- powdered-kunu. Although the observable difference was seen in the moisture and carbohydrate content while there was no significant difference in protein, lipid, ash and crude fibre. Powdered Sorghum also showed the highest amount (%) of protein, lipid and Ash when compared with other grains which is also closely related to the freshly prepared non powdered kunu.

Table 1: Proximate analysis of powdered reconstituted Kunu zaki (P-Kunu) and freshly prepared non powdered Kunu (F-kunu) produced from three different grains (Sorghum, maize and millet)

Proximate	P. Maize	F. Maize	P. Sorghum	F. Sorghum	P. Millet	F. millet
Moisture	9.2±0.09 ^a	39.2±0.09 ^b	9.3±0.08 ^b	38.3±0.08 ^a	8.9±0.09 ^a	38.9±0.09 ^b
Protein	12.3±0.09 ^a	15.3±0.09 ^a	14.5±0.09 ^b	17.5±0.09 ^b	11.2±0.09 ^c	14.2±0.09 ^c
Lipid	1.3±0.08 ^a	1.27±0.08 ^a	1.9±0.09 ^b	1.9±0.09 ^b	1.7±0.08 ^b	1.7±0.08 ^b
Ash	2.8±0.08 ^a	2.77±0.08 ^a	3.8±0.08 ^a	3.8±0.08 ^a	1.9±0.08 ^a	1.9±0.08 ^a
Crude fibre	3.1±0.08 ^b	8.13±0.08 ^a	2.7±0.08 ^a	7.7±0.08 ^b	3.4±0.08 ^b	8.4±0.08 ^a
Carbohydrate	71.4±0.08 ^a	33.4±0.08 ^b	68.4±0.08 ^b	30.8±0.08 ^a	72.9±0.12 ^a	34.96±0.12 ^b

Means with same superscript indicate no significant different at 5% for the attribute

*P.maize, P. sorghum, P. millet= **Powdered** Kunu from maize, Sorghum and millet

*F.maize, F. sorghum, F. millet= **freshly** prepared Kunu not converted to powdered form (conventional mode of traditional preparation of Kunu) produced from maize, Sorghum and millet

The result findings on the sensory properties of powdered reconstituted Kunu and conventional mode of freshly prepared Kunu produced from different grains showed a similar overall acceptability with sorghum having an appealing aroma and a pleasing appearance for

both powdered reconstituted Kunu and conventional mode of freshly prepared. Similarly, aroma, taste and color of millet was found to have no significant different in Kunu from the powdered form and the conventional preparation. The overall acceptability for both powdered reconstituted and conventional mode of preparation was in favor of sorghum when compared with the other grains used.




Fig 1: Sensory properties of powdered reconstituted Kunu and conventional mode of freshly prepared Kunu produced from three different grains

Having being monitored for the emergence of insect pests in the powdered samples of Kunu, the result is presented in the table below. The outcome of the investigation revealed no occurrence and development of flour beetles in the powdered Kunu produced from Maize, Millet and Sorghum.

Table 2: Emergence of flour beetles in different powdered grains stored for 6 months for reconstitution into Kunu (Nigerian Non-alcoholic beverage)

Sample	T. castaneum	C. cephalonica	T. confusum
Powder Maize	—	—	—
Powder millet	—	—	—
Powder Sorghum	—	—	—

⊕ = presence of flour beetles

— = Absence of flour beetles

T. castaneum = *Tribolium castaneum* (Herbst)

C. cephalonica = *Corcyra cephalonica* (Stainton)

T. confusum = *Tribolium confusum* (J. du Val)

Mineral composition of Kunu zaki converted to powder form produced from three different grains (Sorghum, maize and millet) showed a commendable erythropoietin potentials with high amount of Fe, Cu, Co, which are minerals associated with erythropoiesis. Also, the amount of calcium and potassium of the three grains are estimable for several regulation of body system such as the electrolyte regulation and the blood pressure.

Table 4: Mineral composition of Kunu zaki converted to powder form produced from three different grains (Sorghum, maize and millet)

Minerals	Maize	Sorghum	Millet
Cobalt (mg/100g)	0.010±0.00 ^a	0.02±0.03 ^b	0.017±0.003 ^{ab}
Iron (mg/100g)	6.41±0.04 ^a	14.71±0.05 ^b	7.15±0.06 ^c
Zinc (mg/100g)	0.27±0.02 ^a	0.53±0.03 ^b	0.47±0.004 ^b
Copper (mg/100g)	0.17±0.03 ^a	0.50±0.06 ^b	0.37±0.03 ^b
Calcium (mg/100g)	92.0±2.89 ^a	76.0±16 ^b	101.67±2.23 ^c
Magnesium (mg/100g)	2.47±0.02 ^a	4.37±0.08 ^a	1.91±0.04 ^a
Potassium (mg/100g)	26.67±1.67 ^a	35.00±0.00 ^b	38.33±1.67 ^b

Means with same superscript indicate no significant different at 5% for the attribute

Discussion

Low shelf life and spoilage of food commodities have been the bane of productivity in the tropics. This could be attributed to high temperature and humidity of the region which enhances the activities of microbes and pest organisms in storage environment¹³. This has constituted a clog in the wheel of exportation in agrarian countries like Nigeria. In a bid to addressing this challenge, effort was made in the present study to convert the freshly prepared Kunu drinks (produced from three different grains) into powdered form. After 6 months on the shelf, these powders were reconstituted and their organoleptic properties and nutritional status were evaluated and compared with those of freshly prepared Kunu (produced from three different grains). The result showed that the sensory attributes of the powdered reconstituted kunu was not significantly different from those of the freshly prepared drinks. This is in consonance with the result of the proximate analysis which shows no significant difference except in terms of moisture and carbohydrate content. From the forgoing, one may assert that the period of storage did not negatively affect the macro nutrients of the powdered sample. The low moisture content of Kunu-powdered could have discouraged the growth of microbes thereby enabling it to be in storage for six months¹⁴. On the other hand, the increase

in carbohydrate could have resulted from the degradation of polysaccharides inherent in the powders into monosaccharides like glucose and sucrose. It could also be as a result of increase in amylose content¹⁵. Since the dietary fats function to increase food palatability¹⁶, the lipid contents of the powdered coupled with its increased carbohydrate could have made the reconstituted drinks appealing to consumers. This is why the reconstituted kunu powder drink as related to the conventional mode of freshly prepared would be a good source of energy. Moreover, the powdered Kunu was observed to be free from insect pest while being kept in tightly closed glass containers for over 6 months. The drying process employed in the conversion of freshly produced Kunu into powdered form could have helped disinfested the commodities from any insect developmental stage. This is in agreement with the assertion of Ofuya and Lale 2001¹⁷ who stated that temperature has an important quantitative effect upon insect development. Appert also stated that temperature in excess of 35⁰c are lethal for certain species¹⁴. The thickness of the glass container could have also prevented any pest from chewing its way into the commodities. However, in terms of mineral composition, the reconstituted kunu powder drinks were observed to be significantly different from one another. In fact, Kunu-sorghum was observed to have the highest composition of all the minerals investigated except Calcium (76.0±16) which was observed to be relatively more in Millet (101.67±2.23) and maize (92.0±2.89). This finding is in line with the work done by Ogungbemi *et al.* 2017 on freshly made Kunu drinks (prepared from maize, millet and sorghum) but with one striking difference¹⁸. This difference is found in the mineral composition of the reconstituted kunu-sorghum powder. Our study revealed this particular powder to be richer in micronutrients compared to when freshly prepared. This could be as a result of some intrinsic chemical reactions leading to the release of more cations of medicinal benefits. Hence, the reconstituted kunu-sorghum powder could be said to be more nutritious than other kunu drinks. These micro nutrients are constituent of bones, teeth, blood, muscles, hair and nerve cells and according to Rumeza *et al.* 2006 they are very important and essential ingredients of diet required for normal metabolic activities of body tissues¹⁹.

Conclusion

It can be concluded that powdered reconstituted Kunu-zaki showed no significant different in the major nutritional, sensory evaluated properties and proven to have an elongated shelf life for more than 6months with no insect infestation during storage when compared to freshly prepared Kunu.

REFERENCES

1. Adejuyitan, J.A., Adelakun O.E, Olaniyan S.A and Popoola F.I, (2008). Evaluating the quality characteristics of kunu produced by dry-milled sorghum. African Journal of Biotechnology, 7(13): 2244-2247.
2. Ayo J.A. (1998). Effect of *C.farinosa* on the Quality of *Kunun zaki*. M.Sc. Thesis. Enugu State University. Enugu, Nigeria.
3. Adeyemi, I.A., Umar, S. (1994). Effect of method of manufacture on the quantity characteristics of Kunu-zaki a millet based beverage. Nigeria food Journal. 12: 34-40.
4. Elmahmood, A.M and J.H. Doughari, 2007. Microbial quality assessment of Kunu zaki beverages sold in griei town of Adamawa State, Nigeria. African Journal of food Science, pp: 011-015.
5. Ayo J.A., Ayo V.A., Yelmi B., Onuoha G. and Ikani M.O. (2013). Effect of preservatives on microbiological qualities of *kunu zaki*. Int. J. Agric. Sci. Res. Vol. 2(5), pp. 124-130.
6. Ikpoh, I.S., Lennox, J. A., Ekpo, I.A., Agbo, B. E., Henshaw E. E. and Udoekong, N.S. (2013). Microbial Quality Assessment Of Kunu Beverage Locally Prepared And Hawked In Calabar, Cross River State, Nigeria. Glob. J. Biodivers. Sci. Manag., 3(1): 58-61.
7. Maduegwe E.P. (1995). Assessment of Production Practices ad Evaluation of Product Characteristics of *Kunun-zaki*. Unpublished. B.Sc. Thesis, Department of Food Science and Technology, University of Agriculture, Makurdi Benue State.
8. Kordylas, J.M. (1991). Processing and Preservation of Tropical and Subtropical Foods. Published by Macmillam Education Ltd. London.
9. Ikpoh, I.S., Lennox, J. A., Ekpo, I.A., Agbo, B. E., Henshaw E. E. and Udoekong, N.S. (2013). Microbial Quality Assessment of Kunu Beverage Locally Prepared and Hawked In Calabar, Cross River State, Nigeria. Glob. J. Biodivers. Sci. Manag., 3(1): 58-61.
10. AOAC Official methods of analysis. 15th Association of Official Analysis Chemist, Washinton D.C. 2005: 774-784.
11. Pearson D. The chemical analysis of foods. 7th edition. Longman Group Ltd; 1976.
12. Larmond, E. (1977). *Laboratory methods for sensory evaluation of food*. Research Branch, Canada Dept. of Agriculture.

13. De Lima, C.P.F. (1987). Insect pests and postharvest problems in the tropics. Insect Science and its Application 8, 673-676.
14. Appert, J. (1987). The storage of food grains and seeds. Macmillan Publishers Ltd., London. 146pp.
15. Gong K.J. and Chen L.R. (2013). Characterization of carbohydrates and their metabolizing enzymes related to the eating quality of postharvest fresh waxy corn. J Food Biochem. 34:619–627
16. Antia B.S., Akpan E.J., Okon P.A. and Umoren I.U (2006). Nutritive and Anti-Nutritive Evaluation of sweet Potatoes (*Ipomoea batatas*) Leaves. Pakistan Journal of Nutrition 5(2): 166-168.
17. Ofuya T.I and Lale N.E.S. (2001). Pests of stored Cereals and Pulses in Nigeria, biology, ecology and control. Dave Collins Ltd., Akure, Nigeria.
18. Ogungbemi K Alejo A.O., Ilesanmi F.F., Ishola D.T., Afolabi A.A , and Zaka K.O. (2017), "Sensory, Shelf-Life and Nutritional Evaluation of Kunu (Nigeria Non-Alcoholic Beverage) Produced From Different Grains", International Journal of Research Studies in Agricultural Sciences (IJRSAS), vol. 3, no. 9, pp. 20-25, 2017. <http://dx.doi.org/10.20431/2454-6224.0309004>
19. Rumeza H., Zafar I., Mudassar I., Shaheena H. and Masooma R. (2006). Use of vegetales as nutritional food: role in human health. Journal of Agricultural and Biological Science. Vol.1, No.1, July 2006.

Appendix



Pictorial representation of Kunu converted to powdered form.