

Effect of Storage Conditions (Relative Humidity, Packaging materials and Time) on the Chemical properties of Maize- Soy Flour Blend

ABSTRACT

Aims:The aim of this study was to examine the effect of storage conditions on the of maize-soy flour blend

Study Design:Preliminary studies were conducted using ratio blends ranging from 70:30, 75:25, 80:20, 85:15, 90,10, 95:5 and 100.0 of maize flour to Soy flour. This was to ascertain the best blend formulation for the study. The sensory attributes show that the ratio of 85:15 Maize-soy flour blend was preferred. It was packaged in low density polyethylene (LDPE), high density polyethylene (HDPE) and storage at $30.5 \pm 3^{\circ}\text{C}$ and Relative humidity of 57 and 87% for 4 months. Analysis of Proximate composition, pH, total titratable acidity (TTA), Thiobarbuturic acid (TBA) was done on the samples at an interval of month respectively.

Result:Packaging significantly ($p > 0.05$) affected the chemical, qualities of “Soy-fermented maize” flour during storage. Moisture content, titratable acidity (TTA) and Thiobarbuturic acid (TBA) increased with storage period (9.46% – 23.5%, 0.12% - 0.21%, and 0.06 -0.12 respectively) while all other chemical, quality of the soy-fermented maize flour decreased significantly ($p > 0.05$) (pH: 5.18% – 3.45%, Protein: 15.21% – 12.18% Fat: 7.45 – 5.36, Fibre: 3.27% - 1.65%, Ash:1.12% – 0.89%, Carbohydrate: 62.97% - 56.87%.

Conclusion:The samples packaged in HDPE were more acceptable than those in other packaging materials owing to its considerable maintenance of the flour’s quality during and after storage.

Keywords: *Storage, Agidi, Maize- Soy Flour low density polyethylene, high density polyethylene*

1. INTRODUCTION

Agidi is a local West African dish (mostly in Nigeria) made from fermented maize Sorghum or millet known as *ogi*. *Ogi* is one of the popular products consumed widely in Nigeria. It is a fermented starchy mash obtained by soaking, wet milling, wet extraction (filter) and decanting of top water to obtain *ogi* [1]. *Ogi* is cooked with water to produce a semi –solid product called *Agidi* which is also known as *eko* [1]. *Agidi* could be eaten alone or with vegetables soup and/or stew as well as with moi-moi or akara (stemmed or fried bean cake) by both infants and adults. *Agidi* has added advantage over *ogi*, as it could be eaten cold or warm. It could also be prepared and kept for later use, unlike *ogi*, which should be eaten warm, thereby requiring fresh preparation. *Ogi* traditionally, the grains are soaked in water for up to three days, before wet milling and sieving to ferment. For up to three days until sour. It is then boiled as pap, or cooked into a semi –solid product called *Agidi*. Its appearance or color depends on the type of cereal used for production [2].

Earlier attempts made tend to improve the nutritional quality of these maize based “*ogi*” not much was found for *agidi* [3]. *Agidi* is quite low in protein since it is mostly composed starch. Over consumption of such product could lead to problems generally associated with protein malnutrition [4]. Due to its low protein content, soybean was added to improve the nutritional

composition and also add value to *agidi*, since it is cheap and available source of protein. Soybean is a versatile crop with many uses. Among the product are Soymilk Soy-cake, ice cream, Soybean vegetable oil. As a proteinous food, soybean is much better than any other legume in terms of protein quality. The protein content of other legumes varies from 20-25% while that of soybean is about 39% [5]. The meal is rich in mineral element and vitamins such as thiamin riboflavin and niacin.

Storage of maize-soy flour is necessary due to the tedious and cumbersome unit operation methods required for the production of the flour. Storage of the maize-soy flour for the production *agidi* was probably not done in most research articles of agidi production. This was done to ascertain the quality of the storage flour over time in production of agidi with respect to its nutritional and sensory properties as these nutrients depreciate over time.

This study is geared towards find the effects of storage on the quality of maize-soy flour blends and *Agidi* product.

2. MATERIALS AND METHODS

2.1 Procurement of Materials

Maize (*Zea mays*) and Soybean (*Glycine max*) seeds used in this study were purchased from the Teaching and Research Farm of College of Agronomy, University of Agriculture Makurdi Benue State Nigeria.

2.1.1 Preparation of Fermented Maize Flour

The fermented maize flour was prepared by the wet milling process with slight modification [6-8]. As shown in figure 1 .

2.1.2 Preparation of Soy Flour

The soy flour was prepared according to the method reported by [9, 10] with slight modification. As shown in figure 2 . The flour was stored in a refrigerator (4°C) until used.

2.1.3 Preparation of Soy-Agidi

Agidi was prepared according to the method reported by[11] with slight modification. As shown in figure 3 .




Fig1: Flow Chart for the Preparation of Ogi flour (Fermented Maize Flour)

Source: Osungbaro (1998) modified.




Fig 2: Flow Chart for the Preparation of Soy Flour

Source: Amadou, *et al.*, (2009) modified.




Fig 3: Flow Chart for Production of Soy – Agidi

Source: Akpapunam *et al.*, (1997) – modified

2.1.4 Storage Studies

The samples (85:15 maize-soy flour blend) were packed in low density polyethylene film and high-density polyethylene film then stored in two dissectors with relative humidity of 80% and 60% and place in a room at ambient temperature (32 ± 2) for 24 weeks. Sample were withdrawn at four (4) weeks interval to check for chemical analysis.

2.2 Proximate Composition

The protein, moisture, fat, fibre, ash, carbohydrate, pH and titratable acidity were determined according to [12].

2.3 Statistical Analysis

All analyses were carried out in triplicate unless otherwise stated. Statistical significance was established using one-way analysis of variance (ANOVA), and data were reported as the mean standard deviation. Mean comparison and separation was done using Fisher's Least Significant Difference test (LSD) at $p \leq 0.05$. ($P < 0.05$). Statistical analysis was carried out using the SPSS 20 statistical package.

3. RESULTS AND DISCUSSION

3.1 Discussion

3.1.1 Effect storage on the protein quality of soy supplemented maize flour blend

The results for protein for fresh and storage of maize-soy flour are shown in table 1. The protein content decreased significantly ($p < 0.05$) for samples in Low density polyethylene across the four months (15.70 - 13.16), high density polyethylene (15.56 - 13.44) and no package (15.56 - 12.87) for Relative humidity of 57% and there was no significant difference ($p > 0.05$) for samples between packages (Table 1). Also in Relative humidity of 82% there was significant difference for samples in Low density polyethylene, high density polyethylene and no package ($p < 0.05$) as show in Table 1. But there was no significant difference for Samples between packages. There was a decrease in crude protein content for all with sample without package materials recorded the lowest values for both Relative humidity of 57% and 82% respectively. Similar result have been reported by other worker [13].

175 **Table 1: Effect of Storage Conditions (Relative Humidity, Packaging material and Time) on the Crude Protein of Maize- Soy**
176 **Flour Blend**

Relative Humidity	Packaging	Storage Time (in Months)					177
		0	1	2	3	4	LSD 178
57	LDPE	15.61 ^a _a ±0.03	15.70 ^a _a ±0.03	15.53 ^a _a ±0.09	14.76 ^b _a ±0.15	13.16 ^c _a ±0.08	0.56
	HDPE	15.61 ^a _a ±0.03	15.56 ^a _a ±0.14	15.44 ^a _a ±0.06	14.79 ^b _c ±0.07	13.44 ^c _a ±0.48	179 0.56
	No Packaging	15.61 ^a _a ±0.03	15.56 ^a _a ±0.08	15.57 ^a _a ±0.18	14.68 ^b _a ±0.04	12.87 ^c _a ±0.26	0.180
82	LDPE	15.61 ^a _a ±0.03	15.55 ^a _a ±0.07	15.57 ^a _a ±0.08	14.76 ^b _a ±0.09	13.54 ^c _a ±0.12	181 0.56
	HDPE	15.61 ^a _a ±0.03	15.52 ^a _a ±0.16	15.64 ^a _a ±0.08	14.73 ^b _a ±0.11	13.33 ^c _a ±0.67	182 0.56
	No Packaging	15.61 ^a _a ±0.03	15.56 ^a _a ±0.08	15.57 ^a _a ±0.18	14.68 ^b _a ±0.04	12.87 ^c _b ±0.26	0.183
LSD		0.65	0.65	0.65	0.65	0.65	184
							185
							186

187 Key:LDPE: Low density polyethylene, HDPE: High density polyethylene ,Superscript: Separation of mean for months Subscript: Separation of means for
188 packaging materials

190 3.1.2. Effect storage on the moisture content of soy supplemented maize flour blend

191 The result for moisture for fresh and storage of maize-soy flour is shown in table 2. The
192 moisture content increased significantly ($p < 0.05$) as the storage period increased irrespective
193 of the packaging material or the relative humidity. Moisture content was highest in without
194 packaging Material for both Relative humidity of 57% and 82% (9.64 -17.46 and 9.64-23.75) and
195 lowest in high density poly ethylene film (9.60-15.56 and 9.56 -15.59) during the 4 months of
196 storage at ambient condition (Table 2). The increase in the percentage moisture content of
197 stored flour can be attributed to the hygroscopic properties of the flour [14]. and might be due to
198 the fact that at a high humidity, the vapour pressure may have increased which aids water
199 absorption into the samples[15]. Polyethylene films generally have good barrier against
200 moisture [16], but low density polyethylene had higher water vapour permeability compared with
201 high density polyethylene. The result agrees with the earlier findings of [17], who observed
202 higher moisture in low density polyethylene than in high density polyethylene during the storage
203 of African Breadfruit seed flour at room temperature for 12 weeks and also findings of[18], who
204 also found higher moisture in low density polyethylene than in high density polyethylene during
205 the storage of pupuru for 24 weeks.

Table 2: Effect of Storage Conditions (Relative Humidity, Packaging material and Time) on the Moisture of Maize-Soy Flour

Blend

Relative Humidity	Packaging	Storage Time (in Months)					LSD
		0	1	2	3	4	
57	LDPE	9.61 ^d _a ±0.16	9.46 ^d _a ±0.03	10.89 ^c _a ±0.15	13.50 ^b _a ±0.14	16.33 ^a _b ±0.07	0.92
	HDPE	9.61 ^d _a ±0.16	9.60 ^d _a ±0.14	11.72 ^c _a ±0.05	13.63 ^b _a ±0.09	15.56 ^a _c ±0.59	0.92
	No Packaging	9.61 ^d _a ±0.16	9.64 ^d _a ±0.21	11.66 ^c _a ±0.06	14.27 ^b _a ±0.18	17.46 ^a _a ±0.35	0.92
82	LDPE	9.61 ^d _a ±0.16	9.57 ^d _a ±0.16	11.10 ^c _b ±0.59	13.60 ^b _b ±0.06	16.18 ^a _b ±0.43	0.92
	HDPE	9.61 ^d _a ±0.16	9.56 ^d _a ±0.23	12.77 ^c _a ±0.06	13.19 ^b _c ±0.70	15.59 ^a _c ±0.59	0.92
	No Packaging	9.61 ^d _a ±0.16	9.64 ^d _a ±0.23	12.77 ^c _a ±0.23	18.16 ^b _a ±0.54	23.75 ^a _a ±0.49	0.92
LSD		1.06	1.06	1.06	1.06	1.06	

211 Key: LDPE: Low density polyethylene, HDPE: High density polyethylene, Superscript: Separation of mean for months Subscript: Separation of means for
212 packaging materials

3.1.3 Effect storage on the fat content of soy supplemented maize flour blend

The results for crude fat for fresh and storage of maize-soy flour are shown in table 3. There was a progressive decrease in the fat content of all the samples during storage at ambient conditions. The highest decrease in fat was seen in samples without package in both Relative humidity of 57% and 82% as seen in Table 5. And the lowest decrease was found in samples in High density polyethylene. The result agrees with the earlier findings of [19], who also found a steady decrease in fat during storage of cassava chips, cassava flour, yam chips and yam flour for three months. The decrease may be attributed to the lipolytic activity of enzymes i.e. lipase and lipoxidase [20].

230 Table 3: Effect of Storage Conditions (Relative Humidity, Packaging material and Time) on the Crude Fat of
 231 Maize-Soy Flour Blend

Relative Humidity	Packaging	Storage Time (in Months)					LSD
		0	1	2	3	4	
57	LDPE	7.55 ^a _a ±0.08	7.53 ^a _a ±0.03	7.21 ^b _a ±0.2	6.76 ^c _a ±0.08	6.58 ^c _a ±0.03	0.29
	HDPE	7.55 ^a _a ±0.08	7.56 ^a _a ±0.08	7.16 ^b _a ±0.06	6.65 ^c _a ±0.08	6.37 ^c _a ±0.22	0.29
	No Packaging	7.55 ^a _a ±0.08	7.51 ^a _a ±0.02	7.17 ^b _a ±0.06	6.67 ^c _a ±0.07	6.38 ^d _a ±0.19	0.29
82		7.55 ^a _a ±0.08	7.49 ^a _a ±0.01	7.20 ^b _a ±0.11	6.79 ^c _a ±0.03	6.64 ^c _a ±0.14	0.29
	LDPE	7.55 ^a _a ±0.08	7.56 ^a _a ±0.07	7.25 ^b _a ±0.05	6.63 ^c _a ±0.18	6.61 ^c _a ±0.19	0.29
	HDPE	7.55 ^a _a ±0.08	7.45 ^a _a ±0.06	7.16 ^b _a ±0.08	5.69 ^c _b ±0.08	5.00 ^d _b ±0.01	0.29
	No Packaging	7.55 ^a _a ±0.08	7.53 ^a _a ±0.03	7.21 ^b _a ±0.2	6.76 ^c _a ±0.08	6.58 ^c _a ±0.03	0.29
LSD		0.33	0.33	0.33	0.33	0.33	

232 Key: LDPE: Low density polyethylene, HDPE: High density polyethylene ,Superscript: Separation of mean for months Subscript: Separation of means for
 233 packaging materials

3.1.4 Effect storage on the fiber content of soy supplemented maize flour blend

The results for crude fiber for fresh and storage of maize-soy flour are shown in table 4. There was significant difference ($p < 0.05$) for samples in Low density polyethylene across the four months, high density polyethylene, and no package for Relative humidity of 57% and 82% there was no significant difference ($p > 0.05$) for samples between packages. There was a steady decrease in fiber content with samples without packing material having the decreases in both Relative humidity of 57% and 82%. (3.32- 1.86 and 3.23- 1.62). While sample in low density polyethylene had the lowest decrease for relative humidity 57 (3.29 - 2.03) and samples in high density polyethylene had the highest decrease for relative humidity 82 (3.27- 1.86) (Table 4).

These results were contrary to the result obtained by [19], who found a steady increase in fiber during storage of cassava chips, cassava flour, yam chips and yam flour for three months. But was in agreement with [21] who observed fiber decreases during storage of Soup Thickener *Brachystegia enrycoma* (Achi) for 12 weeks.

248 Table 4: Effect of Storage Conditions (Relative Humidity, Packaging material and Time) on the Crude Fiber of Maize-Soy Flour Blend

Relative Humidity	Packaging	Storage Time (in Months)					LSD
		0	1	2	3	4	
57	LDPE	3.30 ^a _a ±0.06	3.29 ^a _a ±0.02	2.54 ^b _a ±0.07	2.16 ^c _a ±0.06	2.03 ^c _a ±0.11	0.30
	HDPE	3.30 ^a _a ±0.06	3.27 ^a _a ±0.04	2.29 ^b _a ±0.01	1.95 ^c _a ±0.12	1.89 ^c _a ±0.19	0.30
	No Packaging	3.30 ^a _a ±0.06	3.32 ^a _a ±0.04	2.38 ^b _a ±0.18	1.89 ^c _a ±0.06	1.86 ^c _a ±0.15	0.30
82	LDPE	3.30 ^a _a ±0.06	3.27 ^a _a ±0.02	2.43 ^b _a ±0.04	1.94 ^c _a ±0.06	1.77 ^c _a ±0.16	0.30
	HDPE	3.30 ^a _a ±0.06	3.27 ^a _a ±0.01	2.45 ^b _a ±0.04	1.91 ^c _a ±0.15	1.86 ^c _a ±0.27	0.30
	No Packaging	3.30 ^a _a ±0.06	3.23 ^a _a ±0.04	2.27 ^b _a ±0.08	1.75 ^c _a ±0.00	1.65 ^c _a ±0.14	0.30
LSD		0.35	0.35	0.35	0.35	0.35	

249 Key: LDPE: Low density polyethylene, HDPE: High density polyethylene, Superscript: Separation of mean for months Subscript: Separation of means for
 250 packaging material

3.1.5 Effect storage on the ash content of soy supplemented maize flour blend

The results for Ash for fresh and storage of maize-soy flour are shown in table 5. There was a significant difference ($p < 0.05$) for samples in Low density polyethylene across the four months, high density polyethylene, and no package for Relative humidity of 57% and 82% there was no significant difference ($p > 0.05$) for samples between packages. There was a steady decrease in ash with samples without packing material having the highest decreases in both Relative humidity of 57% and 82% (1.12-0.99 and 1.13-0.95) and lowest decrease is was recorded in Low density polyethylene for both relative humidity (1.13-1.04 and 1.13-1.05). The findings agreed with result obtained by [21].

268 Table 5: Effect of Storage Conditions (Relative Humidity, Packaging material and Time) on the Ash of Maize-Soy Flour Blend

Relative Humidity	Packaging	Storage Time (in Months)					LSD
		0	1	2	3	4	
57	LDPE	1.14 ^a _a ±0.01	1.13 ^a _a ±0.06	1.07 ^a _a ±0.08	1.06 ^a _b ±0.09	1.04 ^a _a ±0.92	0.23
	HDPE	1.14 ^a _a ±0.01	1.14 ^a _a ±0.01	1.06 ^a _a ±0.08	1.09 ^a _a ±0.00	1.00 ^a _a ±0.01	0.23
	No Packaging	1.14 ^a _a ±0.01	1.12 ^a _a ±0.42	1.09 ^a _a ±0.21	1.01 ^a _a ±0.01	0.99 ^a _a ±0.01	0.23
82	LDPE	1.14 ^a _a ±0.01	1.13 ^a _a ±0.02	1.28 ^a _a ±0.24	0.99 ^a _a ±0.04	1.05 ^a _a ±0.14	0.23
	HDPE	1.14 ^a _a ±0.01	1.14 ^a _a ±0.04	1.15 ^a _a ±0.07	1.00 ^a _a ±0.01	0.88 ^b _a ±0.17	0.23
	No Packaging	1.14 ^a _a ±0.01	1.13 ^a _a ±0.02	1.12 ^a _a ±0.16	1.03 ^a _a ±0.00	0.98 ^a _a ±0.28	0.23
LSD		0.26	0.26	0.26	0.26	0.26	

269 Key: LDPE: Low density polyethylene, HDPE: High density polyethylene ,Superscript: Separation of mean for months Subscript: Separation of means for
 270 packaging material

3.1.6 Effect storage on the carbohydrate content of soy supplemented maize flour blend

The results for carbohydrate for fresh and storage of maize-soy flour are shown in table 6. There was a significant difference ($p < 0.05$) for samples in Low density polyethylene across the four months, high density polyethylene, and no package for Relative humidity of 57% and 82%, there was also significant difference ($p < 0.05$) for samples between packages. There a steady decrease is carbohydrate content with sample with no packaging materials have the lowest decrease for both relative humidity 57 and 82 (62.86-60.42 and 62.99-56.87) and the highest decrease for relative 57 was found in samples in Low density polyethylene (62.9 -61.51) while the highest decrease for Relative humidity of 82% was observed in high density polyethylene (62.96 – 58.87) ((Table 8). The result agrees with the earlier findings of [19], who also found a steady was a decrease in the carbohydrate content of the samples during storage during storage of cassava chips, cassava flour, yam chips and yam flour for three months, which was contrary to the report of [22] who got an increase after the storage of yam chips and flour. Carbohydrate content of the samples might have decreased because of its utilization for growth by the microorganisms present [19].

287 Table 6: Effect of Storage Conditions (Relative Humidity, Packaging material and Time) on the carbohydrate of Maize-Soy Flour
 288 Blend

Relative Humidity	Packaging	Storage Time (in Months)					LSD
		0	1	2	3	4	
57	LDPE	62.97 ^a ±0.06	62.9 ^a ±0.014	62.57 ^a ±0.07	61.91 ^b ±0.09	61.51 ^b ±0.05	0.80
	HDPE	62.97 ^a ±0.06	62.87 ^a ±0.13	62.31 ^a ±0.01	61.96 ^b ±0.17	60.92 ^c ±0.67	0.80
	No Packaging	62.97 ^a ±0.06	62.86 ^a ±0.06	62.08 ^a ±0.13	61.52 ^b ±0.03	60.42 ^c ±0.13	0.80
82	LDPE	62.97 ^a ±0.06	63.01 ^a ±0.11	61.92 ^b ±0.11	61.86 ^b ±0.11	60.83 ^c ±0.25	0.80
	HDPE	62.97 ^a ±0.06	62.96 ^a ±0.92	62.42 ^a ±0.65	62.59 ^a ±0.21	61.67 ^b ±0.25	0.80
	No Packaging	62.97 ^a ±0.06	62.99 ^a ±0.01	61.19 ^b ±0.26	58.80 ^c ±0.66	56.87 ^d ±0.47	0.80
LSD		0.92	0.92	0.92	0.92	0.92	

289 Key: LDPE: Low density polyethylene, HDPE: High density polyethylene ,Superscript: Separation of mean for months Subscript: Separation of means for
 290 packaging material

3.1.7 Effect storage on the pH content of soy supplemented maize flour blend

The pH values of the freshly sample and stored values are shown in table 7.

There was a steady decrease in pH value during the storage months (samples became more acidic). The lowest decrease for pH in Relative humidity of 57% was recorded in samples in Low de Low density polyethylene and highest in Sample No packaging materials. The lowest decrease for pH in Relative humidity of 82% was recorded in samples in No packaging materials and highest in Sample high density polyethylene. These finding are in agreement with earlier [18], who also found higher pH value in low density polyethylene than in high density polyethylene during the storage of pupuru for 24 weeks.

For samples with no low density polyethylene, the samples at Relative humidity of 57% was recorded as samples with the Lowest pH values after storage while samples storage under Relative humidity of 82% had the higher pH values.

For samples with no high-density polyethylene, the samples at Relative humidity of 82% was recorded as samples with the Lowest pH values after storage while samples storage under Relative humidity of 57% had the higher pH values.

For samples with no Packaging material, the samples at Relative humidity of 82% was recorded as samples with the Lowest pH values after storage while samples storage under Relative humidity of 57% had the higher pH values .

In general the was a steady decreases in pH value in all samples the finding is in agreement with[19], who also found a steady decrease in Ph value during storage of cassava chips, cassava flour, yam chips and yam flour for three months. The pH , an indication of microbial proliferation and loads The pH observed value could help in control of microbial load in the flour [23].

314 Table 7: Effect of Storage Conditions (Relative Humidity, Packaging material and Time) on the pH of Maize-Soy Flour Blend

Relative Humidity	Packaging	Storage Time (in Months)					LSD
		0	1	2	3	4	
57	LDPE	5.21 ^a _a ±0.01	5.05 ^a _a ±0.07	4.86 ^{ab} _c ±0.02	3.45 ^c _c ±0.00	3.45 ^c _c ±0.01	0.56
	HDPE	5.21 ^a _a ±0.01	5.18 ^a _a ±0.03	5.14 ^a _a ±0.07	4.15 ^b _b ±0.12	3.80 ^c _b ±0.09	0.56
	No Packaging	5.21 ^a _a ±0.01	5.14 ^a _a ±0.35	5.01 ^b _b ±0.10	4.26 ^b _a ±0.08	4.26 ^b _a ±0.03	0.56
82	LDPE	5.21 ^a _a ±0.01	5.13 ^a _a ±0.21	4.72 ^{ab} _c ±0.01	3.81 ^c _b ±0.06	3.71 ^c _a ±0.01	0.56
	HDPE	5.21 ^a _a ±0.01	5.20 ^a _a ±0.31	5.07 ^a _a ±0.14	3.99 ^b _c ±0.01	3.64 ^b _b ±0.12	0.56
	No Packaging	5.21 ^a _a ±0.01	5.18 ^a _a ±0.01	4.90 ^a _b ±0.02	3.75 ^b _c ±0.35	3.66 ^b _b ±0.07	0.56
LSD		0.07	0.07	0.07	0.07	0.07	

315 Key: LDPE: Low density polyethylene, HDPE: High density polyethylene, Superscript: Separation of mean for months Subscript:

316 Separation of means for packaging material

3.1.8 Effect storage on the Titratable Acidity content of soy supplemented maize flour blend.

The Titratable Acidity values of the freshly sample and stored values are shown in table 8.

There was a steady increase in Titratable Acidity value during the storage months (samples became more acidic). The lowest increase for Titratable Acidity in Relative humidity of 57% was recorded in samples in No packaging materials and highest increase in Samples high density polyethylene. The lowest increase for Titratable Acidity in Relative humidity of 82% was recorded in samples in high density polyethylene and highest increase was found in Samples with No packaging materials. These finding are in agreement with earlier[18], who also found higher Titratable Acidity value in low density polyethylene than in high density polyethylene during the storage of pupuru for 24 weeks.

For samples with no low-density polyethylene, the samples at Relative humidity of 57% was recorded as samples with the Lowest Titratable Acidity values after storage while samples storage under Relative humidity of 82% had the higher Titratable Acidity values.

For samples with no high-density polyethylene, the samples at Relative humidity of 82% was recorded as samples with the Lowest Titratable Acidity values after storage while samples storage under Relative humidity of 57% had the higher Titratable Acidity values.

For samples with no Packaging material, the samples at Relative humidity of 82% was recorded as samples with the Lowest Titratable Acidity values after storage while samples storage under Relative humidity of 57% had the higher Titratable Acidity values .

There was an increase in Titratable Acidity during storage irrespective of packaging materials. The increase in Titratable Acidity with storage period was also observed by [23] who found that Titratable acidity increase during storage of flours from soaked, malted and their blend of millet grains (*Pennisetum glaucum*) for 90 days.

340 Table 8: Effect of Storage Conditions (Relative Humidity, Packaging material and Time) on the Titrable Acidity of Maize-Soy Flour
341 Blend

Relative Humidity	Packaging	Storage Time (in Months)					LSD
		0	1	2	3	4	
57	LDPE	0.12 ^c _a ±0.01	0.12 ^c _b ±0.00	0.13 ^{cb} _b ±0.01	0.15 ^b _c ±0.02	0.18±0.01	0.02
	HDPE	0.12 ^c _a ±0.01	0.13 ^c _a ±0.01	0.13 ^c _b ±0.01	0.17 ^{ab} _b ±0.01	0.19 ^a _a ±0.02	0.02
	No Packaging	0.12 ^c _a ±0.01	0.11 ^c _c ±0.02	0.14 ^b _a ±0.01	0.18 ^a _a ±0.00	0.19 ^a _a ±0.01	0.02
82	LDPE	0.12 ^c _a ±0.01	0.13 ^b _a ±0.02	0.14 ^b _a ±0.03	0.16 ^a _b ±0.01	0.18 ^a _c ±0.01	0.02
	HDPE	0.12 ^c _a ±0.01	0.13 ^c _a ±0.00	0.13 ^c _a ±0.01	0.16 ^b _b ±0.01	0.19 ^a _b ±0.01	0.02
	No Packaging	0.12 ^c _a ±0.01	0.11 ^b _b ±0.01	0.14 ^b _a ±0.12	0.19 ^a _a ±0.00	0.21 ^a _a ±0.02	0.02
LSD		0.01	0.01	0.01	0.01	0.01	

342 Key: LDPE: Low density polyethylene, HDPE: High density polyethylene ,Superscript: Separation of mean for months Subscript:
343 Separation of means for packaging material

4. CONCLUSION

The result of study showed that Increase in moisture content was directly proportional to the increase in storage time, conversely a decrease in protein, carbohydrate, ash, fibre and fat content was observed with increased storage time.

The pH of the samples decreased with an increase in the storage time. An inverse relationship was observed for Titratable acidity.

REFERENCE

1. UMOH, V. and M. FIELDS, *Fermentation of corn for Nigerian agidi*. Journal of Food Science, 1981. **46**(3): p. 903-905.
2. Banigo, E. *Ogi: a Nigerian fermented cereal food*. in *Symposium on Indigenous Fermented Foods*. Bangkok, Thailand. 1977.
3. Eka, O., *Chemical evaluation of nutritive value of soya paps and porridges, the Nigerian weaning foods*. Food Chemistry, 1978. **3**(3): p. 199-206.
4. Grover, Z. and L.C. Ee, *Protein energy malnutrition*. Pediatric Clinics, 2009. **56**(5): p. 1055-1068.
5. Smith, A.K., *Chemical composition of the [soybeans] seed*. Soybeans: chemistry and technology, 1978.
6. Adeyemi, I. and O. Beckley, *Effect of period of maize fermentation and souring on chemical properties and amylograph pasting viscosity of ogi*. Journal of Cereal Science, 1986. **4**(4): p. 353-360.
7. ADENIJI, A.O. and N.N. POTTER, *Properties of OGI powders made from normal, fortified and Opaque-2 corn*. Journal of Food Science, 1978. **43**(5): p. 1571-1574.
8. Taiwo, O., *Physical and nutritive properties of fermented cereal foods*. African Journal of Food Science, 2009. **3**(2): p. 023-027.
9. Amadou, I., et al., *Fermented soybean products: some methods, antioxidants compound extraction and their scavenging activity*. Asian J. Biochem, 2009. **4**(3): p. 68-76.
10. Iwe, M., *The science and technology of soybean*. Rojoint Communication Services Ltd, Uwani, Enugu, Nigeria, 2003: p. 84-110.
11. Akpapunam, M., G. Badifu, and E. Etokudo, *Production and Quality Characteristics of Nigerian Agidi supplemented with soy flour*. Journal of food science and technology, 1997. **34**(2): p. 143-145.
12. Latimer, G.W., *Official methods of analysis of AOAC International*. 2012: AOAC international.
13. Garima, M. and M.A. Anand, *Studies on Effect of HDPE and LDPE on Storage Stability of Weaning Food prepared from Pulse, Banana and Pineapple Pomace*. International Journal of Innovation and Applied Studies, 2014. **7**(2): p. 501.
14. Butt, M.S., et al., *Effect of moisture and packaging on the shelf life of wheat flour*. Internet Journal of Food Safety, 2004. **4**(1): p. 1-6.

15. Akindahunsi, A. and G. Oboh. *Effect of changes in Relative Humidity on the Storage Stability of Micro-Fungi Fermented Gari*. in *Proceedings of 24th Annual Conference of Nigerian Institute of Food Science and Technology*. 2000.
16. Ukpabi, U., R. Omodamiro, and E. Oti. *Feasibility of using sealed polyethylene film in prolonged storage of gari*. in *Proc. 29th Nigerian Institute of Food Science and Technology Annual Conference/AGM*. 2005.
17. Fasasi, O. *Effect of different packaging materials on the chemical composition of African breadfruit seed (Treculia africana) Flour during storage at room temperature*. in *Proceed 27th Annual Conf Nigerian Institute of Food Science and Technol*. 2003.
18. Daramola, O.A., et al., *Effects of packaging material on the quality of pupuru flour during storage*. *African Journal of Food Science*, 2010. **4**(5): p. 258-263.
19. Oyeyiola, G., et al., *A comparative study on the microbiological and nutritional properties of stored chips and flours*. *World Journal of Biological Research*, 2014. **6**(02).
20. Agrahar-Murugkar, D. and K. Jha, *Influence of storage and packaging conditions on the quality of soy flour from sprouted soybean*. *Journal of food science and technology*, 2011. **48**(3): p. 325-328.
21. Nwosu, J., et al., *The effect of storage conditions on the proximate and rheological properties of soup thickener Brachystegia enrycoma (Achi)*. *Report and Opinion*, 2011. **3**(5): p. 52-58.
22. Ojokoh, A. and R. Gabriel, *A comparative study on the storage of yam chips (gbodo) and yam flour (elubo)*. *African Journal of Biotechnology*, 2010. **9**(21): p. 3175-3177.
23. Ogori, A.F., Jatua, M. K., Adamu, L. And Utim, M.S, *Effect of semi-densed polyethylene storage on organoleptic and chemical characteristics of flours from soaked ,malted and their blend of millet grains (pennesitum glacum)*. *American Journal of Biological, Chemical and Pharmaceutical Sciences*, 2013. **1**(7): p. 81-89.