

Effect of ZnO Nanoparticles on Protein Fractions of Some Plant and Animal Protein Sources Using CNCPS Method

ABSTRACT

Aims: This experiment was conducted with the aim of investigating effect of zn nanoparticles on fractions protein of some plant and animal protein sources using cncps method.

Study design: After preparing the desired food items, the protein fractions of feed samples were determined by Licitra et al., 1996.

Methodology: After testing, the amount of non-protein nitrogen (fraction A) in soybean meal, cottonseed meal and rapeseed meal was 8.52, 6.33 and 4.55 %, and in poultry offal meal, fish meal and blood meal in slaughterhouses were 10.38, 13.63 and 16.08% of crude protein, the amount of B1 in soybean meal, cottonseed meal and rapeseed meal was 2.30, 3.32 and 13.68 % respectively, and in poultry offal meal, fish meal and blood meal, in slaughterhouses were 3.45, 7.44 and 7.16 % of crude protein respectively , the amount of B2 in soybean meal, cottonseed meal and rapeseed meal was 80.49, 77.50 and 68.40%, in poultry offal meal, fish meal and blood meal, respectively, 66.36, 55.03 and 61.66 % respectively, the amount of B3 for soybean meal, cottonseed meal and rapeseed meal was 6.24, 2.63 and 9.11 %, respectively, and poultry offal meal, fish meal and blood meal in slaughterhouses were 7.50, 6.74 and 11.91 % of crude protein respectively , the protein C portion assumed in the rumen's indissoluble CNCPS system in soybean meal, cottonseed meal, and rapeseed meal was 2.45, 9.92, and 4.77 %, respectively, in poultry offal meal, fish meal and blood meal, respectively, 12.21 , 17.16 and 3.18 % of crude protein

Results: The results indicate that the use of CNCPS and NRC data for portion fractions of different feeds cannot be considered absolutely, and domestic research and results should be used to extract samples from different regions and different growth conditions, so that the dietary regimens With these foods, it's real and more balanced and with less waste of nutrients.

Keywords: CNCPS, Fractions protein, nZnO, Protein sources.

1. INTRODUCTION

The CNCPS (Cornell Net Carbohydrate and Protein System) system is a semi-mechanical approach that evaluates the rate of degradation of feed in the rumen, passes through the rumen's undigested material, as well as the amounts of energy and protein metabolism used for ruminant tract [2]. The protein fractions in the CNCPS system is A, B1, B2, B3, and C. Non-protein nitrogen (fraction A) is a part of the crude protein that is dissolved quickly (zero time) in the rumen fluid, and its decomposition rate is assumed to be unlimited. Fraction A is chemically part of the crude protein that is dissolved in borate phosphate buffer [1]. The

degradation rate in the rumen is rapid and directly into the rumen ammonia tank [3]. B1 is the percentage of total crude protein dissolved in borate phosphate buffer and precipitated with Trichloroacetic acid (TCA; TCAA; also known as trichloroethanoic acid), and its rate of degradation in the rumen is rapid, the rate of degradation of this part in the rumen is about 200-300 percent, and its degradation in the intestine 100 percent. Fraction B2 is a protein that partly breaks down into the rumen and is calculated as the difference in the total amount of protein A, B1, B3 and C from the crude protein. The fate of this sector is rapidly degradation and the speed of passing through the rumen depends. The digestion of the amino acids of B1 and B2 is 100% and 80% of the B3 sectors. The protein part B3 contains prolamins and fermented proteins, which are found to be extremely low in most of the feed, especially protein. Part C is a non-soluble protein in acidic detergent that is assumed to be in the Cornell net carbohydrate and protein system in the rumen and has a direct correlation with the thermal damage of protein and indigestible protein [4]. In CNCPS, it is assumed that protein C does not decompose in the rumen, this part has a direct relation to heat damage [5]. Therefore, proper and controlled temperature is very important during thermal processes. Increasing cell biodegradability results in the release of proteins bound to the cell wall and decreases the protein C fraction [6]. In CNCPS, the reduction of A and B1 and the increase of protein B2 and B3 are associated with a decrease in protein degradation in the rumen and an increase in RUP [3], which, in the absence of a negative effect on RUP degradation the intestine can have a positive effect on the production and reproductive performance of lactating cows [7].

Table 1: Protein fraction of feed content

Fraction	Grouping*	Style abbreviation	Enzymatic decomposition	Method of estimation and definition
Non-protein nitrogen	A	NPN	impractical	Insoluble and dissolved
True protein	-	TP	-	It is precipitated with trichloroacetic acid
Soluble true protein	B1	BSP	rapid	Buffer solution, but insoluble
Insoluble protein	-	IP	-	Insoluble in buffer
Soluble protein in neutral detergent	B2	IP-NDIP	Variable	The difference between the insoluble in the buffer and the insoluble in the neutral detergent
Insoluble Protein in neutral detergent but soluble in acid detergent	B3	NDIP-ADIP	Variable and Slow	Insoluble in neutral detergent but soluble in acid detergent
Insoluble in acid detergent and attached	C	ADIP or ADIN	Indigestible	Protein has a thermal damage seen and attached to lignin

* Based on Van Soest [8]

This experiment was conducted with the aim of investigating effect of ZnO nanoparticles on fractions protein of some plant and animal protein sources using CNCPS method.

2. MATERIALS AND METHODS

2.1 Determine Different Parts of Nitrogen

The protein fractions of feed samples by [1] as follows were determined.

2.2 NPN or A (Non-Protein Nitrogen):

Initially, 0.5 g of the desired dry feed sample was weighed and spilled into the 125 ml Erlenmeyer. Then, 50 ml of distilled water, and then, 8 ml of sodium tungstate 10% solution was added and placed until the Erlenmeyer remained at 20-25 °C for 30 minutes. After this period of pH, add 10 ml of sulfuric acid to a pH of 0.5 to 2 M and was restarted until the Erlenmeyer remains at room temperature overnight. Pull the filter paper and place it in a cone funnel. First, the paper was wetted, then the filter was done and then left, washed twice with distilled water, and the paper was transferred to the Kjeldahl flask and the remaining nitrogen was estimated to deduct the remaining nitrogen from the total nitrogen feed and NPN fraction calculated.

2.3 BSN (Nitrogen Soluble in Buffer):

Initially, 0.5 g of the desired dry feed sample was weighed and poured into an Erlenmeyer 125 ml and then 50 ml of borate phosphate buffer and then, 1 ml of sodium azide solution was added to it and laid to Erlenmeyer stays at room temperature for 3 hours. After this time, the filter paper was passed and the filter was done and the remainder was washed with 250 ml of distilled water and Nitrogen was estimated in the remainder of the Kjeldahl, which is the same protein as the insoluble, and the soluble protein can also be distinguished by difference. The total crude protein was calculated and the real protein with NPN fraction (A) was obtained from BSN.

$B1 \text{ or } TP = BSN - A$

2.3 ADIN or C (Insoluble Nitrogen in Acid Detergent):

An acid insoluble fiber method was used to dissolve acidic solution in ANKOM method [9] using filter paper 541. The nitrogen present in the sediment is characterized by ADIN, using the Kjeldahl method.

2.4 NDIN (Insoluble Nitrogen in Neutral Detergent):

Using the method of determining the insoluble fiber in neutral detergent in the incubator [9], the amount of nitrogen present in the precipitate was determined on the filter paper as insoluble nitrogen in neutral detergent by the Kjeldahl method.

$A = \text{NPN}$

$B1 = \text{BSN} - \text{NPN}$

$B2 = \text{CP} - (A + B1 + B3 + C)$

$B3 = \text{NDIN} - \text{ADIN}$

$C = \text{ADIN}$

3. RESULTS AND DISCUSSION

3.1 Chemical Composition

The chemical compositions of test feed are presented in Table 2. Blood meal content had higher percentage of protein than any of the other plant and animal protein. The maximum

99 amount of crude fat 31.3% for poultry offal meal (POM) and highest ash content of 20% was
 100 observed for fish meal (FM). Highest of NDF and ADF (70.6% and 58.4%) for cottonseed
 101 meal (CM) and the lowest NDF and ADF were obtained 45.7 and 33.3% for soybean meal
 102 (SM), respectively.
 103

Table 2. Chemical composition of some plant and animal protein sources

Protein sources	DM	CP	EE	Ash	NDF	ADF
Plant						
Soybean meal	92.4	50	1.6	6.1	45.7	33.3
Rapeseed meal	91.4	37	1.2	8	51.5	46.1
Cottonseed meal	93	24	1.4	4.7	70.6	58.4
Animal						
Poultry offal meal	94.4	55	31.3	7.3	48.9	34.8
Fish meal	93.6	50	18.1	20	61.2	40.6
Blood meal	70.6	59	1.6	5	55.3	33.4

*DM = dry matter (percent), CP = crude protein (%DM), EE= crude fat (%DM), Ash = ash (%DM) NDF = Neutral detergent fiber (%), ADF= Acid detergent fiber (%)

104 3.2 Protein Fractions

105

106 The results of various nitrogen fractions based on the CNCPS method in various protein
 107 sources are shown in Table 3. The amount of non-protein nitrogen (part A) in soybean meal,
 108 cottonseed meal and rapeseed meal was 8.52, 6.33 and 4.05 %, and in poultry offal meal,
 109 fish meal and blood meal in slaughterhouses were 10.38, 13.63 and 16.08 % of crude
 110 protein (Table 3). The highest amount of the A part was related to blood meal and the lowest
 111 was related to rapeseed meal ($p < 0.05$), which was different from the reported amounts by
 112 [10], probably due to the method used for measuring non-protein nitrogen, location Crop
 113 cultivation, harvesting method, drying and storage of feed, as well as the type of protein
 114 precipitators in different experiments. Therefore, when using blood meal in order to regulate
 115 the diet due to its high non-protein nitrogen, sufficient amount of energy should be provided
 116 to facilitate the synthesis of microbial protein and to use the NPN fraction well. The
 117 measured values of the true soluble protein (Fraction B1) in soybean meal, cottonseed meal
 118 and rapeseed meal were 2.30, 3.32 and 13.68 % respectively, and in poultry offal meal, fish
 119 meal and blood meal in slaughterhouses were 3.45, 7.44 and 7.16 % of crude protein. There
 120 was a significant difference between the mean of fish meal and blood meal. This part was
 121 the least in soybean meal and the highest in rapeseed meal ($p < 0.05$). The results of this
 122 study were not consistent with the values reported by other researchers [11] and [12], which
 123 probably are part of the difference between various reports related to the use of different
 124 buffers [13]. The protein with a medium degradation function in the rumen (Fraction B2) is in
 125 fact a nitrogen-free solution in neutral detergent, part of which is broken down into the rumen
 126 and part of the intestine, passing through this part of the rumen to the relative rate of

127 digestion and the passage of dependence has it. The amount of B2 in soybean meal,
 128 cottonseed meal, and rapeseed meal was 80.49, 77.50 and 68.40%, respectively, and in
 129 poultry offal meal, fish meal and blood meal samples were 66.36, 55.03 and 61.66 %,
 130 respectively, in crude protein. In fact, the highest proportion of B2 was related to soybean
 131 meal and the lowest was fish meal ($p < 0.05$). [11] The amount of fast digestible protein in the
 132 rumen was 40 % for cottonseed meal and 72.7 % for soybean meal and 72.7 %, [12] for
 133 cottonseed meal 12.29 and soybean meal 4.09 % of crude protein reported. Because this
 134 fraction is calculated from the discrepancy, so all the measurement errors in this fraction are
 135 gathered, which is probably one of the reasons for the difference between the amounts
 136 reported by various researchers. The heating of feedstuffs destroys B2 proteins and makes
 137 them insoluble, in which case fractions B3 and C increase [14]. Low protein digestibility in
 138 rumen (fraction B3) for soybean meal, cottonseed meal and rapeseed meal was 6.24, 2.63
 139 and 9.11 % respectively, and for poultry offal meal, fish meal and blood meal in
 140 slaughterhouses were 7.50, 6.74 and 11.91 % of crude protein. In the present study, the
 141 highest level of B3 was estimated for blood meal, which was higher than other protein
 142 sources in the table ($p < 0.05$). In [11], the amount of protein with ruminal degradation was
 143 10% for cottonseed meal and 0.8% for soybean meal, and [15], for soybean meal, 1% of
 144 crude protein as Fraction B3 reported. Part B3 Protein is very low in most feedstuffs,
 145 especially plant proteins. These proteins are bound to the cell wall and are insoluble in
 146 neutral detergent. The protein C portion assumed in the rumen's indeterminable CNCPS
 147 system in soybean meal, cottonseed meal and rapeseed meal was 2.45, 9.92 and 4.77%,
 148 respectively, in poultry offal meal, fish meal and blood meal, respectively, 12.21, 17.16 and
 149 3.18 % of crude protein. The highest part of C was related to crude protein of fish meal and
 150 the lowest was related to raw protein content of soybean meal ($p < 0.05$). [11] reported the
 151 amount of crude protein C for cottonseed meal 12.7 % and for soybean meal 5 %, while in
 152 [12] for cottonseed meal 12.29 % and for soybean meal, 11.4 % of crude protein is
 153 presented. Fraction C has a very strong relationship with indigestible nitrogen in rumen of
 154 feed, and therefore the proper and controlled temperature during thermal processes is very
 155 important.

Table 3. protein fractions of some protein sources based on the CNCPS (based on dry matter percent)

Chemical composition	Meal type			Powder type			SEM	p-value
	Plant protein source			Animal protein source				
	Soybean	Rapeseed	Cottonseed	Poultry offal	Fish	Blood		
Crude protein	50.09 ^c	23.84 ^e	36.98 ^d	55.00 ^b	50.09 ^c	59.06 ^a	0.22	<0.001
A	8.52 ^{cd}	6.63 ^d	4.05 ^e	10.38 ^c	13.63 ^b	16.08 ^a	0.76	<0.001
B1	2.30 ^c	3.32 ^c	13.67 ^a	3.54 ^c	7.44 ^b	7.16 ^b	0.91	<0.001
B2	80.49 ^a	77.50 ^a	68.40 ^b	66.36 ^{bc}	55.03 ^d	61.66 ^c	1.83	<0.001
B3	6.24 ^{bc}	2.63 ^c	9.11 ^{ab}	7.50 ^b	6.74 ^b	11.91 ^a	1.32	0.008
C	2.45 ^c	9.92 ^b	4.77 ^c	12.22 ^b	17.16 ^a	3.19 ^c	0.16	<0.001

The different letters in each row represent the difference between the averages. A: Non-Protein Nitrogen, B1: The true protein fast parsing in the rumen, B2: The true protein is parsing, B3: The true protein is decomposed, C: Inaccessible protein

4. CONCLUSION

The results show that there is a difference between the average concentration of insoluble fiber in neutral detergent, crude protein, soluble protein, non-protein nitrogen, insoluble protein in neutral detergent and insoluble protein in acidic detergent of feed samples. Therefore, the use of CNCPS and NRC data for portion fractions of various feeds cannot be considered absolutely, and domestic research and results should be used to extract samples from different regions and different growth conditions, so that diets adjusted with these feeds, Realistic and more balanced and with less waste of nutrients.

REFERENCES

1. Licitra C, Hernandez TN, Van Soest PJ. Standardization of procedures for nitrogen fractionation of ruminant feeds. *Anim Feed Sci Technol.* 1996; 57: 347-358.
2. Van Soest PJ, Sniffen CJ, Oconnor JD, Fox DG, Russel JB. A net carbohydrate and protein system for evaluating cattle diets: 2. Carbohydrate and protein availability. *J Anim Sci.* 1992; 70: 3562-3577.
3. Lanzas CL, Tedeschi O, Seo S, Fox DG. Evaluation of protein fractionation systems used in formulating rations for dairy cattle. *J Dairy Sci.* 2007; 90: 507-521.
4. Sniffen CJ, O'Connor JD, Van Soest PJ, Fox DG, Russell J B. A net carbohydrate and protein system for evaluating cattle diets. II. Carbohydrate and protein availability. *J Anim Sci.* 1992; 70: 3562-3577.
5. Mjoun K, Kalscheur KF, Hippen AR, Schingoethe DJ. Ruminal degradability and intestinal digestibility of protein and amino acids in soybean and corn distillers grains products. *J Dairy Sci.* 2010; 93: 4144-4154.
6. Shahbazi HR, Sadeghi AA, Fazaeli H, Raisali G, Chamani M, Shawrang P. Effects of electron beam irradiation on dry matter degradation of wheat straw in the rumen. *Pakistan J Biological Sci.* 2008a; 11(1): 676-679.
7. NRC. Nutrients requirements of dairy cattle. National academy press. Washington, D. C. 2001.
8. Van Soest PJ. *Nutritional Ecology of the Ruminant* (2nd Ed.). Comstock Publishing Associates, Ithaca, NY; 1994.
9. Riasi A, Allaheresani A, Naimipour H, Fathi MH. Comparison of methods for measuring insoluble fibers in neutral detergents and insoluble fiber in acid detergents in forage and feed products. *J Anim Sci.* 2009; 19(1); 91-103.
10. Khezri A, Yousefi Ansari M, Mohammad Abadi MR. Investigating subunits of protein fractions in comparison with soybean meal and cottonseed using SDS-PAGE and CNCPS electrophoresis method, *J Agri Sci.* 2003; 5 (1):12-18.

- 209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
11. MirzaiiAlamoti HR, Amanloo H, Nikkhah A. Protein and Carbohydrate Fractions of Common Feedstuffs in the Cornell Net Carbohydrate and Protein System. *Iranian J Agri Sci.* 2005; 36 (2): 409-414.
 12. Ghoorchi T, Arbabi S. Study of protein Characteristic of five feeds by CNCPS model. *Asian J Anim and Veterinary Advances.* 2010; 5: 584-591.
 13. Krishnamoorthy U, Muscato TV, Sniffen C J, Van Soest PJ. Nitrogen fractions in selected feedstuffs. *J Dairy Sci.* 1982; 65: 217–255.
 14. Arieli A. Whole cottonseed in dairy cattle feeding: a review. *Anim Feed Sci Technol.* 1998; 72: 97-110.
 15. Shannak S , Suedekum KH , Susenbeth A. Estimating ruminal crude protein degradation with in situ and chemical fractionation procedures. *Anim Feed Sci Technol.* 2000; 85: 195-214.

UNDER PEER REVIEW