

**Original Research Article*****Panax sp.* in Tuyen Quang, North Vietnam – A Potential Plant for  
Poverty Reduction****ABSTRACT**

Ginsengs are perennial forest herbs, belonging to genus *Panax* L. A species of ginseng was found in Tuyen Quang province, North Vietnam and named as *Panax sp.* – Tuyen Quang ginseng. Understanding the ecology, morphology, and saponin of Tuyen Quang ginseng becomes important for development, which can contribute to poverty reduction in the province. Field survey was conducted for ecology and samples were collected for morphology description and saponin analysis. The results indicates Tuyen Quang ginseng distributes in evergreen broadleaved forests in elevation of 980-1,200 m above sea level. It grows in forest with canopy cover of >50% and low vegetation cover of >80%. The soil must be acidic with pH of 4.5-5. Soil contains high ratio of sandy particles and is drainage. Morphology indicates some differences of Tuyen Quang ginseng with other described ginsengs in Vietnam. Saponin content of Tuyen Quang ginseng is 13.7%, which is much higher than that of *Panax stipulealatus* (7.1%) and *P. bipinnatifidus* (7.5%), two widely planted and marketed ginsengs in Vietnam. In addition, current price of Tuyen Quang ginseng is up to 2,600 US\$/1 kg. It is concluded that Tuyen Quang ginseng could be considered as a potential forest herb for poverty reduction. However, study on growing techniques should be conducted before practical application.

**Keywords:** *Ethnic community; Panax sp.; poverty reduction; saponin; Tuyen Quang.*

## 26 1. INTRODUCTION

27 The genus *Panax* L. belonging to family Araliaceae is called ginseng. Until recently, 19  
 28 *Panax* species and subspecies have been described worldwide, most of them grow in eastern  
 29 Asia [1]. Ginsengs have been widely used as traditional medicines [2-3]. The ginsenosides  
 30 (triterpene glycosides) are the main biologically-active compounds of *Panax* L, which was  
 31 first found 50 years ago. Until recently, more than 150 different ginsenosides have been  
 32 isolated from different *Panax* species[4-5].

33 In Vietnam the first ginseng was found in 1973 and named as *Panax vietnamensis* Ha et  
 34 Grushv. [6]. In 2003, a subspecies of *P. vietnamensis* was described and named as *P.*  
 35 *vietnamensis* var *fuscidicus* K. Komatsu, S. Zhu & S.Q. Cai. This subspecies has natural  
 36 distribution in south part of Yunnan province, China and Laichau province, North Vietnam  
 37 [7-8]. Until recently, three ginseng species (*P. vietnamensis* Ha et Grushv, *P.*  
 38 *stipulealatus* Tsai & K.m. Feng, and *P. bipinnatifidus* Seem.) and two subspecies (*P.*  
 39 *vietnamensis* var *fuscidicus* and *P. vietnamensis* var. *langbianensis*) have been found in  
 40 Vietnam.

41 Tuyen Quang province locates in North Vietnam with diversity of vegetation types,  
 42 topography, and climate conditions. A ginseng (*Panax* sp.) was found to have natural  
 43 distribution in narrow areas of Lam Binh and Na Hang districts of Tuyen Quang province, so  
 44 called “Tuyen Quang ginseng”. It has been harvested from natural for health improvement  
 45 and marketing to generate income for local ethnic communities for decades. The objectives of  
 46 this study were (1) to describe ecological and morphological characteristics and (2) to  
 47 analyze roots for saponin content of Tuyen Quang ginseng, which could be used to develop  
 48 and manage this economically valuable ginseng sustainably.

49

## 50 2. MATERIALS AND METHOD

### 51 2.1 Materials

52 By interviewing local authorities and ethnic people, it was reported that Tuyen Quang  
 53 ginseng has natural distribution in Sinh Long commune, Na Hang district and Thuong Lam  
 54 commune, Lam Binh district of Tuyen Quang province (Fig. 1). Therefore, those two  
 55 communes were selected for field survey.

56

57 **2.2 Methods**

## 58 2.2.1 Natural and Ecological Characteristics

59 By interviewing local people, who have found and harvested Tuyen Quang ginseng, the areas  
 60 of natural distribution became known. With their field guidance, it was easy for the survey  
 61 team to get the desired survey locations. In the field, if plant of Tuyen Quang ginseng was  
 62 found, natural and ecological characteristics were described including forest type, height of  
 63 forest canopy, vertical structure of forest canopy, forest canopy cover, and low vegetation  
 64 layer. In addition, a soil pit of 1.2 m depth and 60 cm width was dug up for soil profile, and a  
 65 soil sample was collected from 0-20 cm soil depth for analyzing in Lab.

66

## 67 2.2.2 Morphological and Anatomy Characteristics

68 Five fully-developed plants of Tuyen Quang ginseng were randomly selected for  
 69 morphological characteristics including leaves, stem, root, flower, fruit, and seed. Shape,  
 70 color, and dimension of each organ were described by naked-eyes. In addition, slices of stems  
 71 and roots were also used for their structure by observing through microscope [9]. Slices of  
 72 40-60  $\mu\text{m}$  was cut by using special equipment. The slice was then colored by chrysanthemum  
 73 (0.5%) and safranin (1%). It was then cleaned by purified water and observed through  
 74 microscope.

75

## 76 2.2.3 Saponin Content

77 Total saponin was estimated by weighing. Approximate one gram powder of the dried root  
 78 sample were weighed, 100 ml *n*-hexane were then added. The mixture was extracted in  
 79 Soxhlet extraction apparatus in six hours and filtered. The residue was continuously extracted  
 80 by Soxhlet extraction apparatus with 100 ml of 70% methanol for 6 hours. The combined  
 81 filtrate was concentrated under reduced pressure with a rotary evaporator to obtain extract.  
 82 The methanol extract was diluted in 30 ml of water and then fractionated with water saturated  
 83 *n*-butanol until no color was observed in *n*-butanol layer. The filtrated butanol extract was  
 84 then evaporated under reduced pressure to yield butanol extract. This extract was dissolved  
 85 into 10 ml of 70% ethanol then transferred into a porcelain beaker and evaporated solvent to

get extract. The obtained extract was dried in an oven at 105 °C until constant weight. Total saponin content (X) was calculated as:  $X = [(b \times 100)/(m \times (100 - d))] \times 100$ , where b is obtained saponin weight (g), d is moisture of root powder (%), and m is initial weight of root powder (g).

### 3. RESULTS

#### 3.1 Ecology

Tuyen Quang ginseng naturally distributes in evergreen broadleaved forests of both disturbed and undisturbed forests. The height of forest canopy is taller than 15 m. The canopy structure includes two or three layers with the shortest layer of 7-10 m tall and the difference between layers of 4-6 m. Tuyen Quang ginseng prefers to grow in forests with canopy cover of >50%. The species composition of forest canopy, where Tuyen Quang ginseng grows, includes tree species of *Saurauia napaulensis* DC., *Callicarpa arborea* Roxb., *Alniphyllum eberhardtii* Guillaumin, *Schefflera macrophylla* (Dunn) R.Vig., *Machilus chinensis* (Benth.) Hemsl., and *Pavetta indica* L. with density of 650-860 trees/ha. The vegetation layer on forest floor includes plants of *Asarum caudigerum* Hance, *Lophatherum gracile* Brongn., *Alpinia chinensis* (Retz.) Roscoe, *Cyclosorus parasiticus* (L.) Farw., and *Polygonum chiensis* L. with cover >80%. The litter layer on forest floor is thick and wet most time of the year.

Suitable climate conditions for Tuyen Quang ginseng include annual precipitation of 1,700-1,900 mm, annual air humidity of >85%, annual air temperature of 23°C, minimum temperature of >5°C, and maximum temperature of <30°C. Tuyen Quang ginseng distributes on elevation zone of 980-1,200 m above sea level, growing well in mountain foots, flat areas, and well drainage soils.

Soil profile indicates the litter layer of 4-5 cm, humus layer of 2-3 cm with dark color and humidity of 56-60%, and total soil depth of >100 cm. Tuyen Quang ginseng naturally grows in acidic soil with pH of 4.5-5.0, high humus soil of 6.5-9% (Table 1). Content of sand particles is high, indicating drainage soil. Nitrogen is not much required by Tuyen Quang ginseng as it can grow in low N content soil of 0.5%.

#### 3.1 Morphology

Tuyen Quang ginseng is a perennial plant, up to 60 cm tall at maturity (Fig. 2). Root lies horizontally without root branching. Each root contains one aerial stem with leaves. In very rare case, it may contains 2-3 aerial stems.

Root is brown to light yellow (Fig. 2) with numerous scars as results of dead aerial stems of each growing year. Scars arranges alternately in root. The head of root is big and global, while the end of root is much smaller and contains bunch of small roots for absorbing water and nutrient. Inner part of roots is light yellow or violet (Fig. 2). Total root length is 3.2-26.5 cm and diameter is 1.2-3.3 cm.

Aerial stem dies in winter and new stem starts from root head in spring. This creates scars in roots (Fig. 2). Aerial stem is 20-60 cm height and 0.3-0.8 cm diameter. It is green and/or light violet without hairs (Fig. 3). Each aerial stem contains 1-4 main leaves and each main leaf contains 4-6 sub-leaves (Fig. 3). There are tiny hairs covering both sides of sub-leaves (Fig. 3).

Each aerial stem contains an inflorescence on top (Fig. 4), which is 15-25 cm length and contains 40-120 flowers. The inflorescence is 3-4.5 cm in diameter. When ripen, fruits become red. Seeds are white with a triangle shape. Flower season is May-July and season of ripen fruits is October-December.

### 3.3 Anatomy and Saponin

Aerial stem and root of a 6-year old plant of Tuyen Quang ginseng were collected for anatomy and saponin analysis. Results indicates aerial stem of Tuyen Quang Ginseng includes main parts as cutin layer in outermost, epidermis, xylem, phloem, and xylem ray (Fig. 5). While, transverse section of root indicates main layer of epidermis, oxalate, phloem, xylem, and xylem ray (Fig. 6).

Total saponin content of Tuyen Quang ginseng (*Panax* sp.) is 13.7%, which is much higher than that of *P. stipuleanatus* and *P. bipinnatifidus* (Table 2). However, it is lower than *P. vietnamensis* var. *fuscidiscus* (21.9%) and *P. vietnamensis* (22.3%), two well-known ginsengs in Vietnam.

## 4. DISCUSSION

Forest herbs like ginseng can only survive and grow well under shades of other vegetation [10-12]. Therefore, Tuyen Quang ginseng can only be found in evergreen broadleaved with forest canopy cover of >50%, and lower vegetation cover of >80% in the present study. Forest structure including more than two layers of trees, and lower vegetation [13] is an important indicator for the existence of Tuyen Quang ginseng. Low cover of tree layer will allow direct sunlight to forest floor, reducing soil moisture, litter layers, soil organic carbon *etc.*, which is not a favor condition for existence of Tuyen Quang ginseng [14].

Soil with high ratio of sandy particles and high humus content (Table 1) indicates low compact soil, which supports growth of roots in the soil. If soil is compact with low humus content, the soil is high water holding capacity with low soil air. Such kind of soil will never well support growth of plants which have roots as main part of their body. Therefore, selecting suitable soil is important for growing Tuyen Quang ginseng. It must be high humus content and drainage soil.

Comparing morphology among five ginsengs in Vietnam indicates several differences of Tuyen Quang ginseng with others (Table 3). The most different characteristic is number of leaves per aerial stem, which contains 1-4 leaves per stem in Tuyen Quang ginseng compared to 3-5 in others. While, no. flowers were also most numerous of up to 120 flower per inflorescence. It seems that Tuyen Quang ginseng most look like *P. vietnamensis*. However, there is a possibility of a new sub-species found in Tuyen Quang province. Therefore, further study such as DNA analysis is required to identify where the Tuyen Quang ginseng (*Panax* sp.) is a new sub-species of ginseng in Vietnam.

Anatomy of Tuyen Quang ginseng is similar to that of other ginseng and plants [9, 15], which have been widely used as traditional medicines. Root shows clusters of oxalate, which is known as containing high saponin; the main active chemical component of ginseng.

The saponin content of Tuyen Quang ginseng is much higher than that of *P. stipulealatus* and *P. bipinnatifidus* (Table 2), which have been widely grown and marketed for poverty reduction in mountainous areas of Vietnam. In addition, current price of Tuyen Quang ginseng ranges 1,000-2,600 US\$/ 1 kg, depending on size and age of roots [16]. Therefore, growing Tuyen Quang ginseng could also contribute to poverty reduction in Tuyen Quang province. However, before practical application researches on planting Tuyen Quang ginseng should be conducted extensively to issue the applicable growing guideline, ensuring the quality/saponin content of planted ginseng compared to natural ones.

## 5. CONCLUSION AND RECOMMENDATION

Tuyen Quang ginseng –*Panax sp.* has natural distribution in Thuong Lam commune Lam Binh district and Sinh Long commune, Na Hang district of Tuyen Quang province, North Vietnam. The species distributes in evergreen broadleaved forests on the elevation of 980–1,200 m above sea level. The species only distributes in forests with canopy cover >50% and cover of low vegetation layer of >80%. The soil must be acidic with pH of 4.5-5.0 and high ratio of sandy particles, indicating drainage soil.

There is a possibility that Tuyen Quang ginseng is a new sub-species of ginseng because of some differences of its morphological characteristics compared to other ginsengs in Vietnam. However, detail analysis such as DNA application should be conducted for better conclusion.

Tuyen Quang ginseng has high saponin content (13.7%) and is a potential forest herb for poverty reduction to ethnic communities in the province. Study on growing techniques must be conducted, which is valuable for local communities. Selecting vegetation types for growing Tuyen Quang ginseng is important, which must have high canopy cover of > 50% and high cover of low vegetation layer of >80%.

## 196 REFERENCES

- 197 1. Pandey AK, Ali MA. Intraspecific variation in *Panax assamicus* Ban. Populations based  
198 on internal transcribed spacer (ITS) sequences of nrDNA. International Journal  
199 Biotechnology. 2012;11:30–38.
- 200 2. Shim SC, Chang SK, Hur CW, Kim CK. New polyacetylene compounds from *Panax*  
201 *ginseng* C. A. Meyer. Bulletin of Korean Chemical Society. 1987; 8:272–275.
- 202 3. Kochkin DM, Kachala VV, Shashkov AS, Chizhov AO, Chirva VY, Nosov AM. Malonyl-  
203 ginsenoside content of a cell-suspension culture of *Panax japonicus* var. repens.  
204 Phytochemistry. 2013; 93:18–26.
- 205 4. Qi LW, Wang CZ, Yuan CS. Ginsenosides from American ginseng: chemical and  
206 pharmacological diversity. Phytochemistry. 2011; 72:689–699.
- 207 5. Yoshizaki K, Murakami M, Fujino H, Yoshida N, Yahara S. New triterpenoid saponins  
208 from fruit specimens of *Panax japonicus* collected in Toyama prefecture and Hokkaido  
209 (2).Chemical and Pharmaceutical Bulletin. 2012; 60:728–735.
- 210 6. Dung HT, Grushvisky IV. A new species of the genus *Panax* L., Araliaceae in Vietnam:  
211 *Panax vietnamensis* Haet Grushv. Botany Journal Vietnam.1985; 70:518–522.
- 212 7. Zhu S, Fushimi H, Qing CS, Biao CH, Komatsu K. A new variety of the genus *Panax* from  
213 Southern Yunnan, China and its nucleotide sequences of 18S ribosomal RNA gene and  
214 matK gene. Journal of Japanese Botany. 2003; 78: 86–94.
- 215 8. Phan KL, Le TS, Phan KL, Vo DD, Phan VT. Lai Chau ginseng *Panax vietnamensis* var.  
216 *fuscidiscus* K. Komatsu, S. Zhu & S.Q. Cai I. morphology, ecology, distribution and  
217 conservation status. Proceeding of the 2nd VAST-KAST Workshop on Biodiversity and  
218 Bio-active compounds.2013; pp. 65–73.
- 219 9. Liu EH, Qi LW, Li K, Chu C, Li P. Recent advances in quality control of traditional  
220 Chinese medicines. Combinatorial Chemistry & High Throughput Screening. 2010;  
221 13:869–884.
- 222 10. Anderson RC, Fralish JS, Armstrong JE, Benjamin PK. The ecology and biology of  
223 *Panax quinquefolium* L. (Araliaceae) in Illinois. American Midland Naturalist.1993;  
224 129:357–372.
- 225 11. McKenzie D, Halpen CB, Nelson CR. Overstory influences on herb and shrub  
226 communities in mature forests of western Washington, U.S.A. Canadian Journal ofForest  
227 Research. 2000; 30:1655–1666.
- 228 12. Pabst RJ, Spies T. Distribution of herbs and shrubs in relation to landform and canopy  
229 cover in riparian forests of coastal Oregon. Canadian Journal of Botany. 1997; 76:298–  
230 315.
- 231 13. Turner IM, Tan HTW, Chua KS. Relationships between herb layer and canopy  
232 composition in a tropical rain forest successional mosaic in Singapore. Journal  
233 ofTropical Ecology1996; 12:843–851.
- 234 14. Beyfuss R. Growing ginseng fact sheet. USDA, Natural Resources Conservation Service  
235 401. 2000. <http://hwwff.cce.cornell.edu/docs/GinFs.pdf>(accessed: 1 May, 2018)
- 236 15. Tengku AS, Tengku M, Humera N, Ratni SJ, Khatijah H, Mohd RAR, Aishad A, Jean-  
237 FredericFW. Chemical and pharmacognostical characterization of two Malaysian plants  
238 both known as Ajisamat. Revista Brasileira de Farmacognosia. 2013; 23:724–730.
- 239 16. Pham QT. Growing and developing Laichau ginseng. Scientific Report. Silviculture  
240 Research Institute, Vietnamese Academy of Forest Sciences. Hanoi. 2018.



**Table 1.** Physical and chemical characteristics ( $\pm$ SE) of soil samples

Research site	Density (g/cm <sup>3</sup> )	Silt (< 0.002 mm; %)	Lam (0.002-0.02 mm; %)	Sand (2-0.02 mm; %)	pH	Humus content (%)	N (%)	K <sub>2</sub> O (%)
Thuong Lam, Lam Binh	1,135.8 $\pm$ 27.0	12.7 $\pm$ 0.9	33.0 $\pm$ 1.9	54.3 $\pm$ 2.7	4.5 $\pm$ 0.3	6.7 $\pm$ 0.9	0.5 $\pm$ 0.1	2.5 $\pm$ 0.3
Sinh Long, Na Hang	1,156.5 $\pm$ 8.5	4.3 $\pm$ 0.0	34.9 $\pm$ 1.4	60.8 $\pm$ 1.4	4.9 $\pm$ 0.3	8.9 $\pm$ 1.1	0.6 $\pm$ 0.0	3.2 $\pm$ 0.3

UNDER PEER REVIEW

244 **Table 2.** Saponin content of ginsengs in Vietnam

Species	Root source	Saponin content (%)
<i>Panax sp.</i> - Tuyen Quang ginseng (study species)	Natural, Tuyen Quang province	13.7
<i>Panax stipulealatus</i> Tsai & K.m. Feng	Growing in garden of local people in Laichau Province, five years old	7.1*
<i>Panax bipinnatifidus</i> Seem.,	Growing in garden of local people in Laichau Province, five years old	7.5*
<i>Panax vietnamensis</i> var <i>fuscidicus</i> K. Komatsu, S. Zhu & S.Q. Cai.	Growing in garden of local people in Laichau Province, five years old	21.9*
<i>Panax vietnamensis</i> Ha et Grushv	Natural, central Vietnam	22.3*


245 \* cited from Pham et al. [16].

246 **Table 3.** Morphological characteristics of five ginsengs in Vietnam.

	Species				
	<i>P. stipuleanatus</i>	<i>P. vietnamensis</i>	<i>P. vietnamensis</i> var. <i>fuscidiscus</i>	<i>P. notoginseng</i>	<i>Panax sp.</i> ( <i>Tuyen Quang ginseng</i> )
Root distribution	Horizontal	Horizontal	Horizontal	Vertical	Horizontal
Root scar	In line	Alternate	Alternate	Random	Alternate
Color of inner part of root	White green	Yellow or light violet	Yellow or light violet	Yellow or light violet	Yellow or light violet
Aerial stem	Violet or green	Violet	Violet	Violet	Green or light violet
Leaves per stem	3	3-5	3-5	4	1-4
No. flowers	60-80	50-120	40-120	40-120	40-120
Flower color	White green	White green	White green	White green	White green
Stamen	1	1-2	1-2	2-3	1-2
Pistil	1-2	1-2	1-3	2-3	1-2
Color of ripen fruit	Red	Red with black dots on top	Red with black dots on top	Red	Red
Shape of fruit	Kidney shape	Kidney shape	Kidney shape	Half global or triangle	Kidney shape or half global
Seeds per fruit	1-2	1-2	1-3	2-3	1-2
Shape of seed	Kidney shape	Kidney shape	Kidney shape	Triangle	Triangle

247

248



**Fig. 1.**(a) Map of Vietnam, (b) Tuyen Quang province, (c) Lam Binh and Na Hang districts – field survey sites.

254



255



256


257 **Fig. 2.** Full plant of Tuyen Quang ginseng (above left), root (above right), and root transverse  
258 sections (below; light yellow in left and violet in right).




**Fig. 3.** Aerial stem (1), sub-leaves (2) back side of leaf (3), front side of leaf (4), petal (5), and bract (5).



**Fig. 4.** Inflorescence (1), bract (2), palea (3), sepal back side of leaf (3), front side of leaf (4), petal (5), stamen (6), pistil (7), stigma splits (8), fruit lengthwise (9), fruit (10).



**Fig. 5.** Transverse section of aerial stem. Cutin (1), epidermis (2), collenchyma (3), axial parenchyma (4), sclerenchyma (5), xylem (6), phloem (7), xylem ray (8).



**Fig. 6.** Transverse section of root. Epidermis (1), axial parenchyma (2), clusters of calcium oxalate (3), phloem (4), xylem (5), xylem rays (6).