

2

3 **Effect of drying on the composition of secondary**

4 **metabolites in extracts from floral parts of *Curcuma***

5 ***longa* L.**

6

7 **ABSTRACT**

8 **Background:** *Curcuma longa* is a rhizomatous herbaceous plant of Zingiberaceae family, originating  
9 from South Asia and very widespread in hot and rainy regions of the globe. The rhizomes are very  
10 popular spice and used as food additives for its coloring, aromatic, food preservation and nutritional  
11 properties.

12 **Aim:** The aim of this study was to assess the drying effect of floral parts of *C. longa* at laboratory and  
13 incubator temperature (35°C) on the composition of secondary metabolites in general and  
14 polyphenolic compounds in particular.

15 **Place and Duration of Study:** The study was carried out at the Laboratory of Natural Products,  
16 Department of Chemistry, Faculty of Sciences, University of Kinshasa between November 20 and  
17 December 10, 2016.

18 **Methodology:** Different parts of *C. longa* floral parts were collected. Petals were collected every day  
19 while sepals were collected 20 days after the first petal appeared. The phytochemical screening was  
20 used as per the standards protocol and it was assessed between floral parts dried in the room  
21 temperature and floral parts dried in the incubator at 35 °C.

22 **Results:** The findings revealed the presence of total polyphenols, flavonoids, tannins in petal extracts  
23 dried in an incubator at 35°C. However, flavonoids and tannins were not detected in extracts from  
24 petals dried at laboratory temperature. Phytochemical screening findings of the organic phase of *C.*  
25 *longa* floral parts revealed the presence of free quinones and terpenes in sepal extracts dried in the  
26 incubator at 35°C while those of extracts from the same part of the dried plant in the laboratory  
27 temperature revealed the presence of terpenes and steroids.

28 **Conclusion:** Future studies should carry out a similar study using the spectrophotometry method to  
29 determine polyphenolic compounds and confirm our hypothesis on the degradation of polyphenolic  
30 compounds during the drying of *C. longa* floral parts at laboratory temperature.

31

32 **Keywords:** *Curcuma longa*, Phytochemical screening, Effect of drying, Laboratory, Incubator

33

34 **1. INTRODUCTION**

35 *Curcuma longa* or saffron is a rhizomatous herbaceous plant, perennial, 50 cm to 1m high, of the  
36 Zingiberaceae family, originating from South Asia and very widespread in hot and rainy regions of the  
37 globe such as Asia, Africa and Oceania [1]. It is widely cultivated in India but also to a lesser extent in  
38 China, Taiwan, Japan, Burma, Indonesia and Africa. The intense cultivation of this plant is due to its  
39 rhizomes [1-2]. The latter dried and powdered are a very popular spice and are used as food additives  
40 for its coloring, aromatic, food preservation and nutritional properties. In addition, *C. longa* rhizomes  
41 are a key ingredient in traditional South American and Asian medicines such as Ayurvedic medicine  
42 (Indian medicine). Thus, *C. longa* is a remedy against gastrointestinal, digestive disorders,  
43 inflammatory diseases, skin diseases [1-5].

44

45 Numerous studies correlated with ancestral knowledge have demonstrated the preventive action of *C.*  
46 *longa* rhizomes on many diseases such as cancers, cardiovascular diseases [6]. Several studies have  
47 shown that extracts from *C. longa* rhizomes possess a large pharmacological potential namely: anti-  
48 cancer, anti-inflammatory, healing, cholesterol-lowering, hypoglycemic, anti-Alzheimer's,  
49 antiplasmodial, anti-inflammatory, antioxidant, antibacterial, antifungal, anti-venomous, antipyretic,

50 analgesic, inhibits the action of HIV-1 integrase, and HIV-1 integrase protein replication, protects  
 51 against diabetic retinopathy and many other pathologies [2-5, 7-12]. In addition to rhizomes, the other  
 52 organs of the plant (leaves, roots and floral parts) are less used. The leaves of *C. longa* are  
 53 sometimes used in cooking or to extract essential oils [5]. Ritwiz *et al.* [13] have shown that extracts  
 54 from the leaves of *C. longa* have antioxidant, antibacterial potential and can modulate immunological  
 55 properties. Meanwhile Mbadiko *et al.* [11-12], report that total methanol extracts of rhizomes, roots,  
 56 leaves and floral parts (petals and sepals) possess an antisickling activity. Most of the studies carried  
 57 out so far focused only on the rhizomes of the plant, studies on the leaves, roots and floral parts of *C.*  
 58 *longa* remain poor or less reported. This justifies our interest in carrying out a phytochemical study on  
 59 the extracts of the floral parts (petals and sepals) of *C. longa*.

60  
 61 In addition, Mbadiko *et al.* [11-12] reported that floral parts of *C. longa* contained low levels of  
 62 secondary metabolites (polyphenolic compounds) and thought this was related to the drying effect.  
 63 These authors reported that during the drying of *C. longa* floral parts, and in particular the petals at  
 64 laboratory temperature, the latter tended to soften and lose color, suggesting degradation of the  
 65 phytoconstituents under cellular conditions. The negative impact of post-harvest treatments of plant  
 66 samples on phytoconstituent composition has also been reported by Singh *et al.* [14]. The degradation  
 67 of polyphenolic compounds in plants under cellular conditions, i.e. fresh samples, has also been  
 68 reported by Yan *et al.*, [15] and Chang *et al.* [16]. Thus, this study is part of a context to assess the  
 69 effect of post-harvest treatments, in particular drying on the composition of secondary metabolites in  
 70 extracts from *C. longa* floral parts.

71

72 **2. MATERIALS AND METHODS**

73 **2.1 Collection of plant material**

74 As biological material, the floral parts of *C. longa* were used. It was observed that *C. longa* renews its  
 75 petals every 24 hours. Thus, petals were collected every day (between November 20 and December  
 76 10, 2016). The sepals were collected 20 days after the first petal appeared.

77

78 **2.2 Packaging of plant material**

79 After the collection, the petals and sepals were cleaned, washed quickly with tap water. Some of our  
 80 samples were dried at laboratory temperature and the other part at the incubator (Melag Nurfur  
 81 Wechselstrom brand) at 35°C.

82

83 **2.3 Preparation of aqueous and organic extracts**

84 The aqueous extracts were obtained by macerating 5g of the powder of our four samples each in 50  
 85 mL of distilled water during 24 hours at room temperature, then filtered using filter paper (Whatman  
 86 n°1). Maceration of 2g of the powder from our samples in 20 mL of ethyl acetate during 24 hours of  
 87 incubation then filtered using a filter paper (Whatman n°1).

88

89 **2.4 Phytochemical screening**

90 Phytochemical screening represents all the qualitative techniques used to determine or identify the  
 91 different chemical groups (secondary metabolites) contained in an extract. These chemical groups are  
 92 identified by means of coloring and precipitation reactions that take place by adding specific reagents  
 93 [11, 17].

94

95 **3. RESULTS AND DISCUSSION**

96 The results of the phytochemical screening of aqueous and organic extracts of *C. longa* floral parts  
 97 dried in the incubator at 35 °C are presented in the following tables.

98

99

100 **Table 1 : Phytochemical screening of aqueous phase of floral parts dried at the incubator at 35**  
 101 **°C**

Secondary metabolites	Aqueous phase	
	Petals	Sepals
Saponines	-	-
Total polyphenols	+	+
Flavonoids	+	+
Anthocyanins	-	+

Leuco-anthocyanins	-	-
Alkaloids	-	+
Bound Quinones	-	-
Tannins	+	+

102 Legend : + :presence of the phytoconstituent, - :absence of the phytoconstituent

103

104 Table 1 shows that the aqueous phases of different floral parts (petals and sepals) of *C. longa* dried in  
 105 the incubator at 35 °C differ in their composition in secondary metabolites. Total polyphenols,  
 106 flavonoids, and tannins were detected in petal extracts. On the other hand, the analysis on sepal  
 107 extracts revealed the presence of total polyphenols, flavonoids, anthocyanins, alkaloids and tannins. It  
 108 should be noted that saponins were not detected in all extracts.

109

110 **Table 2. Phytochemical screening of organic phases of floral parts of *C. longa* dried at the**  
 111 **incubator at 35 °C**

112

Secondary Metabolites	Organic phase	
	Petals	Sepals
Terpenes	+	+
Steroids	-	-
Free Quinones	+	+

113 Legend : + :presence of the phytoconstituent, - :absence of the phytoconstituent

114

115 Table 2 shows that the organic phase of extracts from different floral parts of *C. longa* dried in the  
 116 incubator at 35 °C has the same chemical profile. The presence of free terpenes and quinones has  
 117 been detected in both parts of the plant.

118

119 **Table 3 : Phytochemical screening of aqueous phase of *C. longa* floral parts dried at the**  
 120 **laboratory temperature**

121

Secondary metabolites	Aqueous phase	
	Petals	Sepals
Saponines	-	-
Total polyphenols	+	+
Flavonoids	-	-
Anthocyanins	-	+
Leuco-anthocyanins	-	-
Alkaloids	+	+
Bound Quinones	-	-
Tannins	-	+

122 Legend : + :presence of the phytoconstituent, - :absence of the phytoconstituent

123

124 Table 3 shows that extracts of *C. longa* petals possess total polyphenols and alkaloids while the  
 125 presence of total polyphenols, anthocyanins, tannins and alkaloids have been highlighted in sepal  
 126 extracts.

127

128

129

130

131

132

133 **Table 4 : Phytochemical screening of organic phase of *C. longa* floral parts dried at the**  
 134 **laboratory temperature**

135

Secondary Metabolites	Organic phase	
	Petals	Sepals
Terpenes	-	+
Steroids	-	+
Free Quinones	-	-

136 Legend : + :presence of the phytoconstituent, - :absence of the phytoconstituent.

137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196

Table 4 shows the presence of terpenes and steroids in petal extracts. None of these compounds were detected in the petal extracts.

## DISCUSSION

### - Aqueous phase

Phytochemical analyses in this study revealed the presence of total polyphenols, flavonoids, tannins in petal extracts dried in an incubator at 35°C. However, flavonoids and tannins were not detected in extracts from petals dried at laboratory temperature. This implies their degradation under laboratory conditions. In fact, since biochemical reactions necessarily occur in an aqueous environment, drying at a temperature that does not remove moisture at a short time would favor the action of certain enzymes that are activated during drying [19], could be at the origin of the degradation of certain secondary metabolites. As mentioned above, the degradation of polyphenolic compounds in fresh plant samples has also been reported by Yan *et al.*, [15] and Chang *et al.* [16].

In addition, phytochemical screening of sepal extracts dried in the incubator at 35°C revealed the presence of total polyphenols, flavonoids, anthocyanins, tannins and alkaloids while those of sepals dried at laboratory temperature showed the presence of total polyphenols, anthocyanins, tannins and alkaloids. The absence of flavonoids in sepals and petals dried at laboratory temperature would suggest their degradation during drying; this would at the same time justify the discoloration of these samples during drying. Indeed, flavonoids are pigments responsible for the yellow, orange and red discoloration of different plant organs [18]. Further studies are needed to confirm this hypothesis.

### - Organic phase

Phytochemical screening findings of the organic phase of *C. longa* floral parts revealed the presence of free quinones and terpenes in sepal extracts dried in the incubator at 35°C while those of extracts from the same part of the dried plant in the laboratory temperature revealed the presence of terpenes and steroids. Phytochemical screening of petal extracts dried in the incubator at 35°C showed the presence of free quinones and terpenes. These compounds were not detected in petal extracts dried at laboratory temperature. The absence of quinones in sepal and petal extracts dried at laboratory temperature could justify the discoloration of these organs during drying at room temperature and would affirm their degradation during drying. Boukri [18] reported that quinones are colored and bright substances, usually red, yellow or orange.

## CONCLUSION AND RECOMMENDATIONS

The purpose of this study was to assess the drying effect of floral parts (petals and sepals) at laboratory or incubator temperature at 35°C on the composition of secondary metabolites in general and polyphenolic compounds in particular. In light of the findings obtained, we believe that drying the floral parts (petals and sepals) of *C. longa* at laboratory temperature would promote the degradation of certain polyphenolic compounds (flavonoids and quinones). This would prevent the action of certain enzymes that would be activated after harvest or during drying and would prevent the degradation of polyphenolic compounds. Samples should also be cut into small pieces before drying to increase the surface area of contact of the samples with heat and rapidly reduce moisture. The study on the drying effect on the composition of secondary metabolites has not yet been reported in the literature.

Future studies should carry out a similar study using the spectrophotometry method to determine polyphenolic compounds and confirm our hypothesis on the degradation of polyphenolic compounds during the drying of *C. longa* floral parts at laboratory temperature.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

- 198 [1]. Savina P. Le Curcuma, un agent naturel contre le vieillissement. Master en Biologie-gestion,  
199 Université de Rennes1, UFR Sciences de la vie et de l'environnement, 2014.
- 200 [2]. Cheikh ZA. Etudes chimiques et biologiques d'*Aframomum sceptrum* (Zingibéraceae) et de la  
201 curcumine. Université paris-sud, Faculté de pharmacie de chatnay-Melabry, 2013, 161pp.  
202 ([http:// www.tel.archives-ouvertes.fr](http://www.tel.archives-ouvertes.fr))
- 203 [3]. Hombourger C. Le Curcuma de l'épice au médicament. Thèse de Doctorat, Université Henri  
204 Poincare-Nancy 1, Faculté de pharmacie, 2010, 222p. ([http://www.culture.gouv.fr/culture/infos-  
205 pratique/droits/protection.htm](http://www.culture.gouv.fr/culture/infos-pratique/droits/protection.htm))
- 206 [4]. Perry MC. Evaluation de la Curcumine comme agent anticancéreux dans le traitement des  
207 tumeurs cérébrales. Mémoire de Master en chimie, Université du Québec à Montréal, 2008,  
208 97pp.
- 209 [5]. Vaquier ARL. Intérêt d'un nouveau nutriment a visée anti-inflammatoire dans la gestion de  
210 troubles locomoteurs de cheval. Aspects bibliographiques et études cliniques. Ecole vétérinaire  
211 d'Alfort, Faculté de Médecine de Cretal, 2010, 199 pp.
- 212 [6]. Aggarwal BB, Kumar A, Bharti AC. Anticancer potential of curcumin : Preclinical and clinical  
213 studies. *Anticancer Research*, 2003, 23, 363-398.
- 214 [7]. Araujo CC, Leon LL. Biological activities of *Curcuma longa* L. *Mem Inst Oswaldo Cruz*, 2001;  
215 96(5): 723-728.
- 216 [8]. Anbujeba J, Suraj R., Renitha G., Anandrarajagopal K., Gnamakumari Anita A.V., Promwichit P.  
217 *In vitro* antimicrobial evaluation of *Zingibera ffcinale*, *Curcuma longa* and *Alpinia galangal*  
218 Extracts as Natural food preservatives. *America Journal of Food Technology*, 2002; 4 (5):192-  
219 200. ([http: www.Scialert.net](http://www.Scialert.net)).
- 220 [9]. Goel A, Kunnumkkara B, Aggarwal BB. Curcumin as «Curecumin»: From kitchen to clinic.  
221 *Biochem. Pharmacol.*, 2008; 75: 787 – 809.
- 222 [10]. Sivabalan S, Anuradha CV. A Comparative study on the antioxidant and glucose lowering  
223 effects of Curcumin Analogue trough in vitro essay. *International Journal of Pharmacology*,  
224 2010, 6 (5): 664-669 pp. ([http: www.Scialert.net](http://www.Scialert.net)).
- 225 [11]. Mbadiko CM, Ngbolua KN, Mpiana PT, Mutambel ' H, Kikakedimau RN, Makengo KG, Pambu  
226 AL, Kemfine LL., Bongo GN, Mbemba TF. Phytochemical screening and assessment of anti-  
227 sickling activity of total methanolic extracts of different organs of *Curcuma longa* L.  
228 (Zingiberaceae). *The Pharmaceutical and Chemical Journal*, 2017, 4(1):32-40.
- 229 [12]. Mbadiko CM, Ngbolua KN, Mpiana PT, Tshilanda DD, Makengo KG, Pambu LA, Kemfine LL,  
230 Bamvingana C, Bongo. GN and Mbemba FT. Assessment of the antisickling activity of total  
231 methanolic extracts from the rhizomes and roots of *C. longa* and the effect of photodegradation  
232 on the antisickling activity. *Journal of Advancement in Medical and Life Sciences* 2017, 5(1).
- 233 [13]. Ritwiz BS, Kusha BS, Ranjeet D, Ganesh N. *Curcuma longa* leaves exhibits a potential  
234 antioxidant, antibacterial and immunomodulating properties. *International Journal of*  
235 *Phytomedicine*, 2001; 3 (11), 270-278 <http://www.arjournals.org/index.php/ijpm/index>.
- 236 [14]. Singh G, Kapoor IPS, Pratibha S, Carola HS, Marina de Lampasona P, Catalan CAN.  
237 Comparative study of chemical composition and antioxidant activity of fresh and dry rhizomes of  
238 turmeric (*Curcuma longa* Linn.). *Food and Chemical Toxicology*, 2010, 48 (2010) 1026–1031.
- 239 [15]. Yan SW and Asmah R. Comparison of total phenolic contents and antioxidant activities of  
240 turmeric leaf, pandan leaf and torch ginger flower. *International Food Research Journal*, 2010 ;  
241 17: 417-423.
- 242 [16]. Chang CH, Lin HY, Chang CY and Liu YC. Comparisons on the antioxidant properties of fresh,  
243 freeze-dried and hot-air-dried tomatoes. *Journal of Food Engineering*, 2006, 77: 478-485.
- 244 [17]. Harbone JB. Methods of extraction and Isolation 'Phytochemical Methods'. 3rd edition  
245 Chapman and Hall, London, 1998, 60 - 66.
- 246 [18]. Boukri NEH. Contribution à l'étude phytochimiques des extraits bruts des épices contenus dans  
247 le mélange Ras-el-hanout. Mémoire de Master, Université KasdiMerbah Ouargla, Faculté des  
248 Sciences de la Nature et de la Vie, 2014, 99 pp.
- 249 [19]. Tula YM. Activité anti drépanocytaire et stabilité des anthocyanes extraits des bulbes d'*Hypoxis*  
250 *angustifolia* et de feuilles d'*Ipomoea batatas*, Mémoire de licence, Faculté des Sciences,  
251 Département de chimie, Université de Kinshasa, 2011.
- 252