Original Research Article

Effect of silicon and phosphorus additions and their interactions on wheat plants grown on a clay soil

Abstract

A pot experiment was conducted in clay soil collected from Agricultural Research Center farm, Giza governorate, Egypt. Wheat grains (Triticumaestivum L., Giza 168) were cultivated to study the effect of silicate and phosphate ions as well as their interactions on the growth and nutritional status of the growing plants, beside their availability in the studied soil. Silicon (Si) in the form of sodium metasilicatepenta-hydrate (Na₂SiO₃.5H₂O) was added at a rate of 0, 200, 300 and 400 mg Si kg⁻¹ soil, and phosphorus (P) in the form of calcium super phosphatewas given at a rate of 0, 3.5, 7.0, 10.0 and 13.0 mg P kg⁻¹ soilto represent 0, 25, 50, 75 and 100% of the recommended rate of P fertilization by the Egyptian Ministry of Agriculture for wheat cultivation. Also, the experiment included combinations between all these concentrations of Si and P. Obtained results showed that, Si and P availability increased in the studied soil with increasing either Si or P concentrations added. This means that P availability in soil as an essential element for plant growth can be improved by addition of Si. Also, Si increased in plant with increasing applied Si concentrations. Interaction between Si and P generally increased all parameters of plant growth; such responses were significant for fresh and dry weights of wheat plants at booting stage. It could be recommended that selecting good P fertilization design, including time and rate of addition, goes along with values of available Si in the soil.

Keywords: Silicon, Phosphorus, Interactions, Wheat plant, Clay soil.

Introduction

Silicon (Si) is a basic mineral formatting element, which considered the second most abundant element in the earth's crusts (Tubana et al., 2016). It is known as non-essential element for higher plants, while lately it confirmed its role in enhancing crop yield by promoting several desirable plant physiological processes. Silicon fertilization has been reported to result in increased soil exchange capacity, improved soil aggregation; consequently improved water and air regimes, transformation of P-containing minerals and formation of alumino-silicates and heavy metal silicates (Matichenkov and Bocharnikova, 2001). The amount of literature documenting the benefits of Si on plants is vast and primarily highlights the value of Si fertilization in maintaining plant productivity under stressed conditions (Epstein, 1999 and Li et al., 2007). The established Si-induced mechanisms to improve plants' resistance to biotic and abiotic stresses take place in the soil, the root system, and inside the plant.Silicon deposition of silica in shoots and leaf epidermis, also known as the mechanical barrier hypothesis (Rodrigues et al., 2015), enhances the plant's mechanical strength and protective layer. This barrier is believed to help promote the plant's resistance to pathogens, improve plants' tolerance to abiotic stresses and increase photosynthetic rates (Datnoff et al., 2009 andRizwan et al., 2012).

Phosphorus (P) is a macro-nutrient that has the most influence on plantsgrowth and productivity, and, according to statistics of Egyptian Ministry of Agriculture, more than 50% of the cultivated areas are in P-deficient soils(Academy of Scientific Research and Technology, 2017). The low availability of P in those soils is one of the greatest challenges in the fertilization of those soils. There are various factors interfere with theavailability of P in the soil, including the environmental factors that control the activity of microorganisms which immobilize or liberate theorthophosphate ions, the physiochemical andmineralogical properties of the soil (Santos et al., 2008) and the silicon-phosphorus interactions(Carvalho et al., 2001 and Tokura et al., 2011). One of the most effective solution to raise the P availability is exploit the relationship between it and Siconcentration in the soil. Ma and Takahashi (1990) studied the effect of Si on the growth and P uptake of rice and found that, addition of Si raised the optimum P level for rice and encouraged the plant growth and yield. Alovisi et al. (2014) studied the Si-P interactions in some soils cultivated with bean plantsin Brazil and found that, application of silicate could result in the increase of P available to plants, as based on the fact that the silicate anion occupies the adsorption sites of the phosphate anion, which reflect on enhancing plant growth and yield and increasing the content of itsdifferent parts with P.

Wheat (*Triticumaestivum* L.) is the most important cereal crop in Egypt where it provides more than 30% of the population's calorie intake (Academy of Scientific Research and Technology, 2017).Works related to the use of Si to enhance P fertilization efficiency and uptake in wheat plant are rare in Egypt to date. Thus, the main objective of this investigation is to study the effect of silicate and phosphate ions as well as their interactions on the nutritional status of growing wheat plants, beside their availability in the studied clay soil.

Materials and Methods

A pot experiment was conducted in clay soil classified as *Entisol*, collected from Agricultural Research Center farm, Giza governorate, Egypt $(29^{\circ} 58' 55.046'' \text{ N} \text{ and } 31^{\circ} 12' 49.272'' \text{ E})$. Soil samples (0-30 cm) were air-dried, crushed and finely ground, then sieved to pass through a 2 mm sieve. Some physical and chemical characteristics of the studied soil samples were achieved as described by Klute (1986) and Page et al. (1982) and the results are presented in Table 1. The study was conducted by growing wheat grains (*Triticumaestivum* L., Giza 168) under open field conditions. Twenty grains were sown per pot (15.5 cm interdiameter and 21 cm height, containing 4 holes); each pot had 5 kg of the studied soil samples after preparation.

Phosphorus treatments were performed using calcium super phosphate (15% P₂O₅) which given at a rate of 0, 3.5, 7.0, 10.0 and 13.0 mg P kg⁻¹ soil to represent 0, 25, 50, 75 and 100% of the recommended rate of P fertilization by the Egyptian Ministry of Agriculture for wheat cultivation. It added before plant cultivation and mixed uniformly with the studied soil in pots. Silicon in the form of sodium metasilicatepenta-hydrate (Na₂SiO₃.5H₂O) was added at a rate of 200, 300 and 400 mg Si kg⁻¹ soil, in addition to the control treatment (without any Si additions). Silicon concentrations were applied to the soil one time after sowing the grains of wheat plants. Also, the experiment included combinations between all these concentrations. The main plots were assigned to the P additions and the subplots were occupied by the Si applications.

Nitrogen was added in the form of ammonium nitrate (33.5% N) at a rate of 100 kg N fed⁻¹ for wheat cultivation. It was added in three spilt equal doses after 2, 4 and 7 weeks from sowing. Potassium was added in the form of potassium sulfate (48% K₂O) at a rate of 50 kg K₂O fed⁻¹ for wheat plants. It was added one time after 4 weeks from cultivation. The moisture content in the studied soil were kept to the field capacityusing tap water till the end of the experiment.

Plants were harvested at booting growth stage (60 days from sowing), divided into root and shoot, and weighted the fresh and dry weights. Plant samples representing each studied treatment were oven dried at 70°C for 24 h, then ground in a stainless mill to be finally digested using a mixture of sulfuric acid and hydrogen peroxide according to the method described by Black et al. (1965). Aliquots were finally taken out and analyzed for total Si and P usingSpectrophotometer (Page et al., 1982).

After plant harvest, soil samples were subjected to analyses the availability of both studied elements. The available Si was extracted by 0.5 M acetic acid as described by Korndorferet al. (2001) and determined colorimetrically as yellow silico-molybdic acid according to Page et al. (1982). Available P was extracted by using sodium bicarbonate (NaHCO₃) 0.5 M at pH 8.5 according to Watanabe and Olsen (1965) and determined using Spectrophotometer by using ascorbic acid method.

The obtained data of soil and plant were statistically analyzed according to the method described by Gomez and Gomez (1984) using the LSD at 5% level of significance. Correlation coefficients were calculated between available Si and P in soil and their contents in plant using linear equation.

	Soluble ions (meq L^{-1})	
6.22	Ca ²⁺	4.18
2.66	Mg ²⁺	2.98
26.1	Na ⁺	2.75
65.0	K ⁺	1.91
Clay	HCO ₃ -	5.61
42.8	Cl	2.29
4.12	SO_4^{2-}	3.92
38.7	Available elements (mg kg ⁻¹))
0.96	Ν	55.2
3.33	Р	28.5
7.80	K	189
1.12	Si	268
	 6.22 2.66 26.1 65.0 Clay 42.8 4.12 38.7 0.96 3.33 7.80 1.12 	Soluble ions (meq L ⁻¹) 6.22 Ca^{2+} 2.66 Mg^{2+} 26.1 Na^+ 65.0 K^+ Clay $HCO_3^ 42.8$ $CI^ 4.12$ SO_4^{2-} 38.7 Available elements (mg kg ⁻¹) 0.96 N 3.33 P 7.80 K 1.12 Si

Table 1. Some physical and chemical characteristics of the studied soil.

Carbonate ions were not detected.

Results and Discussion

Availability of silicon and phosphorus in soil

Data presented in Table 2 showed the available concentration of Si and P in the studied clay soil at booting growth stage of wheat plants as affected by their ground application. Results revealed that available Si increased with increasing its applied dosein soil, with significant increases as compared to the control treatment. The highest increase (56%) occurred due to application of 400 mg Si kg⁻¹, as compared to control. This result

agreed with those of Lee and Kim (2007) who found that available Si in soil significantly increased with increasing the rate of silicate application and consequently affected the absorbed Si by plants. Also, the available P increased with increasing its applied concentration in soil, with significant differences as compared to the control treatment. The application of 3.5 mg P kg⁻¹ soil gave the lowest increase (10%) in available P, while the highest increase (22.9%) was given by applying the rate of 13 mg P kg⁻¹ soil as compared to control. These findings agreed with those obtained by Opala (2017) who stated that soil available P was positively significant affected by increasing P application in the soil.

Regarding the interaction between the applied Si and P concentrations in soil, data illustrated in Fig. 1 showed that, Si application increased the Si availability in soil under conditions of various P concentrations (Fig. 1A), as compared to the control treatment.Phosphorus application increased the P availability in soil under various Si concentrations (Fig. 1B), as compared to control. Phosphorus availability increased with increasing the concentration of either Si or P (Fig. 1C), as compared to control. Silicon availability increased with increasing either P or Si (Fig. 1D), as compared to control. Ma and Takahashi (1990) reported that Si can improve soil P bioavailability through sorptive interaction between the two elements and enhancement phosphate solubility.

Treatment	Available Si	Treatment	Available P
mg Si kg ⁻¹	mg Si kg ⁻¹	mg P kg ⁻¹	mg P kg ⁻¹
0.00	300	0.00	23.0
200	438	3.50	30.8
300	450	7.00	31.9
400	468	10.0	33.6
		13.0	34.4
LSD _{0.05}	13.4	LSD _{0.05}	5.90

Table 2. Available Si and P in the studied soil at booting growth stage of wheat plants as affected by theirground applications.



Fig 1. Available concentration (mg kg⁻¹) of Si and P as affected by their interactions in the studied soil at booting growth stage of wheat plants.

Vegetative growth parameters of wheat plants as affected by Si and P additions and their interactions

Data in Table 3 showed the effect of either Si or P on fresh and dry weights along with moisture content in shoots and roots of wheat plants at booting growth stage. Results showed that application of 400 mg Si kg⁻¹ soil gave the highest increase percent for fresh and dry weights of either shoots (38.1 and 48.4%, respectively) or roots of wheat plants (24.7 and 54.5%, respectively), as compared to the control treatment. These findings agreed with those obtained by Zia et al. (2017) who reported that shoot and root dry weight of wheat plants increased with Si applied. In that sense, Soratto et al. (2012) stated that Si played a favorable role in plant growth, mineral nutrition, and mechanical strength. The increase in plant growth may be attributed to the changes in physiological and morphological properties which are facilitated by the presence of Si. On contrary, the treatment of 400 mg Si kg⁻¹ soil gave the lowest value of moisture content in shoots and roots of wheat plants at booting growth stage (after 60 days from cultivation), and this confirmed increasing the dry matter content by applying this treatment. Regarding the effect of P, results in Table 3 showed that, the treatment of 13 mg P kg⁻¹ soil gave the highest % increase in the fresh and dry weights of

shoots (31 and 67%, respectively) or roots of wheat plants (78 and 154%, respectively), as compared to control. These results agreed with those obtained by Zhang et al. (2016) who reported that dry weight of shoots and roots of wheat plants increased substantially as the P application rate increased. Increasing the P application rate to the soil greatly increased root dry weight, root length and density, and consequently root surface area of the cultivated plants. Herein, Marschner (1986) stated that the increment in growth and yield due to P fertilization may be attributed to the activation of metabolic processes, where its role in building phospholipids and nucleic acid is known. Moreover, this treatment (13 mg P kg⁻¹ soil) gave the lowest value of moisture content in shoots and roots of wheat plants.

Traatmont	Fresh	weight	Dry w	veight	Moisture	e content
	g pl	g plant ⁻¹ g plant ⁻¹		g plant ⁻¹		o
ту ку	Shoots	Roots	Shoots	Roots	Shoots	Roots
Control	6.99	0.85	1.26	0.22	82.0	74.0
			Si Conce	ntrations		
Si ₂₀₀	7.25	0.88	1.36	0.25	81.0	72.0
Si ₃₀₀	8.28	0.95	1.50	0.27	82.0	72.0
Si ₄₀₀	9.65	1.06	1.87	0.34	81.0	68.0
LSD _{0.05}	0.55	0.16	0.33	0.02	3.08	7.05
			P Conce	ntrations		
P _{3.5}	8.47	1.17	1.77	0.23	79.0	80.0
P _{7.0}	8.61	1.23	1.81	0.29	79.0	76.0
P _{10.0}	9.00	1.33	2.01	0.32	78.0	70.0
P _{13.0}	9.18	1.52	2.11	0.56	77.0	63.0
LSD _{0.05}	0.90	0.22	0.30	0.06	4.81	5.35

Table 3. Fresh and dry weights of shoots and roots of wheat plants at booting growth stage as affected by Si and P applications to the studied soil.

Data in Table 4 showed that the interaction between Si and P applications, generally, increased the growth parameters of wheat plants. This may be due to some sort of synergetic interaction between them. Also, the treatment of 400 mg Si kg⁻¹ in combination with 13 mg P kg⁻¹ soil gave the highest % increase in fresh and dry weights of shoots (50 and 62%, respectively) and roots of wheat plants (109 and 186%, respectively), as compared to the

control treatment. Regarding the moisture content in shoots and roots of wheat plants at booting growth stage, there are either no significant differences among the studied treatments as compared to the control treatment or the control treatment has the highest values. In this concern, Yang et al. (2008) reported that Si promote the growth of maize seedling, dry matter accumulation of different organs and root/ shoot ratio under low P stress. Also, Owino-gerroin and Gascho (2005) found that application of silicate and phosphate increased the shoot and root dry weights of maize plants and the greatest increases were observed when silicate was applied together with phosphate.

Tracto	Treatment		weight	Dry w	Dry weight		Moisture content	
Treau	-1	g pla	g plant ⁻¹		g plant ⁻¹		%	
mg I	(g	Shoots	Roots	Shoots	Roots	Shoots	Roots	
P ₀ ×	Si ₀	6.99	0.85	1.26	0.22	82.0	74.0	
	Si ₂₀₀	7.80	0.91	1.26	0.26	82.2	71.3	
P _{3.5}	Si ₃₀₀	7.77	1.07	1.48	0.31	80.8	71.4	
	Si ₄₀₀	8.41	1.25	1.67	0.36	80.2	71.4	
	Si ₂₀₀	7.78	1.08	1.49	0.31	80.8	71.1	
P _{7.0}	Si ₃₀₀	8.12	1.24	1.55	0.34	80.9	72.9	
	Si ₄₀₀	9.77	1.48	1.82	0.45	81.4	69.5	
	Si ₂₀₀	8.43	1.21	1.67	0.37	79.8	69.3	
P _{10.0}	Si ₃₀₀	8.83	1.38	1.76	0.39	80.0	71.6	
	Si ₄₀₀	10.0	1.51	1.97	0.49	80.4	67.3	
	Si ₂₀₀	8.66	1.37	1.87	0.39	78.4	71.6	
P _{13.0}	Si ₃₀₀	9.71	1.47	1.94	0.41	80.0	72.3	
	Si ₄₀₀	10.5	1.78	2.04	0.63	80.6	64.7	
LSD	0.05	0.96	0.11	0.14	0.05	2.77	3.69	

Table 4. Effect of Si and P interactions on fresh and dry weights of shoots and roots of wheat plants at booting growth stage as affected by their applications to the studied soil.

Silicon and phosphorus concentrations in wheat plants at booting growth stage

Data in Table 5 showed that Si concentration and its total content in shoots and roots of wheat plants at booting growth stage were increased by increasing its applied concentrations in the studied soil. This response was clearer with roots than shoots. Also, data illustrated in Fig. 2A and B showed strong correlations between the available Si in soil and its concentration in both shoots and roots (R^2 = 0.93 and 0.99) of wheat plants, respectively. These results were in coincidence with those obtained by Mali and Aery (2008) who reported that there was a significant direct relationship between tissue (shoot and root) Si content and its applied rate in the soil. Sarto et al. (2014) and Zia et al. (2017) added that plant supplied with exogenous Si had high concentration of Si in shoot. Also, the accumulation of Si in shoots and roots of wheat plants increased with increasing the applied rate as compared to the control treatment. Regarding the P concentration and its total content in shoots and roots of wheat plants, data in Table 5 showed that, there were significant increases in plant organs with P which related positively with its applied concentration between available P in soil and its concentration in shoots of grown wheat plants was equal 0.83 and in roots = 0.85. These results were in harmony with those obtained by Zhang et al. (2016)who reported that the concentration of P in different plant organs increased with increasing its applied rate is applied rate in the soil.

Treatment	Concen	Concentration		ontent
ma ka ⁻¹	mg	g ⁻¹	mg pl	lant ⁻¹
nig kg	Shoots	Roots	Shoots	Roots
	Si C	oncentrati	ons	
 Si ₀	3.50	4.60	4.41	1.01
Si ₂₀₀	5.72	6.73	7.78	1.68
Si ₃₀₀	6.35	6.93	10.9	1.87
Si ₄₀₀	7.30	7.07	11.9	2.40
LSD _{0.05}	0.97	0.40	1.50	0.37
	P C	oncentrati	ons	
\mathbf{P}_0	4.05	3.90	5.10	0.86
P _{3.5}	5.80	4.40	10.3	1.01
P _{7.0}	6.00	4.70	10.9	1.36
P _{10.0}	6.80	4.90	13.7	1.57
P _{13.0}	7.50	5.30	15.8	2.97

Table 5. Concentration and total content of Si and P in shoots and roots of wheat plants at booting growth stage as affected by their applications to the studied soil.



Fig. 2. The relationship between available Si concentration (mg kg⁻¹) in the studied soil and its concentration (mg g⁻¹) in shoots and roots of the growing wheat plants.



Fig. 3. The relationship between available P concentration in the studied soil and its concentration in shoots and roots of the growing wheat plants.

Data in Table 6 showed the effect of the interaction between Si and P on concentration and total content of Si in shoots and roots of wheat plants at booting growth stage. Results showed that the treatment of 400 mg Si kg⁻¹ in combination with 13 mg P kg⁻¹ soil gave the highest concentration and total content of Si in shoots and roots of wheat plants. Similar results were found with P concentration and total content per plant as affected by the same interaction in the soil (Table 7). Data in Table 8 showed positive and, in general, strong correlation between available concentration of Si and P in the soil, and their concentrations in shoots and roots of the grown wheat plants. These data agreed with those obtained by Roy et al. (1971) who found that Si application tended to increase P content in the green tops. These data suggested that Si enhanced P mobilization from metabolically less active to metabolically more active tissue. In addition, Zia et al. (2017) found that Si application markedly increased the concentration of P in shoots. The increase in Si concentration in root medium may increase the uptake of P. It may be due to desorption of P from adsorption sites within the soil. Bai et al. (2013) reported that the critical levels of soil Olsen-P for optimal crop yield ranged from 10.9 to 21.4 mg kg⁻¹, and these levels were highly affected by crop specie, soil type, pH and soil organic matter content. While the critical levels of soil available Si extracted by 0.5 M acetic acid for rice plants ranged from 44 to 116 ppm, according toNarayanaswamy (2007). Considering these levels into account, the applied P in the studied soil till 13 mg kg⁻¹ considered adequate, while applying Si to the studied soil till 400 mg kg⁻¹considered high but not within the toxic range according to the previous studies of Alovisi et al. (2014).

Trees	Treatment mg kg ⁻¹		Concentration of Si		ntent of Si
ma			$mg g^{-1}$		plant ⁻¹
ing			Roots	Shoots	Roots
P ₀ >	<si<sub>0</si<sub>	3.50	4.60	4.41	1.01
	Si ₂₀₀	3.64	6.07	4.59	1.58
P _{3.5}	Si ₃₀₀	4.11	6.72	6.08	2.08
	Si ₄₀₀	5.67	7.32	9.47	2.64
	Si ₂₀₀	3.53	6.43	5.26	1.99
P _{7.0}	Si ₃₀₀	4.60	6.77	7.13	2.28
	Si ₄₀₀	5.89	7.63	10.7	3.43
	Si ₂₀₀	3.60	6.90	6.01	2.55
P _{10.0}	Si ₃₀₀	4.02	7.13	7.08	2.78
	Si ₄₀₀	6.04	7.69	11.9	3.77
P _{13.0}	Si ₂₀₀	6.02	6.98	11.3	2.72

Table 6. Concentration and total content of Si in shoots and roots of wheat plants at booting growth stage as affected by Si and P interactions in the studied soil.

Si ₃₀₀	6.07	7.31	11.8	3.00
Si_{400}	6.33	8.64	12.9	5.44
LSD _{0.05}	1.23	0.73	1.93	0.52

Table 7. Concentration and total content of P in shoots and roots of wheat plants at booting growth stage as affected by Si and P interactions in the studied soil.

Tract	Tractores		Concentration of P		ntent of P	
		mg	mg g ⁻¹		mg plant ⁻¹	
ing i	xg	Shoots	Roots	Shoots	Roots	
P ₀ ×	Si ₀	4.05	3.90	5.10	0.86	
	Si ₂₀₀	4.15	4.33	5.23	1.13	
P _{3.5}	Si ₃₀₀	5.37	4.54	7.95	1.41	
	Si ₄₀₀	5.88	4.51	9.82	1.62	
P _{7.0} S	Si ₂₀₀	5.10	4.77	7.60	1.48	
	Si ₃₀₀	5.53	5.49	8.57	1.87	
	Si ₄₀₀	6.12	5.22	11.1	2.35	
	Si ₂₀₀	5.66	4.80	9.45	1.78	
P _{10.0}	Si ₃₀₀	5.78	5.48	10.2	2.14	
	Si ₄₀₀	6.01	5.72	11.8	2.80	
Si ₂₀₀		5.82	5.18	10.9	2.02	
P _{13.0} Si ₃₀ Si ₄₀	Si ₃₀₀	6.05	5.07	11.7	2.08	
	Si ₄₀₀	6.41	5.08	13.1	3.20	
LSD	0.05	0.62	0.94	1.29	0.40	

Table 8. Relationship between Si and P concentrations in shoots and roots of wheat plant, and their available concentrations in the studied soil.

Treatment	\mathbb{R}^2 ((Si)	R^2	(P)
Treatment	Shoots	Roots	Shoots	Roots
Si + P _{3.5}	0.97	0.80	0.97	0.80
$Si + P_{7.0}$	0.97	0.99	0.97	0.50
$\mathrm{Si} + \mathrm{P}_{\mathrm{10.0}}$	0.99	0.98	0.80	0.93
$Si + P_{13.0}$	0.98	0.96	0.99	0.93

$P + Si_{200}$	0.50	0.90	0.80	0.63
$P + Si_{300}$	0.20	0.97	0.96	0.20
$P + Si_{400}$	0.90	0.62	0.50	0.30

Conclusion

It could be concluded that application of Si to the clay soil can release P into the soil solution and make it readily available to the plant absorption. So, it could be recommended that selecting good P fertilization design, including time and rate of addition, goes along with values of available Si in the studied soil. In addition, Si fertilization with suitable rate is very important to encourage wheat plant growth and productivity. The treatment of 400 mg Si kg⁻¹ in combination with 13 mg P kg⁻¹ soil gave the highest growth parameters and total content of both Si and P in the different wheat plants organs.

References

- Academy of Scientific Research and Technology (2017). The national campaign for upraising wheat production.<u>http://www.asrt.sei.eg/index.php/food-agriculture/132-</u> thenationalcampaign
- Alovisi, A.M.T., Neto, A.E.F., Carneiro, L.F., Curi, N. and Alovisi, A.A. (2014). Siliconphosphorus interactions in soils cultivated with bean plants. *ActaScientiarumAgron.***36**, 79-86.
- Bai, Z., Li, H., Yang, X., Zhou, B., Shi, X., Wang, B., Li, D. et al. (2013). The critical soil P levels for crop yield, soil fertilityand environmental safety in different soil types. *Plant Soil*, **372**,27-37.
- Black, C.A., Evans, D.D., White, J.L., Ensminger, L.E., Clark, F.E. and Dinauer, R.C. (1965). *Methods of Soil Analysis*. Am. Soc. Agron., Madison, Wisconsin, USA.
- Carvalho, R., FurtiniNeto, A.E., Santos, C.D., Fernandes, L.A., Curi, N. and Rodrigues, D.C. (2001).Interaçõessilício-fósforoem solos cultivados comeucaliptoem casa de vegetação. *PesquisaAgropecuáriaBrasileira*, **36**, 557-565.
- Datnoff, L.E., Rodrigues, F.A. and Seebold, K.W. (2009). Silicon and plant disease. In: *Mineral Nutrition and Plant Disease*.Datnoff, L.E., Elmer, W.H. and Huber,D.M. (Eds.). The American Phytopathological Society Press, St. Paul, MN, pp. 233–246.
- Epstein, E. (1999). Silicon. Annual Rev. Plant Physiol. Plant Mol. Biol. 50, 641-664.

- Gomez, K.A. and Gomez, A.A. (1984). *Statistical Procedures for Agricultural Research*. John Wiley and Sons, Inc., New York, USA.
- Klute, A. (1986). Methods of Soil Analysis, part I, 2nd ed., Madison, Wisconsin, USA.
- Korndorfer, G.H., Snyder, M., Ulloa, G. andDatnoff, L.E. (2001). Calibration of soil and plant silicon analysis for rice production. *J. Plant Nutr*.**24**, 1071-1084.
- Lee, Y.B. and Kim, P.J. (2007). Reduction of phosphate adsorption by ion competition with silicate in soil. *Korean J. Environ. Agric*.26, 286-293.
- Li, Q.F., Ma, C.C. and Shang,Q.L. (2007). Effects of silicon on photosynthesis and antioxidative enzymes of maize under drought stress. *Ying Yong Sheng Tai XueBao.*18, 531–536.
- Ma, J., Takahashi, E. (1990). Effect of silicon on the growth and phosphorus uptake of rice. *Plant Soil*, **126**, 115-119.
- Mali, M. and Aery, N.C. (2008). Influence of silicon on growth, relative water contents and uptake of silicon, calcium and potassium in wheat grown in nutrient solution. J. *Plant Nutr.* **31**, 1867-1876.
- Marschner, H. (1986). *Mineral Nutrition of Higher Plants*. Inst. Plant Nutr., Univ. Hohenheim, FRG Academic Press, Harcout Brace & Company, London, UK.
- Matichenkov, V.V. and Bocharnikova, E.A. (2001). The relationship between silicon and soil physical and chemical properties. *Studies Plant Sci.* **8**, 209-219.
- Narayanaswamy, C. (2007). Calibration and categorization of plantavailable soil silicon in different riceecosystems for evaluating drought and disease resistance in rice genotypes. Ph. D. Thesis, Department of Soil Science and Agricultural Chemistry, University of Agricultural Sciences, Gkvk, Bangalore.
- Opala, P.A. (2017). Influence of lime and phosphorus application rates on growth of maize in an acid soil, *Adv. Agric*.2017, 5 pages. <u>https://doi.org/10.1155/2017/7083206</u>
- Owino-Gerroh, C. and Gascho, G.J. (2005). Effect of silicon on low pH soil phosphorus sorption and on uptake and growth of maize. *Commun. Soil Sci. Plant Anal.***35**, 2369-2378.
- Page, A.L., Miller, R.H. and Keeney, D.R. (1982). *Methods of Soil Analysis*, part II, 2nd ed., Wisconsin, USA.
- Rizwan, M., Meunier, J.,Miche, H. and Keller, C. (2012). Effect of silicon on reducing cadmium toxicity in durum wheat (*Triticumturgidum* L. cv. Claudio W.) grown in a soil with aged contamination. J. Hazard. Mater.209–210, 326–334.

- Rodrigues, F.A., Resende, R.S., Dallgnol, L.J. and Datnoff, L.E. (2015). Silicon potentiates host defense mechanisms against infection by plant pathogens. In: *Silicon and Plant Disease*. Rodrigues, F.A. and Datnoff, L.E. (Eds.). Springer International Publishing, Switzerland, p. 109–138.
- Roy, A.C., Ali, M.Y., Fox, R.L. and Silva, J.A. (1971). Influence of calcium silicate on phosphate solubility and availability in Hawaiian Latosols. In: International Symposium on Soil Fertility Evaluation, New Delhi, p. 757-768.
- Santos, D.R., Gatiboni, L.C. and Kaminski, J.K (2008). Fatoresqueafetam a disponibilidade e o manejo daadubação fos fatadaem solos sob sistemaplantio direto. *Ciência Rural*, **38**, 576-586.
- Sarto, M.V.M., Lana, M.C., Rampim, L., Rosset, J.S., Wobeto, J.R., Ecco, M., Bassegio, D. and da Costa, P.F. (2014).Effect of silicate on nutrition and yield of wheat. *African J. Agric. Res.* 9, 956-962.
- Soratto, R.P., Crusciol, C.A.C., Castro, G.S.A., Costa, C.H.M. and Ferrari, J. (2012). Leaf application of silicic acid to white oat and wheat. *R. Bras. Ci. Solo*, **36**, 1538-1544.
- Tokura, A.M., FurtiniNeto, A.E., Carneiro, L.F., Curi, N., Santos, J.Z.L. and Alovisi, A.A. (2011). Dinâmica das formas de fósforoem solos de textura e mineralogiacontrastantescultivados com arroz. *ActaScientiarumAgron.*33, 171-179.
- Tubana, B.S., Babu, T. and Datnoff, L.E. (2016). A review of silicon in soils and plants and its role in us agriculture: History and future perspectives. *Soil Sci.* **181**, 1-19.
- Watanabe, F.C. and Olsen, S.R. (1965). Test of an ascorbic acid method for determining phosphorus in water and NaHCO₃ extracts from soils. *Soil Sci. Soc. Amer. Proc.*29, 677-678.
- Yang, X., Roonasi, P. and Holmgren, A. (2008). A study of sodium silicate in aqueous solution and sorbed by synthetic magnetite using in situ ATR-FTIR spectroscopy. J. Colloid Interface Sci. 328, 41-47.
- Zhang, W., Liu, D., Liu, Y., Cui, Z., Chen, Z., and Zou, C. (2016). Zinc uptake and accumulation in winter wheat relative to changes in root morphology and mycorrhizal colonization following varying phosphorus application on calcareous soil. *Field Crop Res.* **197**, 74-82.
- Zia, Z., Bakhat, H.F., Saqib, Z.A., Mustafa, G., Muhammad, S. and Ashraf, R. (2017). Effect of water management and silicon on germination, growth, phosphorus and arsenic uptake in rice. *Ecotoxicol. Environ.Safety*, 144, 11-18.