

1 Gene clone and bioinformatics analysis
2 of subtilisin-like protease in *Cucumis sativus*

13 **Background:** Cucumber target leaf spot (TLS), caused by *Corynespora cassiicola*, is a serious disease in cucumber
14 (*Cucumis sativus*) production worldwide. Therefore, cultivating new varieties of TLS resistance of *C. sativus* is an
15 important goal of cucumber breeding. Previous studies have shown that subtilisin-like protease (SUBP) plays an
16 important role in response to *C. Cassiicola* infection in resistant plants.

17 **Objective:** In this study, the full-length cDNA of the *CsSUBP* gene was cloned, and the prokaryotic expression vector
18 was successfully constructed in order to study the effects of subtilisin. Futhermore, vital clues regarding *CsSUBP*
19 gene involved in TLS resistance of *C. sativus* are gained from the bioinformatics assay.

20 **Method:** The *CsSUBP* gene was identified by sequencing with the intermediate vector pMD18 by designing specific
21 primers and PCR amplification techniques. The prokaryotic expression vector pET30a-*CsSUBP* was further
22 constructed and identified by colony PCR and *EcoR* V and *Sall* double digestion.

23 **Result:** The primary structure of *CsSUBP* was predicted and analyzed by bioinformatics analysis. The results showed
24 that *CsSUBP* was weakly acidic protein, N-terminal signal peptide region, including a Inhibitor_I9 domain domain
25 which may be the primary disease resistance domain of the protein.

26 **Conclusion:** The pET30a-*CsSUBP* prokaryotic expression vector was constructed successfully. This study is
27 convenient for the study of prokaryotic expression and its kinase activity.

28 *Key words:* Subtilisin protease; Gene clone; Bioinformatics assay; Prokaryotic expression
29

31

32 1. INTRODUCTION

33 Cucumber (*Cucumis sativus*) is one of the largest vegetable crops in China. In recent years,
34 cucumber target leaf spot (TLS), caused by *Corynespora cassiicola*, is a major disease in cucumber
35 production[1]. At present, the prevention and treatment of TLS are mainly chemical control,
36 chemical pesticide can be used to inhibit the activity of pathogens. However, *C. cassiicola* can easily
37 mutate with environmental changes and develop resistance to various chemical pesticide, so it is
38 difficult to prevent and eliminate the disease efficiently[2]. The best method for controlling the

39 disease in cucumber is the use of resistant cultivars, but few high resistant varieties of cucumber can
 40 enhance resistance of *C. cassiicola* at present. Therefore, understanding the molecular mechanisms
 41 and identifying the molecular components involved in the defense responses of the TLS resistant
 42 cucumber cultivars will provide a promising approach to restraining this disease.

43 In our previous study, a kind of subtilisin-like protease (SUBP) from resistant cucumber plants
 44 (Jingyou 38) after the fungus inoculation were explored using iTRAQ quantitative proteomics
 45 technique. It is speculated that the SUBP may play an important role in cucumber resistance to early
 46 invasion of *C. cassiicola*.

47 SUBP is a serine protease widely found in plants, bacteria, fungi, and parasites [3-8]. Such proteases
 48 have a typical Asp/Ser/His catalytic domain in structure. Studies have shown that it is closely related
 49 to the stability of the fungal phenotype and the pathogenicity of the pathogen. It plays an important
 50 immune stimulating role in the interaction between plants and pathogens [9]. SUBP is also involved
 51 in plant cell programmed cell death (PCD) processes[10]. A number of studies have demonstrated
 52 the role of SUBP in plant and pathogen interactions. For example, two subtilisin-like proteins,
 53 SAS-1 and SAS-2, in oat (*Avena sativa*) exhibit specific activity of mammalian caspase apoptosis
 54 protease and participate in the signal transduction of PCD during the non-affinity interaction of
 55 *oat-Cochliobolus victoriae*, thereby inducing HR production [11-12]. Overexpression of Arabidopsis
 56 subtilases gene *AtSBT3.3* activates downstream immune signaling and induces innate immune
 57 response in *Arabidopsis*[13].

58 In this study, the cDNA sequence of *CsSUBP* gene was cloned and the prokaryotic expression vector
 59 was constructed. The primary structure of *CsSUBP* was predicted and analyzed using bioinformatics
 60 analysis methods. On this basis, we can further explore the defense function and molecular
 61 mechanisms of SUBP against *C. cassiicola* in cucumber.

62 2. MATERIALS AND METHODS

63 2.1 Plant Materials

64 The cucumber varieties used in the experiments were Jinyou 38 (preliminary experiments showed
 65 that this variety was resistant to TLS), which were planted in a greenhouse at 28°C under 16:8
 66 light/dark cycles.

67 2.2 Full-length cDNA cloning

68 Two candidate genes were screened through preliminary test on the basis of the proteomics analysis.
 69 The cDNA sequences of *CsSUBP* (A0A0A0K993) from resistant cucumber leaves were clone. The
 70 sequence of *CsSUBP* were aligned with cucumber genome database using the service provided by
 71 <http://cucurbitgenomics.org/BLAST>. The primers (table 1) were designed by using the DNAMAN.
 72 The PCR fragments amplified from the cDNAs were cloned into TA vectors pMD18-T and were
 73 sequenced.

74 **Table 1. *CsSUBP* gene primer sequences**

Gene name	Primer sequence (5' to 3')
<i>CsSUBP-F</i>	GATATCATGTCCGGCCAATCCACAGCCCTG
<i>CsSUBP-R</i>	GTTCGACCTAATGAAGCCTTGCTGCTCCTCCT

75 2.3 Construction of recombinant prokaryotic expression vector pET30a-*CsSUBP*

76 Recombinant plasmid pMD18-T-*CsSUBP* and expression vector pET30a were digested by *EcoR* V
 77 and *Sal*I respectively. The double-digested DNA fragment had the same sticky ends as the pET30a

78 vector backbone, and the two fragments were ligated into a complete recombinant plasmid using T4
 79 ligase and ligated overnight at 16°C and then trans-formed into *E. coli* DH5α competent cells, blue
 80 and white screening. Recombinant prokaryotic expression vector was identified by direct colony
 81 PCR. Recombinant prokaryotic expression vector pET30a-*CsSUBP* plasmid was extracted and the
 82 product was digested by *EcoR V* and *Sall* respectively. It was digested for 3 h at 37°C in a constant
 83 temperature metal bath and electrophoresed on a 1.5% agarose gel. Plasmids with positive clones
 84 were sent for bioinformatics sequencing.

85 **2.4 Bioinformatics analysis**

86 The specific bioinformatics [14-17] analysis contents and tools used are as follows see table 2.

87 **Table 2 Bioinformatics analysis content and Tools website**

Analyze the content	Name of software	Bioinformatics Analysis Tools Website
Primary structure	ProtParam	http://web.expasy.org/protparam/
Secondary structure	SOPMA	http://npsa-pbil.ibcp.fr/cgi-bin/npsa_automat.pl?page=/NPSA/npsa_sopma.html
Hydrophobicity	ProtScale	http://web.expasy.org/protscale/
Signal peptide	SignalP 4.0	http://www.cbs.dtu.dk/services/SignalP/
Functional domains	SMART	http://smart.embl-heidelberg.de/


88

89 **3. RESULTS**

90 **3.1 Amplification of *CsSUBP***

91 The reverse transcription of cDNA was used as the template. PCR amplification product was
 92 subjected to 1.5% agarose gel electrophoresis and a fragment of approximately 282 bp was obtained
 93 as expected (Fig.1)

94



95


96 **Fig. 1 Electrophoresis profile of PCR products of *CsSUBP* Gene. A1-A4: *CsSUBP* PCR amplification products;**
 97 **M: DL2000 DNA Marker**

98 **3.2 Construction of Prokaryotic Expression Vector**


99 To construct the prokaryotic expression vector, we recycle the small fragments of
 100 pMD18-T-*CsSUBP* (Fig. 2, 3) and large fragments of pET30a plasmid after double digest with *EcoR*
 101 V and *SaII*. The colony PCR products of recombinant plasmid pET30a-*CsSUBP* via agarose gel
 102 electrophoresis analysis showed 282 bp of the target gene band (Fig.4, 5). Sequencing results
 103 showed that the gene fragments was consistent with the target gene sequence in GenBank (Fig. 6).
 104



105
 106 **Fig. 2 The Colony PCR of recombinant plasmid pMD18-*CsSUBP*. B1-B3,B5-B8: Colony PCR of *CsSUBP* gene;**
 107 **B4: water; M: DNA marker DL2000**
 108




109
 110 **Fig. 3 Double digestion of recombined vector pMD18-*CsSUBP* with restriction enzymes.C1-C4: Double**
 111 **digestion products; M: DNA marker DL2000**
 112



113
 114 **Fig. 4 The Colony PCR of recombinant plasmid of pET30a-*CsSUBP*. D1-D3, D6: Colony PCR products of**
 115 ***CsSUBP* gene; D4, D5: Water control; M: DNA marker DL200**

116



117

118

119

Fig. 5 Double digestion of recombinant vector pET30a-CsSUBP with restriction enzymes. E1, E2: Double digestion products; M: DNA marker DL2000

SUBP.seq	0
subp-6-M13F_-47__B04.ab1RC	GCTTTACACTTATGCTTCCGGTCCGATGTTGTGTGGAATTCGTAGGCGGATACAAITTCACACAGGAPACAGGTATGACCATGATACCAATTCGAGG	800
Consensus	
SUBP.seq	67
subp-6-M13F_-47__B04.ab1RC	ATGTCGGCCAAATCCACACCGGTGCACATCGTTTACACTGAGCGGCTCACATCAAGAACCCGAGG	900
Consensus	
SUBP.seq	167
subp-6-M13F_-47__B04.ab1RC	CCATACATCCGACCCCTACCTTCTCTTGGCAGTGAAGCGCTGCTACGGAGCGCTTGGTATAGCTATAAATGCTGCCACTGGCTTCTCTGG	1000
Consensus	
SUBP.seq	267
subp-6-M13F_-47__B04.ab1RC	TCGGCTTACCTCCGATCAGGTTGCGGACATTACCAAACACCCAGGAGTATTCACGTTGTCCAPAGCGGTAATAATAAGCTGCATTCTGGAGCAGGAGCA	1100
Consensus	
SUBP.seq	282
subp-6-M13F_-47__B04.ab1RC	GCPAGGCTTCATTAG	1167
Consensus	

120

121

122

Fig.6 Sequencing results recombinant vector pET30a-CsSUBP.

3.3 Bioinformatics Analysis of CsSUBP proteins

123

124

125

126

127

128

129


130


Some information about the CsSUBP are obtained by some online tools (shown in table2). The protein molecular weight is 23146.02, chemical formula is C₈₃₄H₁₃₈₈N₂₈₂O₃₄₈S₆₉, isoelectric point (pI) is 5.28. It is weakly acidic and hydrophobic average coefficient is 0.794. It is speculated that CsSUBP maybe a hydrophobic protein (shown in Fig 8). Neither signal peptide nor transmembrane region is found in the CsSUBP (shown in Fig 9). Secondary structure of CsSUBP is analysed by SOPMA. The result showed that CsSUBP consists of 39.78% alpha helix (h), 16.13% extended strand (e), 11.83% beta turn (t) and 32.26% random coil (c)(Fig.7 and Table 3).

131



132







133
 134 **Fig. 7 The secondary structure of CsSUBP protein. h: Alpha helix; t: Beta turn; e: Extended strand; c:**
 135 **Random coil**

136 **Table 3 Prediction the secondary structure of CsSUBP protein**

Name	Number	Percentage
Alpha helix	37	39.78%
3 ₁₀ helix	0	0.00%
Pi helix	0	0.00%
Beta bridge	0	0.00%
Extended strand	15	16.13%
Beta turn	11	11.83%
Bend region	0	0.00%
Random coil	30	32.26%
Ambiguous states	0	0.00%
Other states	0	0.00%



137
138 **Fig. 8 Prediction the hydrophobicity of CsSUBP protein**
139



140
141 **Fig. 9 Prediction the signal peptide of CsSUBP protein**

142 A major functional domains of *CsSUBP*-encoded proteins is found by SMART analysis. The results
143 are shown in Table 4.

144 **Table 4 Functional domain analysis of *CsSUBP*-encoded proteins**

Name	Position	E-value
Inhibitor_I9 domain	8(Starting site)-84(Termination site)	4.8e-15

145 Inhibitor_I9 domain, which belongs to a member of the MEROPS family of protease inhibitors I9, a
146 protease pro-peptide inhibitor (sometimes referred to as an activated peptide) that regulates the
147 folding and activity of the proenzyme or zymogen. The anterior segment extends into the enzyme
148 moiety and shields the substrate binding site, thereby promotes enzyme inhibition. Although it often
149 has lower sequence identity, several such propeptides have similar topologies. The propeptide region
150 has an open-loop, sandwich-like anti-parallel α/β , it also has two α -helices and four β -strands with a
151 $2x(\beta/\alpha/\beta)$ topology.

152 *CsSUBP* also belongs to the N-terminal propeptide domain of the meropsis family S8A, a subtilisin
153 peptidase. The use of the subtilisin propeptide as a chaperone helps to fold the mature peptidase. The

154 propeptide is removed by proteolytic cleavage to remove the activating enzyme.

155 4. CONCLUSION

156 We have successfully amplified and purified of *CsSUBP* gene in cucumber coding
sequence, and 157 constructed the recombinant prokaryotic expression vector
pET30a-*CsSUBP*. Bioinformatics 158 analysis shows that *CsSUBP* belongs to the hydrophobic
protein and has no signal peptide at the 159 N-terminus. It contains a functional domain . Its
protein chemical formula is $C_{834}H_{1388}N_{282}O_{348}S_{69}$, 160 an isoelectric point (PI) is 5.28 and a
molecular weight is 23146.02. Our study is a preparation for 161 revealing the functions and
mechanisms of *CsSUBP* against *C. cassiicola* in cucumber.

169 REFERENCES

- 170 1. Hasama W, Morita S, Kato T. Seed transmission of *Corynespora melonis*, causal fungus of target leaf spot, on
171 cucumber . *Jpn.j.phytopathol.* 2009; 59(2):175-179.
- 172 2. Qi Y X, Zhang X, Pu J J, et al. Morphological and molecular analysis of genetic variability within isolates of
173 *Corynespora cassiicola* from different hosts . *European Journal of Plant Pathology.* 2011; 130(1):83-95.
- 174 3. Siezen R J, De W V, Leunissen J A, et al. Homology modelling and protein engineering strategy of subtilases, the
175 family of subtilisin-like serine proteinases . *Protein Engineering.*1991;4(7): 719.
- 176 4.Ksiazek M, Karim A Y, Bryzek D, et al. Mirolase, a novel subtilisin-like serine protease from the
177 periodontopathogen *Tannerella forsythia*. *Biological Chemistry.* 2015; 396(3): 261-275.
- 178 5. Duan X, Zhang Z, Wang J, et al. Characterization of a novel cotton subtilase gene *GbSBT1* in response to
179 extracellular stimulations and its role in *Verticillium* resistance . *Plos One.* 2016; 11(4): e0153988.
- 180 6. Meyer M, Huttenlocher F, Cedzich A, et al. The subtilisin-like protease SBT3 contributes to insect resistance in
181 tomato . *Journal of Experimental Botany.* 2016;67(14): 4325-4338.
- 182 7. Mizutani O, Shiina M, Yoshimi A, et al. Substantial decrease in cell wall α -1, 3-glucan caused by disruption of
183 the *kexB* gene encoding a subtilisin-like processing protease in *Aspergillus oryza e*. *Bioscience, biotechnology, and*
184 *biochemistry.* 2016; 80(9): 1781-1791.
- 185 8. Pepe A, Frey M E, Munoz F, et al. Fibrin (ogen) olytic and antiplatelet activities of a subtilisin-like protease from
186 *Solanum tuberosum* (StSBTc-3). *Biochimie.*2016;125: 163-170.
- 187 9. Figueiredo A, Monteiro F, Sebastiana M. Subtilisin-like proteases in plant-pathogen recognition and immune
188 priming: a perspective . *Front Plant Sci.*2014;5(739): 739.
- 189 10. Vartapetian A B, Tuzhikov A I, Chichkova N V, et al. A plant alternative to animal caspases: subtilisin-like
190 proteases. *Cell death and differentiation.* 2011; 18(8): 1289.
- 191 11. Rantong G, Evans R, Gunawardena A H. Lace plant ethylene receptors, AmERS1a and AmERS1c, regulate
192 ethylene-induced programmed cell death during leaf morphogenesis . *Plant molecular biology.* 2015; 89(3): 215-227.
- 193 12. Coffeen W C, Wolpert T J. Purification and characterization of serine proteases that exhibit caspase-like activity
194 and are associated with programmed cell death in *Avena sativa*. *The Plant Cell.* 2004; 16(4): 857-873.
- 195 13. Ramírez V, López A, Mauch-Mani B, et al. An extracellular subtilase switch for immune priming in *Arabidopsis* .
196 *PLoS Pathogens.* 2013;9(6): e1003445.

- 197 14. Zhang B, Pan X, Cannon C H, et al. Conservation and divergence of plant microRNA genes . The Plant Journal.
198 2006; 46(2): 243-259.
- 199 15. Zhang B, Wang Q, Wang K, et al. Identification of cotton microRNAs and their targets. Gene. 2007; 397(1):
200 26-37.
- 201 16. Yin Z, Li C, Han X, et al. Identification of conserved microRNAs and their target genes in tomato(*Lycopersicon*
202 *esculentum*) . Gene. 2008; 414(1): 60-66.
- 203 17. Sunkar R, Jagadeeswaran G. In silico identification of conserved microRNAs in large number of diverse plant
204 species . BMC Plant Biology. 2008; 8(1):37.