

Original Research Article

Nematode Parasites of Anurans from Three Cocoa farms in Ondo State, Nigeria.

Abstract

The research was designed to investigate the parasitic fauna of Anurans from cocoa farms in Ondo state. Amphibians are one of the most threatened groups of vertebrates. Many reasons are attributed to the decline of amphibian species such as global warming, habitat destruction and modification, others include: exploitation, pesticide use, introduced species, ultraviolet-B radiation (UV-B), pollution, parasites and diseases. A total of 31 frogs from 4 genera, *Hemisus*, *Ptychadena*, *Rana* and *Xenopus* and 7 toads from 1 genus, *Sclerophrys* were examined. 9 frogs were collected from Oluwateru farm at Iwoye Village; 7 frogs were collected from Folorunso farm at Ako-Igbatoro and 15 frogs from Obodulu farm in Idanre. 5 toads were collected from Oluwateru farm at Iwoye Village and 2 toads from Obodulu farm in Idanre. In all cases collection was done between 20:00hrs and 05:00hrs. Collected specimens were transported in sealed but ventilated containers to the laboratory where identification was done to species level. The frogs were anaesthetized until death in absolute chloroform soaked in cotton wool placed inside kill-jar for 3 minutes in the laboratory. The gastrointestinal tracts were cut open and the contents of the various sections were put into separate Petri dishes containing normal saline. The skin and the bladder were observed directly under a dissecting microscope for the presence of cysts and monogeneans. The parasites were fixed and preserved in 70% alcohol following standard procedure. Parasites recovered from the gastrointestinal tracts of the anurans include *Cosmocerca ornata*, Deising, 1861, *Cosmocerca cummutata*, Diesing, 1851 *Paracosmocerca mucronata*, Kung and Wu, 1945, *Ampliceacum africanum*, Taylor, 1924, *Gendria liberrei* Bain and Philippon, 1969 and *Chenospirura asturi* Hsu, 1957 Others were *Procamallus brevis* Kung, 1948 and *Camallanus dimotrivii* Durette-Desset and Batcharov, 1974. Some of the parasites are zoonotic while a few others are established parasites of African fishes and water Birds raising probable public health concerns from the findings. Further works aimed at unravelling the biodiversity of hosts and parasites in the lush ecosystem of Ondo state, as well as identification of organisms involved in the life cycle are noted.

Key words: Anurans, Parasitic fauna, Cocoa farms, Ondo State, Nigeria

Introduction

Amphibians are a class of tetrapods that evolved from lobe-finned fish and primitive tetrapods about 340 million years ago (San Mauro *et al.*, 2005). Anurans are usually less than 65cm in length and most species breed in aquatic environments. The most common of them in tropical

Africa are the Frogs (Ford and Cannatella, 1993; Rodel, 2000). They can be herbivorous or omnivorous and are consumed by both vertebrates and invertebrates. They are also used as pest controller and play an important role as bio-indicators (Gonwouo and Rodel, 2008). A parasitic organism lives on or in another organism (host) and obtains its food, protection, transportation and also performs its essential metabolism through the host. Monogeneans, common parasites of fishes, may externally infect aquatic life stages of amphibians. Some cestodes, acanthocephalans, and hirudineans may also reside on or in adults anurans, generally as internal infections (Johnson *et al.*, 1999; Poynton and Whitaker, 2001). Many types of helminths may infect amphibians, but many nematodes or roundworms, are common helminths that infect amphibians from egg to adult stages and affect a variety of organs and tissues. Anurans have the capacity to carry extremely high parasite loads. As a resource for parasitological studies, there has been a number of significant papers and reviews of parasite groups of amphibians over the past century (Baylis, 1929; Kerve, 1930; Walton, 1932; Walton, 1933; Southwell and Kirshner, 1937; Bain and Phillipon, 1969; Kiesecker, 2002; Ford *et al.*, 2004). Some reports had been written about the parasites of Amphibians in Nigeria by Thurston (1967, 1970), Avery (1970), Jackson and Tinsley (1995a, 1995b) and Aisien *et al.* (2001, 2003, 2004, 2009, 2010), but not as extensive as in some parts of the world. Interestingly, tropical Nigeria has a limited number of publications in this field and none seem to have been done in Ondo State, which is a typical rainforest region of the country. In this region, many humans consume some of the amphibian species and the latter can also be found in some relative abundance despite the associated threats. This present research investigated the nematode parasites of anurans found in some Cocoa farms in Ondo State, Nigeria.

Materials and Methods

Ondo State is situated in the south western part of Nigeria with geographical coordinates of 5°45'N, 4°20'E and 7°52'N, 6°05'E (Wikipedia, 2014). The state is bordered by Ekiti State in the north, Osun State by the west, Edo State at the eastern end, Ogun State and the Atlantic Ocean in the southern area. The study sites were cocoa farms with fresh flowing stream, making them good sites for amphibian habitat. The three cocoa farms where amphibians were collected include:

- a. Oluwateru Family Farm at Iwoye Village (7°25'N, 5°20'E), situated in Akure South Local Government Area (LGA), about 10 kilometres outside Akure township towards Ondo town
- b. Folorunso Family Farm at Ako-Igbatoro Village (7°09'N, 5°37'E), situated in Akure South LGA about 5 kilometres along Igbatoro road from Akure metropolis
- c. Obodulu Cocoa farm (7°24'N, 5°19'E), situated in Idanre LGA, around a rocky farmland in Idanre

The frogs and toads were collected between 20:00 and 05:00hrs in ponds, streams, underneath leaf litters and on trees. The specimens were handpicked and transported in sealed but ventilated containers to the Laboratory. Each container held specimens of averagely same size to prevent injury or death resulting from aggression. Safety precautions were put in place particularly

against snake bites during collection by wearing thick boots and usage of hand gloves. Other measures included usage of whistles by all on site, keeping of bitter kola in the pockets (it is believed by the locals that it scares away snakes), usage of back packs instead of hand bags, sticks to remove leaf litters coupled with sharpened machetes and torch for proper illumination. A local who is very familiar with each respective site was engaged as a guide. An average of 12 specimens were collected per site. For proper identification, the specimens were first anaesthetized for 3 minutes in absolute chloroform soaked in cotton wool inside a kill-jar in the Laboratory. The smooth vein length (SVL) of each of the specimens were measured for taxonomic reasons and the sex of the animals were determined. The specimens were identified to the species level using identification keys by Rodel (2000). The specimens were examined for parasites 12 to 18 hours after collection. Dissections of the specimens were done 3-5 minutes after anaesthetizing so as to recover live parasites. The various sections of the gastrointestinal tract were cut out systematically i.e. the Stomach, Oesophagus, and the intestine and put inside separate saline solutions in Petri dishes. The skin and the bladder were observed directly under a dissecting microscope to view the presence of monogeneans and cysts. The organs were teased using dissecting needle to facilitate the escape of the parasites into normal saline, then the Petri dishes were examined under a dissecting microscope. The parasites were lifted off the saline solution using Forceps/Pasteur pipette and placed inside another petri dish of saline solution before they were fixed for observation. The parasites were fixed by placing each of them inside small sterilized stainless steel vials, containing 70% alcohol and the container heated to make the parasite stretch out from the usual coiled position. (Aisien *et al.*, 2001; 2009; 2010). The preservation of the parasites was done by removing them from the hot alcohol and placing them inside vials containing 70% alcohol. The recovered nematodes were cleared in lactophenol (Aisien *et al.*, 2001; 2009) followed by examination under the dissecting microscope. The nematodes were identified using taxonomic keys provided by Yamaguti (1961). The prevalence rate was calculated as a percentage of the number of a particular host species infected with the specific helminth parasite divided by the total number of hosts examined, and mean intensity of infection was taken as the total number of parasites per host, and this was done for the whole animal population collected either infected or uninfected (Anderson, 1993).

Ethical Considerations: Care were taken not to sacrifice more animals than necessary for the research following the standard procedures as established by the International Society of Applied Ethology (ISAE, 2002; Sherwin *et al.*, 2003). In addition to this, the study conformed with the practice of reducing the number of amphibians used in research work to the smallest minimum possible as proposed by Herpetological Animal Care and Use Committee (HACC) of the American Society of Ichthyologists and Herpetologists (Beaupre *et al.*, 2004).

Results

A total of 31 frogs and 7 toads were collected from the sites. The species of frogs encountered during the study include: *Ptychadena longirostris* Peters, 1870, *Ptychadena mascareniensis* Dumeril and Birron, 1841, *Rana galamensis* Dumeril and Bibroni, 1841, *Ptychadena retropunctata* Angel, 1949, *Xenopus muelleri* Peters, 1844, *Hemisus marmoratus* Peters, 1855, *Ptychadena bibroni* Hallowell, 1845 and *Ptychadena pumilio* Boulenger, 1920. The toads encountered were *Sclerophrys maculata* Hallowell, 1854 and *Sclerophrys pentoni* Anderson, 1893.

The gut contents of *Rana galamensis* and *Ptychadena* spp were filled mostly with grasses and insects while *Xenopus muelleri* and *Hemisus marmoratus* had Tadpoles in their gut. The contents were from the Oesophagus and the stomach while in the intestine, liquid to semi liquid matter was seen and could not be traced to any specific food substance. The gut contents of the *Sclerophrys* spp were insects e.g. grasshoppers, bugs and crickets. The contents were from the Oesophagus and the stomach while in the intestine, liquid or semi liquid matter were seen and could not be traced to any specific type of food substance. The gut contents were examined so as to help in knowing possible intermediate hosts of the parasites encountered. The parasites found in the alimentary canal of the frogs and toads were measured to help in taxonomic description and recorded (Table 1). They include: *Cosmocerca cummutata* Diesing, 1851, *Cosmocerca ornata*, *Procamallus brevis* Kung, 1948, *Camallanus dimotrivii* Durette- Desset and Batcharov, 1974, *Chenospirura asturi* Hsu, 1957, *Gendria liberrei*, *Paracosmocerca mucronata* Kung and Wu, 1945, *Ampliceacum africanum* Taylor, 1924 and Ascaridoid larvae Blanchard, 1849. The parasites were all nematodes belonging to 3 Orders and 5 families (Fig. 1). Frogs in Idanre were more infected than the other locations as highlighted in Table 2 and Iwoye had the highest number of infected toads (Table 3). The mean intensity was generally between 0.75-3.75 except for *Hemisus marmoratus* that had about 86 Ascaridoid larvae per host (Table 4). The overall prevalence for frogs recorded showed that *Sclerophrys maculata* had the highest infection rate in the study.

Discussion

7 toads from genus *Sclerophrys* and 31 frogs belonging to 4 genera, *Hemisus*, *Ptychadena*, *Rana* and *Xenopus* were examined out of the several seen for conservation reasons. The prevalence of infection in the observed species when compared to earlier reports in tropical Nigeria (Aisien *et al.*, 2001 and 2009) showed that the frogs in this study had low worm burden. Previous studies from Nigeria also recorded the presence of cestodes and trematodes but only nematodes were encountered in this study.

Table 1: Parasitic Species Recovered from Frogs and Toads from Different Sites

Parasites	Host		Site
	Frog	Toad	
<i>Ampliceacum africanum</i>	Nil	<i>Sclerophrys maculata</i> ; <i>Sclerophrys pentoni</i>	Stomach, Oesophagus, Small intestine and Body Cavity
<i>Chenospirura asturi</i>	Nil	<i>Sclerophrys maculata</i>	Small intestine
<i>Camallanus dimitrovi</i>	Nil	<i>Sclerophrys maculata</i> ; <i>Sclerophrys pentoni</i>	Stomach and Small intestine
<i>Cosmocerca cummutata</i>	<i>Ptychadena pumilio</i> ; <i>Xenopus muelleri</i>	<i>Sclerophrys maculata</i>	Small intestine
<i>Cosmocerca ornata</i>	<i>Ptychadena pumilio</i> ; <i>Hemissus memoratus</i>	nil	Small intestine
<i>Procamallus brevis</i>	Nil	<i>Sclerophrysmaculata</i>	Stomach and Small intestine
<i>Gendria liberrei</i>	Nil	<i>Sclerophrysmaculata</i>	Small intestine
<i>Chenospirura asturi</i>		<i>Sclerophrysmaculata</i>	Body Cavity
<i>Paracosmocerca mucronata</i>	Nil	<i>Sclerophrys pentoni</i>	Body cavity
Ascarididoid larvae	<i>Hemissus memoratus</i> ; <i>Ptychadena bibroni</i>	nil	Small intestine

Table 2: Prevalence and Mean Intensity of Infection in Frogs at the Different Sites

Parasites	Host	Location					
		Iwoye		Ako-Igbatoro		Idanre	
		Prevalence	Mean intensity	Prevalence	Mean intensity	Prevalence	Mean Intensity
<i>Cosmocerca</i>	<i>Xenopus muelleri</i>	100%	1.00	-	-	-	-
<i>cummutata</i>	<i>Ptychadena pumilio</i>		-	-	-	100%	2.00
<i>Cosmocerca</i>	<i>Ptychadena pumilio</i>	-	-	-	-	100%	2.00
<i>ornata</i>	<i>Hemissus memoratus</i>	-	-	-	-	50%	0.5
Ascarididoid	<i>Hemissus memoratus</i>	-	-	-	-	100%	86
larvae	<i>Ptychadena bibroni</i>	-	-	-	-	25%	0.25




Fig 1:Pedigree of the Nematode Parasites encountered in the Anuran hosts

Adapted using taxonomic keys from Yamaguti (1961)

Table 3: Prevalence and Mean Intensity of Infection in Toads at the Different Sites

Parasites	Host	Location			
		Iwoye		Idanre	
		Prevalence	Mean intensity	Prevalence	Mean Intensity
<i>Ampliceacum africanum</i>	<i>Sclerophrys maculata</i>	50%	0.75	50%	0.50
	<i>Sclerophrys pentoni</i>	-	-	100%	1.00
<i>Camallanus dimitrovi</i>	<i>Sclerophrys maculata</i>	50%	3.75	-	-
	<i>Sclerophrys pentoni</i>	-	-	100%	3.00
<i>Cosmocerca cummutata</i>	<i>Sclerophrys maculata</i>	25%	1.75	-	-
<i>Procamallus brevis</i>	<i>Sclerophrys maculata</i>	25%	0.75	-	-
<i>Gendria liberrei</i>	<i>Sclerophrys maculata</i>	25%	0.50	-	-
<i>Chenospirura asturi</i>	<i>Sclerophrys maculata</i>	25%	1.25	-	-
<i>Paracoscocerca mucronata</i>	<i>Sclerophrys pentoni</i>	.	-	100%	1.00

Table 4: Overall Prevalence of infection in the examined Animals

	Host Species									
	<i>Sclerophrys maculata</i>	<i>Sclerophrys pentoni</i>	<i>Xenopus muelleri</i>	<i>Hemius memoratus</i>	<i>Rana galamensis</i>	<i>P. bibroni</i>	<i>P. pumilio</i>	<i>P. retropunctata</i>	<i>P. longritoitis</i>	<i>P. mascarensis</i>
<i>Cosmocerca cummutata</i>	-	-	11.11%	-	-	-	28.57%	-	-	-
<i>Cosmocerca ornata</i>	-	-	-	33.33%	-	-	14.29%	-	-	-
Ascarididoid larvae	-	-	-	33.33%	-	20%	-	-	-	-
<i>Ampliceacum africanum</i>	50%	100%	-	-	-	-	-	-	-	-
<i>Camallanus dimitrovi</i>	33.33%	100%	-	-	-	-	-	-	-	-
<i>Procamallus brevis</i>	16.67%	-	-	-	-	-	-	-	-	-
<i>Cosmocerca cummutata</i>	16.67%	-	-	-	-	-	-	-	-	-
<i>Cosmocerca ornata</i>	-	-	-	-	-	-	-	-	-	-
Ascarididoid larvae	-	-	-	-	-	-	-	-	-	-
<i>Gendria liberrei</i>	16.67%	-	-	-	-	-	-	-	-	-
<i>Chenospirura asturi</i>	16.67%	-	-	-	-	-	-	-	-	-
<i>Paracosmocerca mucronata</i>	-	100%	-	-	-	-	-	-	-	-

Two of the parasites *Gendria liberrei* and *Chenuspirura asturi* have been reported before in tropical Africa but are reported for first time in Nigeria in this study. The remaining parasites have been reported before in tropical Africa and in Nigeria. *Cosmocerca cummutata* recovered from the intestine of female *Ptychadena pumilio* in the Idanre farm had been reported in Brazil and Europe as a parasite of North American frogs (Walton, 1933), it was reported in Congo (Vuylesteke, 1964), and in Northern Nigeria as parasite of *Xenopus muelleri* (Avery, 1970). Same parasite was also reported in Sudan as a parasite of *Sclerophrys regularis* and *Dicroglossus occipitalis* (Pike, 1979) and in Turkey as a parasite of the tree frog, *Hyla arborea* (Dusen and Oz, 2004). The report of this parasite in *Ptychadena pumilio* is new for the parasite.

Cosmocerca ornata was recovered from the gut of male *Hemissus memoratus* from the Idanre farm. Puylaert (1970), reported it in Senegal, Avery (1971), reported it in Nigeria from *Xenopus* spp, Baker (1981), observed it in South Africa from the native South African frogs. Pike (1981), reported it from *Dicroglossus occipitalis* and *Sclerophrys regularis* in Sudan, Moravec *et al.* (1987), described it in their findings in Amphibian and reptile parasites in Egypt while Moravec and Barus (1990), encountered and reported it in Zambia and Uganda. Aisien *et al.* (2001 and 2009), also reported it in Nigeria from *Sclerophrys regularis*, *D. occipitalis*, *X. muelleri* and from *Hemissus memoratus*.

Paracosmocerca mucronata was gotten from the intestine of the only *Sclerophrys pentoni* encountered in the study, Aisien *et al.* (2001), reported it in Nigeria as a parasite of *Xenopus muelleri*, this is the second time this parasite will be reported in another part of the Country. *Chenuspirura asturi* was recovered from *Sclerophrys maculata* in Iwoye farm. Hsu (1957), reported the parasite as a parasite of water Birds. *Camallanus dimitrovi* recovered from *Sclerophrys pentoni* has been reported extensively in the West African axis of the tropics. Durette-Desset and Bacharov (1974), reported it in Togo in a general review of amphibian parasites, Jackson and Tinsley (1995a) reported it in Nigeria from *Xenopus* spp and Aisien *et al.* (2001 and 2009) reported it in Nigeria from *Dicroglossus occipitalis* and from *Hoplobatrachus occipitalis*. *Procamallanus brevis* recovered from *Sclerophrys maculata* in this research was first reported in Tanzania (Baylis, 1929) and Avery (1971) reported it in the northern part of Nigeria and in both cases it was recovered from *Xenopus* spp.

Ampliceacum africanum, recovered from *Sclerophrys maculata* and *Sclerophrys pentoni*, was first reported from the mountainous region of the present day Tanzania from *Sclerophrys maculata* (Baylis, 1929). Baker (1987), reported it in his synopsis of the nematode parasitic in amphibians and reptiles, from some East and West African countries and Aisien *et al.* (2001), established its occurrence in Nigeria in *Sclerophrys maculata* and *D. occipitalis*. *Gendria liberrei* was recovered from *Sclerophrys maculata*. Bain and Phillipon (1969), reported the latter in Togo as a parasite of tilapia fish. This is the first time the parasite will be reported in Nigeria.

The life cycle of *Cosmocerca cummutata*, *Cosmocerca ornata*, and *Paracosmocerca mucronata* of the family Oxyuridae starts by the female producing thousands of eggs in the large intestine of

its host making its host's rectum itch. The host scratches the area and transfers the eggs to the mouth where they travel to the intestine. Another way is by retrofection where eggs that are not transferred to extremities will hatch and crawl back into the intestines (Schmidt and Roberts, 1989). Olsen (1986), reported that *Chenuspirura asturi* a Spiruridae, undergoes indirect life cycle by using an arthropod intermediate host, most especially bugs or grasshoppers, while frogs and other vertebrates serve as definitive hosts.

The life cycle of *Camallanus dimitrovi* and *Procamallus brevis* of family Camallanidae involves a cyclopoid copepod crustacean as an intermediate host where development continues in the intestines of a vertebrate namely freshwater fishes and turtles (Schmidt and Roberts, 1989). Females with fully developed first-stage larva burst from cheeks of a definitive host, releasing the larva, which are eaten by copepods that are then eaten by a definitive host (Olsen, 1986). Fertilization occurs when migrating from intestines of the definitive host to its head, after which all the males die (Schmidt, 1992). *Ampliceacum africanum* and Ascarididoid larvae of the family Quimperiidae undergo viviparous direct life cycle in the stomach of the host. *Gendria liberrei* lays egg inside the host (Olsen, 1986) and the eggs of this nematode can be effectively transferred from its natural host to a paratenic host i.e. amphibians and reptiles through the bite of Black flies which dwells in fast flowing streams with the natural hosts (fishes) Bain and Philippon (1969).

The reported pathological effects of the parasites vary. Olsen (1986), reported that *Cosmocerca cummutata* and *Paracosmocerca mucronata* changes the host's colour, causes ulceration and corrosion of alimentary canal. *Chenuspirura asturi* escalates the effects of other helminths and obstructs the intestinal passage, leading to serious mechanical damages (Olsen, 1986). *Camallanus dimitrovi* causes lesions, haemorrhage, mechanical damage, and associated diseases (Olsen, 1986; Schmidt and Roberts, 1989). The recorded pathological effects of *Procamallanus brevis* are lesions, haemorrhage, mechanical damage, and associated diseases whereas, *Ampliceacum africanum*, causes major lesions and mechanical obstruction leading to a kwashiorkor like appearance of the host (Olsen, 1986). Ascarididoid larvae causes varying degree of mechanical damages depending on the numbers and stages of development while *Gendria liberrei* has no recorded pathological effect on the hosts.

Some of the observed parasites are zoonotic. *Cosmocerca cummutata*, *Paracosmocerca mucronata*, *Chenuspirura asturi* and *Ampliceacum africanum* are zoonotic (Olsen, 1986). *Gendria liberrei* generally are parasites of African fishes and associated animals. Dwellers and Farmers in the farm areas are prone to zoonotic infections since they get in contact with the anurans through various activities like farming, hunting (some of the anurans are edible), fishing in the streams especially during rainy seasons.

The current study has described for the first time, the parasitic fauna of amphibians from the tropical rainforest of Ondo State, Nigeria and probable public health concerns from the findings.

Further works aimed at unravelling the biodiversity of hosts and parasites in the lush ecosystem, as well as identification of organisms involved in the life cycle continues.

References

- Aisien, M. S. O., Du Preez, L. H., Imasuen, A. A. (2010) *Polystoma okomuensis* sp. (Monogenea: Polystomatidae) from Boulenger's striped frog, *Phlyctimantis boulengeri* (Perret, 1986) in Nigeria. *Journal of Helminthology*: 1-7.
- Aisien, S. O., Ajakaiye, F. B and Braimoh, K (2003) Helminth fauna of anurans from the savannah-mosaic zone of south-western Nigeria. *Acta Parasitologica*. 48: 47-54.
- Aisien, S. O., Ayeni, F and Ilechie, I. (2004) Helminth fauna of anurans from the Guinea savanna of new Bussa, Nigeria. *Africa Zoology*. 39:133-136.
- Aisien, S. O., Ogoannah, S. O and Imasuen, A. A. (2009). Helminth parasites of amphibians from a rainforest reserve in south western Nigeria. *Africa Zoology*. 44(1): 1-7.
- Aisien, S. O., Ugbo, A. D., Ilavbare, A. and Ogunbor, O. (2001). Endoparasites of Amphibians from South-Western Nigeria. *Acta Parasitologica*. 46(4): 299-305.
- Anderson, R. M. (1993). Epidemiology. In: *Modern Parasitology, A textbook of Parasitology*, (ed.) F.E.G. Cox, Blackwell Scientific, London: 75-116.
- Avery, R. A. (1971). A preliminary list of parasites collected from reptiles and amphibians in Northern Nigeria. *British Journal of Herpetology*. 4: 217-219.
- Bain, O and Philippon, B (1969) Researches on larvae of Ascaridida Nematodes found in *Simolium dimnosum*. *Annals de parasitologie*(Paris). 44: 147-156
- Bain, O and Philippon, B (1969) Researches on larvae of Ascaridida Nematodes found in *Simolium dimnosum*. *Annals de parasitologie*(Paris). 44: 147-156
- Baylis, H. A. (1929). Some parasitic Nematodes from Uluguru and Usambara Mountains, Tanganyika territory, Tanzania. *Ann and Magazine of Natural History*. 10 (4): 372-391.
- Baylis, H. A. (1929). Some parasitic Nematodes from Uluguru and Usambara Mountains, Tanganyika territory, Tanzania. *Ann and Magazine of Natural History*. 10 (4): 372-391.
- Beaupre, S. J., Jacobson, E. R., Lillywhite, H. B. and Zamudio, K. (2004). Guidelines for use of live amphibians and reptiles in field and laboratory research. Second Edition, Revised by the Herpetological Animal Care and Use Committee (HACC) of the American Society of Ichthyologists and Herpetologists: 1-42.
- Durette-Desset, M., and Bacharov, G. (1974). 2 Nematodes parasites of Amphibians from Togo. *Annals de parasitologie Humaine et Comparee*. 49 (5), 567-576.
- Dusen, S. and Oz, M. (2004). Helminth parasites of the tree frog, *Hyla arborea* (Anura: Hylidae) from Southwest Turkey. *Comparative Parasitology*. 71(2): 258-261.
- Ford, L.S. and Cannatella, D. C. (1993). The major clades of frogs. *Herpetological Monographs* 7: 94-117.
- Ford, T. R., Dillehay, D. L. and Mook, D. M. (2004). Cutaneous ascariasis in the African clawed frog (*Xenopus laevis*). *Comparative Medicine*. 54(6): 713-717.

- Gonwouo, L. N. and Rödel M. O. (2008). The importance of frogs to the livelihood of the Bakossi people around Mount Manengouba, Cameroon, with special consideration of the Hairy Frog, *Trichobatrachus robustus*. *Salamandra*. 44: 23–34.
- ISAE (2002). Guidelines for Ethical treatment of Animals in Applied Animal Behaviour and Welfare Research, Prepared by International Society of Applied Ethology (ISAE) Ethical Committee. [http://www.researchgate.net/publications/401541403_ISAE_2002_Proceedings_of_the_36th International Congress of the ISAE](http://www.researchgate.net/publications/401541403_ISAE_2002_Proceedings_of_the_36th_International_Congress_of_the_ISAE). Retrieved 2015-01-02
- Jackson, J. A and Tinsley, R. C. (1995a). Evolutionary relationships, host range and Geographical Distribution of *Camallanus* Railliet and Henry 1956 species (Nematoda; Camallanidae) from the clawed toads of the genus *Xenopus* (Anura; Pipidae). *Systematic Parasitology*. 32:1-21.
- Jackson, J. A and Tinsley, R. C. (1995b). Representatives of *Batrachocamallanus* n. g. (Nematoda: Procamallaninae) from *Xenopus tropicus*. (Anura: Pipidae): geographical distribution, host range and evolutionary relationships. *Systematic Parasitology*. 32: 1-21
- Jackson, J. A and Tinsley, R. C. (1998a). Hymenochirine anurans (Pipidae) as transport hosts in camallanid nematode life cycles. *Systematic Parasitology*. 39: 141-151.
- Jackson, J. A and Tinsley, R. C. (1998b). Paramphistomedigenaeans from *Xenopus* spp (Pipidae) in Africa: taxonomy, host specificity and biogeography. *Systematic Parasitology*. 40: 143-160.
- Jackson, J. A. Tinsley, R. C. Du Preez, L. H and Henderson, A. C (2001). A redescription of *Chabaudus liberrei* (Bain and Philippon, 1969) (Nematoda: Seuratoidea) from *Xenopus* spp. in Swaziland. *Systematic Parasitology*. 50(2):81-89.
- Johnson P. T. J., Lunde, K. B., Ritchie, E. G, Launer, A. E. (1999). The effect of trematode infection on amphibian limb development and survivorship. *Science*. 284:802–804.
- Karve, J. N., (1930). Some parasitic nematodes of toads and frogs. *Annual Tropical Medicine Parasitology*. 24(4) 481-491.
- Kiesecker, J. M. (2002). Synergism between trematode infection and pesticide exposure: a link to amphibian limb deformities in nature? *Proceedings of the National Academy of Sciences of the United States of America*. 99(15): 9900-9904.
- Olsen, O.W. (1986). *Animal Parasites: Their Biology and Life Cycles*. 3rd Edition. New York: Courier Dover Publications: 44-468
- Pike, A.W. (1979). Helminth parasites of the amphibians *Dicroglossus occipitalis* (Gunthor) and *Bufo regularis* (Reuss), in Khartoun, Republic of Sudan. *Journal of Natural History*. 13: 337-376.
- Poynton, J. A and Whiteaker, B. R. (2001) Protozoa and metazoan infecting amphibians In Wright K. M Witetaker B. R., eds . *Amphibian Medicine and captive Husbandary*. Malabar F. L: Krieger Publishing Company: 193-221
- Rodel, M. (2000). Herpetofauna of West Africa. In: *Amphibians of West Africa vol 1*, (ed.) Micheal Meyers: 36-267.
- San Mauro, D., Vences, M., Alcobendas, M., Zardoya, R., and Meyer, A. (2005). "Initial Diversification of living amphibians predated the breakup of Pangaea". *The American Naturalist*. 165 (5): 590–599.

- Schmidt, G. D. & Roberts, L. S. (1989). *Foundations of Parasitology*: Fourth Edition. St. Louis: Times Mirror/Mosby College Publishing: 80-102
- Sherwin, C. M., Christiansen, S. B., Duncan, I. J. H., Erhard, H. W., Lay, D. C., Mench, J. A., O'Connor, C. E. and Petherick C. J. (2003). Guidelines for the ethical use of animals in applied animal behaviour research. *Applied Animal Behaviour science*. 81: 291-305.
- Southwell, T. and Kirschner, A. (1937). On some parasitic worms found in *Xenopus laevis*, the South Africa clawed toad, *Annals of Tropical Medicine and Parasitology*. (31). 245-265
- Thurston, J.P. (1967). The morphology and life cycle of *Cephalochlamys namaquensis* from *Xenopus muelleri* and *Xenopus laevis*. *Parasitology*. (57): 187-200
- Thurston, J.P. (1970) Studies on some protozoans and helminth parasites of *Xenopus*, the African Clawed toad. *Revue de Zoologie et de Botanique Africaines*. (82): 349-369
- Vuylesteke, C. (1964). Mission, de Zoologie in'edicala au Maniema (Congo, L'opddville). Vermes Nematoda, *Annals Mus. R. Afr. Cent Se'r*. 3 (4): 41-66.
- Walton, A. C. (1932). A new Nematode (*Camallanus multiriga*) parasite in a West Africa frog. *Annal magazine of natural history*. 9: 151-154.
- Walton, A. C. (1933). The nematode as parasites of amphibians. *Journal of Parasitology*. 20 (1): 1-43.
- Yamaguti S. (1961). *Systema Helminthum*. Volume III. The nematodes of vertebrates part I and II. Interscience Publishers, Inc., New York: 1- 600