

2  
3 **Acid Phosphatases from the Breadfruit**  
4 ***Artocarpuscommunis* Seeds as Novel Plant**  
5 **Phosphorylating Biocatalysts**

6  
7  
8  
9  
10 **ABSTRACT**  
11

**Aims:** Investigation on the phosphotransferase activity of two non-specific acid phosphatases (EC 3.1.3.2) designated as AP1 and AP2, previously isolated from breadfruit (*Artocarpuscommunis*) seeds for further biotechnological and industrial applications.

**Methodology:** Transphosphorylation reactions were tested with sodium pyrophosphate as the phosphoryl donor and phenol as its acceptor. Transfer products were quantified by using high performance liquid chromatography.

**Results:** The two acid phosphatases were able to catalyse phosphoconjugates synthesis using pyrophosphate as the phosphoryl donor and phenol as acceptor. The optimal conditions of transphosphorylation reactions indicated that this synthesis was highly dependent on pH, temperature, time course, donor and acceptor concentrations and enzyme amount. A very short period (1.25 h) was observed for these synthesis reactions catalysed by acid phosphatases isolated from breadfruit (*Artocarpuscommunis*) seeds. This suggested energy saving during biotransformation processes. The high average yields of 84.20 and 66.78 % obtained for AP1 and AP2, respectively, made them useful to phosphorylate a wide range of nucleophile compounds such as nucleotides often used as food additives and pharmaceutical intermediates.

**Conclusion:** The acid phosphatase AP1 would be the most promising on the basis the better synthesis product yield (84.20 %). The two biocatalysts could be considered as new valuable tools for bioprocesses.

12  
13 *Keywords: Plant acid phosphatases, transphosphorylation, breadfruit Artocarpuscommunis*  
14 *seeds, phosphoconjugate synthesis, biocatalysts*

15  
16  
17 **1. INTRODUCTION**

18  
19 Transphosphorylation is an efficient process for the large-scale production of new  
20 phosphorylated conjugates with improved properties [1, 2, 3]. This reaction is essentially  
21 reversible, and the equilibrium position depends on the conditions and the amount of  
22 reagents and products present in the reaction mixture [1].

23 Kinases are well-known phosphorylating enzymes which transfers a phosphate unit from  
24 ATP to a variety of acceptors, but the large-scale application is impeded by the need of  
25 regenerating ATP and in addition these enzymes are specific for the substrate to be  
26 phosphorylated [4]. Nowadays, it has been shown that acid phosphatases (orthophosphoric-  
27 monoester phosphohydrolase (EC 3.1.3.2)) catalyzing the hydrolysis of a broad and  
28 overlapping range of phosphomonoesters are also able to carry out transphosphorylation  
29 reactions in which a phosphate unit is transferred from a donor (phosphomonoesters or  
30 pyrophosphate P<sub>Pi</sub>) to an acceptor alcohol [5]. This enzymatic phosphorylation process

31 shows more advantages than the chemical one that uses phosphoryl chloride (POCl<sub>3</sub>) [6]. It  
32 is well known that enzymes tolerate environmental stress and thanks to their broad substrate  
33 specificity, they are able to accept a wide range of cheap phosphoryl donors and acceptors  
34 [6, 7]. In contrast, the chemical methods should be explored with caution due to high number  
35 of by-products obtained in typical reactions. Current chemical routes to phosphate esters  
36 synthesis often proceed via the corresponding phosphate di or triesters and require harsh  
37 reagents and conditions leading to hardly separable product mixtures [8].  
38 Thus, Bacterial phosphatases from the class A nonspecific acid phosphatase family were  
39 mostly used, and *Shigella flexneri* [7, 9], *Salmonella enterica* [7], *Morganella morganii* [5, 1]  
40 and *Escherichia coli* [2] were prominent sources of the enzyme. The  
41 phosphoryltransferase activities of nonspecific acid phosphatases from these  
42 microorganisms were exploited to produce nucleotides as inosine-5 monophosphate (5-IMP)  
43 and guanosine-5 monophosphate (5-GMP) using pyrophosphate (PP<sub>i</sub>) as a phosphate  
44 donor. So, this production of phosphoconjugates is another key area of interest in  
45 biotechnology, because they have often used as food additives and as pharmaceutical  
46 synthetic intermediates [5].  
47 To our knowledge, there are few reports on the transphosphorylation abilities of plant  
48 nonspecific acid phosphatases. Nevertheless, due to their wider availability and lesser cost,  
49 plant phosphatases could be explored as good substitutes for biotechnological phosphate  
50 ester synthesis. Thus, Koffi et al. [3] isolated four non-specific acid phosphatases from the  
51 neglected crop *Lagenaria siceraria* seeds exhibiting high phosphoryltransferase activities. In  
52 search of new biocatalysts with improved transphosphorylation yield, two acid phosphatases  
53 from breadfruit (*Artocarpus communis*) seeds [10] were investigated. In this paper, we report  
54 on the phosphoryltransferase activity of these enzymes.

55  
56  
57

## 58 **2. MATERIAL AND METHODS**

59  
60

### 60 **2.1 Enzymes**

61 The acid phosphatases AP1 and AP2 (EC 3.1.3.2) used for transphosphorylation reactions  
62 originated from the purified enzyme collection of Laboratoire de Biochimie et Technologies  
63 des Aliments, UFR Sciences et technologies des Aliments, Université Nangui Abrogoua  
64 (Abidjan, Côte d'Ivoire). These phosphatases were purified as described previously [10].

65  
66

### 66 **2.2 Chemicals**

67 Sodium pyrophosphate (donor), phenol (acceptor) and tyrosine (internal standard) were  
68 purchased from Sigma Aldrich. All other reagents used were of analytical grade.

69  
70

### 70 **2.3 Transphosphorylation reactions**

71 The ability of phosphatases AP1 and AP2 from *A. communis* seeds [10] to catalyse  
72 transphosphorylation reactions was tested with sodium pyrophosphate as the phosphoryl  
73 donor and phenol as its acceptor. In typical experiment, transphosphorylation reactions were  
74 carried out at 37 °C in a total reaction mixture of 250 µL containing 400 mM of sodium  
75 acetate buffer pH 4.0. The reactions were stopped by immersion in boiling water for 3 min  
76 followed by cooling in ice bath. Prior to each HPLC analysis, the reaction mixtures were  
77 filtered using Ultrafree-MC filter (0.45 µm) devices (Millipore). Tyrosine was used as the  
78 internal standard to correct chromatographic product areas. Twenty microliter (20 µL)  
79 aliquots of each reaction mixture always containing the internal standard (2 mM final  
80 concentration) were analysed quantitatively by HPLC. The column used was

81 SPHERECLONE 5  $\mu$  ODS (2) (250 mm  $\times$  4.60 mm; Phenomenex) and phenolic compounds  
82 were detected at 254 nm with a SPECTRA SYSTEM UV 1000 detector. The elution was  
83 done with a BECKMAN 114 M solvent delivery module pump, at a flow rate of 0.5 mL min<sup>-1</sup>  
84 by using a degassed mixture of  
85 acetonitrile/water in the ratio 50:50 (v/v) as mobile phase. The chromatograms were  
86 obtained with a SHIMADZU C -R8A CHROMATOPAC V1.04 integrator. The detailed  
87 experimental conditions for studying parameters likely to affect the transphosphorylation  
88 reactions (pH, time, donor concentration, acceptor concentration and enzyme amount) are  
89 given below. For the determination of each parameter, reactions were carried out in  
90 triplicate.

#### 91 **2.4 Determination of optimum pH**

93 The pH values were determined at 25 °C. For determination of optimum pH,  
94 transphosphorylation reactions were performed by incubating at 37 °C for 1 h each  
95 phosphatase (25  $\mu$ L) in a pH range of 3.6 to 5.6 (sodium acetate buffer, 400 mM), with 100  
96 mM final concentration of sodium pyrophosphate and 5 mM final concentration of phenol.

#### 97 **2.5 Determination of optimum time**

99 To determine the optimum time of transphosphorylation, the optimum pH determined for  
100 each enzyme was fixed. The transphosphorylation reactions were performed at different  
101 times ranging from 0 to 6 h (at 37 °C), in 400 mM sodium acetate buffer at appropriate  
102 optimum pH, with 25  $\mu$ L enzyme, 100 mM final concentration of sodium pyrophosphate and  
103 5 mM final concentration phenol.

#### 104 **2.6 Determination of phosphoryl donor optimum concentration**

106 The influence of phosphoryl donor concentrations (0 to 300 mM) on the transphosphorylation  
107 reactions was determined under the optimum conditions of pH and time. Phosphatases (25  
108  $\mu$ L) were separately incubated at 37 °C, with different concentrations of sodium  
109 pyrophosphate ranging from 0 to 300 mM final concentration and 5 mM phenol.

#### 110 **2.7 Determination of phosphoryl acceptor optimum concentration**

112 The influence of phosphoryl acceptor concentrations (0 to 10 mM) on the  
113 transphosphorylation reactions was determined under the optimum conditions of pH, time  
114 and phosphoryl donor concentration. The phosphatases (25  $\mu$ L) were separately incubated  
115 at 37 °C, with different concentrations of sodium pyrophosphate ranging from 0 to 300 mM  
116 final concentration and 5 mM phenol.

#### 117 **2.8 Determination of enzyme amount**

119 For this study, the optimal condition of pH, time, donor concentration and acceptor  
120 concentration determined for each enzyme were fixed. Only the amount of each  
121 phosphatase varied from 0 to 10  $\mu$ g. In the determinations described above, 25  $\mu$ L of  
122 enzyme corresponded to 5  $\mu$ g, and 2.1  $\mu$ g of protein, respectively, for AP1 and AP2. All the  
123 reactions were stopped by immersion in boiling water for 3 min and the products quantified  
124 by HPLC as described in the typical transphosphorylation reaction.

#### 125 **2.9 Transphosphorylation reactions in optimum conditions**

126

127 Ultimately, the optimal conditions of pH, enzyme amount, donor and acceptor concentrations  
128 and time were fulfilled to perform a unique transphosphorylation reaction with phosphatases  
129 AP1 and AP2 purified from *A. communis* seeds. These reactions were also carried out in  
130 triplicate with regard to the typical conditions described in the other experiments, and the  
131 synthesized products were quantified by HPLC as described previously.  
132

### 133 **2.10 Estimation of the yield of transphosphorylation**


134 One mol of the synthesized product corresponds to 1 mol of phenol used as acceptor in the  
135 reaction mixture. Therefore, the starting phenol concentration deficit at the end of each  
136 reaction is typical of the amount of acceptor necessary for the product synthesis. After  
137 adjusting areas with the internal standard (tyrosine), transphosphorylation yields were  
138 determined as follows: (area of initial phenol – area of remaining phenol)/area of initial  
139 phenol × 100.  
140

## 141 **3. RESULTS AND DISCUSSION**

142  
143 The two non-specific acid phosphatases AP1 and AP2 purified from *A. communis* seeds [10]  
144 were tested in transphosphorylation reaction of pyrophosphate at pH 4. The retention time of  
145 the newly synthesized product was found to be similar to that of phenylphosphate (Data not  
146 shown). This shows clearly that the studied enzymes were able to catalyse the  
147 phosphorylation of pyrophosphate (PPi) to phenylphosphate. Pyrophosphate was earlier  
148 used as donor in many transphosphorylation reactions [3, 11] due to the fact that it is a safe  
149 and inexpensive compound which can be used in large excess for various manufacturing  
150 processes [12]. Also, this molecule is easily synthesized from phosphate groups [13].  
151 Therefore, an efficient phosphorylation process could be achieved by recycling  
152 pyrophosphate to phosphate group by-products for further transfer in transphosphorylation  
153 reactions.

154 In order to optimize the phenylphosphate formation five parameters were investigated: the  
155 pH, time, donor concentration, acceptor concentration and enzyme amount dependency.

156 In Figure 1 the pH dependency of the phosphotransferase reaction is depicted. The highest  
 157 yield of phenylphosphate (69.25 and 56.75 % respectively for AP1 and AP2) was obtained at  
 158 pH 3.6. It is noteworthy that there are few reports on phosphotransferase activity of plant



159 acid phosphatases, but it seems that these biocatalysts exhibit better transfer activity at low  
 160 pHs compared to their hydrolysis activity (optima pHs sited between 5 and 6). A similar  
 161 behaviour has been observed by KOFFI et al. [3] for four non-specific acid phosphatases  
 162 isolated from the cucurbit *L.siceraria* seeds. According to Huber et al. [14], the dissimilarity  
 163 between the optima pHs for phosphorylation and dephosphorylation could be due to ionized  
 164 groups sited in the active site of the enzymes which would favour synthesis and  
 165 disadvantage hydrolysis in more acidic conditions.

166


167 **Fig. 1. Effect of pH on transphosphorylation reaction catalyzed by the acid**  
 168 **phosphatases (AP1 and AP2) from breadfruit (*A.communis*) seeds**

169 Bars represent  $\pm$  SE

170

171 The enzyme activities were also measured at various temperatures from 30 to 65°C. AP1  
 172 had maximum activity at 40°C whereas AP2 shows maximum activity at 35°C. In these  
 173 conditions, optima yields of 73.58 and 59.75 % were obtained for phosphatases AP1 and  
 174 AP2, respectively (Figure 2). The optimal temperatures of transphosphorylation were lower  
 175 than that obtained for dephosphorylation (55°C). So, these temperatures would be  
 176 advantageous since the studied enzymes have been earlier shown to be stable at  
 177 temperatures below 55°C [10]. Asano et al. [5] obtained their best transphosphorylation rate  
 178 at 40°C during the synthesis of Inosine-5'-monophosphate using a phosphorylating enzyme  
 179 isolated from bacteria *M. morgani*. The general observation would be that both the  
 180 phosphorylation and dephosphorylation temperatures do not coincide. It appears that the  
 181 optimum phosphorylation temperature is generally lower (ranging from 30 to 40°C)  
 182 compared to that of dephosphorylation (above 45°C).


183



184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201

**Fig. 2. Effect of temperature on transphosphorylation reaction catalyzed by the acid phosphatases (AP1 and AP2) from breadfruit (*A. communis*).**  
Bars represent  $\pm$  SE


The time course of phenylphosphate synthesis by acid phosphatases AP1 and AP2 isolated from bread fruit *A. communis* seeds is depicted in Figure 3. The two enzymes show maximum transphosphorylation yield at 1.25 h. This time course of reaction is very low compared with those obtained for 5'-inosinic acid synthesis as reported by Mihara et al. [2]. These authors have obtained their better phosphotransferase activity at time courses ranging from 18 to 36 h. Ishikawa et al. [12] have reported phosphorylation times sited between 5 and 15 h for inosine-5'-monophosphate (5'-IMP) synthesis. Therefore, the very short period of synthesis reactions catalysed by the studied acid phosphatases may constitute an interesting feature insofar as this would make it possible to save energy in biotransformation processes using these biocatalysts. Also, it should be remembered that this reaction time is very well included in the stability time (fully active for more than 150 min) of these enzymes as describe by Konan et al. [10].



202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221

**Fig. 3. Time course of phenylphosphate synthesis by the acid phosphatases (AP1 and AP2) from breadfruit (*A. communis*) seeds**  
Bars represent  $\pm$  SE


The yields of phenylphosphate produced by nonspecific acid phosphatases AP1 and AP2 were highly dependent on donor and acceptor concentration (Figures 4 and 5). Similar behaviours were observed by Van Herk et al. [9]. In our study, the maximum concentrations of phosphorylated product were obtained from 200 mM pyrophosphate and 12.5 mM phenol for both AP1 and AP2. In these conditions, optimum yields of 82.39 and 65.71 % were obtained for AP1 and AP2, respectively. As presented in figures 4 and 5, it seems that higher concentrations of donor and acceptor than those obtained in this study would inhibit the phosphoryltransferase activities. This would explain the decrease in the transfer rate since it is well known that high substrate loads are required to achieve high product titers, while kinetic control is crucial to keep the phosphorylation/hydrolysis ratio as high as possible due to the competition between hydrolysis and transphosphorylation in aqueous medium [15]. A common mechanistic feature of phosphate-transferring phosphatases is the formation of a covalent phospho-enzyme intermediate, which undergoes either nucleophilic attack by water (hydrolysis) or accepts a suitable nucleophile (transphosphorylation).



222  
223  
224  
225  
226  
227  
228

**Fig. 4. Effect of phosphoryl donor (sodium pyrophosphate) on transphosphorylation reaction catalyzed by the acid phosphatases (AP1 and AP2) from breadfruit (*A. communis*) seeds**

Bars represent  $\pm$  SE




229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245

**Fig. 5. Effect of phosphoryl acceptor (phenol) on transphosphorylation reaction catalyzed by the acid phosphatases (AP1 and AP2) from breadfruit (*A. communis*) seeds**

Bars represent  $\pm$  SE.

Figure 6 shows effect of enzyme amounts on transphosphorylation reactions. This result indicates that high amounts of enzymes would be unfavourable to the phosphoryl transfer reaction. It appears that these high enzyme amounts result in the release of high amounts of inorganic phosphate, which could inhibit the phosphatase [16]. The phosphotransferase activities were optimal with enzyme amounts of 6  $\mu$ g for both AP1 and AP2, and average yields of 83.60 and 65.10 % were respectively obtained.



247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

**Fig. 6. Effect of enzymatic unit on transphosphorylation reaction catalyzed by the acid phosphatases (AP1 and AP2) from breadfruit (*A. communis*) seeds.**

Bars represent  $\pm$  SE.

Table 1 summarized the average yields of transfer at the different stages as well as those obtained under the optimal conditions. During this study, yields significantly improved (about 10 to 15 %). In optimal conditions yields of 84.20 and 66.78 % were obtained for AP1 and AP2, respectively. These values are higher than those reported for nonspecific acid phosphatases from seeds of the orphan crop *L. siceraria* [3] and for the nonspecific acid phosphatase from *Salmonella typhimurium* LT2 [17]. Although they are plant acid phosphatases, the studied enzymes exhibit good transphosphorylation potentials compared to bacterial phosphoryl transfer enzymes which were often modified to improve their transferase properties. Regarding these interesting phosphoryltransferase activities, it is likely that nonspecific acid phosphatases from the breadfruit *A. communis* seeds would be useful to phosphorylate a wide range of nucleophile compounds such as nucleotides often used as food additives and pharmaceutical intermediates.

**Table 1. Transfer product yields in optimal conditions of the transphosphorylation assay catalyzed by acid phosphatases AP1 and AP2 from breadfruit (*A. communis*) seeds**

Parameters	Transphosphorylation yield	
	AP1	AP2
pH	69.25 $\pm$ 0.8	56.75 $\pm$ 1.1
Temperature	73.58 $\pm$ 0.7	59.75 $\pm$ 2.2
Reaction time	76.98 $\pm$ 1.2	60.50 $\pm$ 0.9
Donor	79.08 $\pm$ 1.5	62.47 $\pm$ 0.9
Acceptor	82.39 $\pm$ 0.5	65.71 $\pm$ 0.7
Enzyme unit	83.60 $\pm$ 0.9	65.10 $\pm$ 1.0
Optimum conditions	84.20 $\pm$ 0.6	66.78 $\pm$ 0.5

269

270

271

272

273

Values given are the average from at least three experiments

274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324

#### 4. CONCLUSION

The data obtained in this study showed that acid phosphatases from seeds of the breadfruit *A. communis* may be good alternative biocatalysts to form phosphoconjugates by applying them in synthesis of various phosphorylated compounds such as nucleotides often used as food additives and pharmaceutical intermediates. Shorter time course reactions (1.25 h at the most) were observed for optimum synthesis, suggesting energy saving during biotransformation processes. The acid phosphatase AP1 would be the most promising on the basis the better synthesis product yield (84.20 %). The two biocatalysts could be considered as new valuable tools for bioprocesses.

#### REFERENCES

- [1] Mihara Y, Utagawa T, Yamada H, Asano Y. Acid phosphatase/ phosphotransferases from enteric bacteria. *J BiosciBioeng*2001; 2:50-54.
- [2]Mihara Y, Ishikawa K, Suzuki E, Asano Y. Improving the Pyrophosphate-inosine Phosphotransferase Activity of *Escherichia blattae* Acid Phosphatase by Sequential Site-directed Mutagenesis. *BiosciBiotechnolBiochem* 2004; 68(5): 1046-1050.
- [3]Koffi DM, Faulet BM, Gonnety JT, Bédikou ME, Kouamé LP, Zoro Bi IA, Niamké SL. Novel Plant Acid Phosphatases from the Orphan Crop *Lagenariasiceraria*(Molina) Standl. for Phosphate Ester Synthesis. *Philip Agric Scientist* 2012; 95(1): 14-21.
- [4] Faber K. *Biotransformation in organic chemistry* 2000; pp 123–134, 4th edition, Springer, Berlin.
- [5] Asano Y, Mihara Y, Yamada H. A novel selective nucleoside phosphorylation enzyme from *Morganella morganii*. *J BiosciBioeng*1999; 87: 732-738.
- [6] Yoshikawa M, Kato T, Takenishi T. Studies of phosphorylation. III. Selective phosphorylation of unprotected nucleosides. *Bull Chem Soc Jpn* 1969; 42: 3505-3508.
- [7] Tanaka N, Hasan Z, Hartog AF, Van Herk T, Wever R, Sanders RJ, Phosphorylation and dephosphorylation of polyhydroxy compounds by class A bacterial acid phosphatases. *Org Biomol Chem* 2003;1: 2833-2839.
- [8]Tasnádi G, Lukesch M, Zechner M, Jud W, Hall M, Ditrich K, Baldenius K, Hartog AF, Wever R, Faber K. Exploiting Acid Phosphatases in the Synthesis of Phosphorylated Monoalcohols and Diols. *Eur J Org Chem* 2016; 2016(1): 45-50.
- [9] Van Herk T, Hartog AF, Van Der Burg AM, Wever R. Regioselective phosphorylation of carbohydrates and various alcohols by bacterial acid phosphatases; probing the substrate specificity of the enzyme from *Shigella flexneri*. *Adv Synth Catal* 2005; 347: 1155-1162.
- [10] Konan HK, Yapi DYA, Yué Bi CY, Koné TFM, Kouadio PEJN, Kouamé LP. Biochemical Characterization of Two Acid Phosphatases Purified from Breadfruit (*Artocarpuscommunis*) Seeds. *Chiang Mai J Sci* 2016; 43(5): 1102-1113.
- [11]Babich L, Peralta JLVM, Hartog AF, Wever R. Phosphorylation by Alkaline Phosphatase: Immobilization and Synthetic Potential. *Int J Chem* 2013; 5(3): 87-98.
- [12] Ishikawa K, Mihara Y, Shimba N, Ohtsu N, Kwasaki H, Suzuki EI, Asano Y. Enhancement of nucleoside phosphorylation activity in an acid phosphatase. *Protein Eng* 2002; 15(7): 539-543.
- [13] Staffel T. *Condensed sodium phosphates* 1991. Weinheim: VCH Verlag.
- [14] Huber RE, Gaunt MT, Sept RL, Babiak MJ. Differences in the effects of pH on the hydrolytic and transgalactosylic reactions of beta-galactosidases (*Escherichia coli*). *Can J Biochem Cell Biol* 1983; 61: 198-206.

- 325 [15]Mihara Y, Utagawa T, Yamada H, Asano Y. Phosphorylation of nucleosides by the  
326 mutated acid phosphatase from *Morganella morganii*. Appl Environ Microbiol 2000; 66:  
327 2811-2816.
- 328 [16] Femley HN, Walker PG. Studies on alkaline phosphatase. Inhibition by phosphate  
329 derivatives and substrate specificity. Biochem J 1967; 104: 1011-1018.
- 330 [17] Dissing K, Uerkvitz W. Class B nonspecific acid phosphatase from *Salmonella*  
331 *typhimurium* LT2 phosphotransferase activity, stability and thiol group reactivity.  
332 Enzyme Microb Technol 2006; 38: 683-688.  
333