

# Using Plant growth-promoting fungi (PGPF), as a biofertilizer and biocontrol agents against *Tetranychus cucurbitacearum* on Nubian watermelon (*Citrullus lanatus* L.)

## ABSTRACT

Plant growth-promoting fungi (PGPF) have attracted considerable interest as biofertilizers and biocontrol due to their multiple beneficial effects on plant quantity and quality as well as their positive relationship with the ecological environment. The objective of this study was to determine the efficacy of different concentrations 25, 50, 75 and 100% from cultural filtrate of *Trichoderma viride* and *T. harzianum* to induce the two-spotted spider mite, *Tetranychus cucurbitacearum* (*In vitro*), and their ability to improve the growth dynamics of Nubian watermelon plants in field experiment during two growing summer seasons of 2017 and 2018.

In general the effect of tested concentrations of *T. viride* were non effective on egg deposition by adult females after five days from treatment, while in *T. harzianum*, the concentration 75% was the most effective than the other concentrations. Also, egg hatchability % decreased with increased of two egg age (1 – 2 day old) at treatment by concentrations 25 and 50% of both fungal. In field experiment, the test compound (vertimec) was the more effective against egg stage of spider mite *T. cucurbitacearum* than motile stages of both *T. viride* and *T. harzianum* treatments. Also, plants inoculated with *T. harzianum* showed increases in vegetative growth parameters included numbers of leaves, Leaf dry weight, stem length and numbers of branches and biochemical analysis of leaves included chlorophyll content and percentages of NPK at 30 and 60 days from sowing during both seasons compared to uninoculated control plants. Also, enzymes activities, treatment T2 (inoculated with *T. viride*) recorded the highest values at all growth stages, which recorded 155.77, 257.29 and 114.62 mg TPF g<sup>-1</sup> soil day<sup>-1</sup> for dehydrogenase and 113.79, 201.03 and 115.24 mg NH<sub>4</sub><sup>+</sup>- N g<sup>-1</sup> soil d<sup>-1</sup> for urease at 30, 60 and harvest during 2017 growing season, respectively. The same trend was observed in total count of fungi during both seasons.

For fruit yield, *T. viride* (T2) had significantly the highest number of fruits per plant, number of seeds per fruit, fruit weight (g) and dry weight of 100 seeds (g) which recorded 1.92, 273.07, 1126 g and 306.29 g as compared to untreated control treatment, which attained 1.21, 185.08, 526.66 g and 14.41 g at 2017 season, respectively. Therefore, these results reflected to increase fruit yield (Kg/m<sup>2</sup>), seed yield (g/m<sup>2</sup>) and weight of yield (ton fed.<sup>-1</sup>) during both seasons.

**Keywords:** PGPF; *Trichoderma viride*; *Trichoderma harzianum*; *Tetranychus cucurbitacearum*; Nubian watermelon; yield

## INTRODUCTION

Nowadays, agricultural sector has been facing the destructive activities of numerous pests like insects and weeds leading to significant decrease in many agricultural crops [1]. For example, in insect pests the two-spotted spider mite, *Tetranychus cucurbitacearum*, consider a polyphase pest which can infest the plants and pierces the epidermis and feeds on the contents of mesophyll cells that results in chlorosis and due to a decrease in total chlorophyll content and an eventual loss of photosynthetic capacity and sometimes the plants will often die [2, 3].

To cope that pest, various approaches can be used such as chemical control which it is better to use some chemicals in controlling these pests. However, it considers expensive and not environmental friendly. So, many researchers have used biological control by use of living organisms to reduce damage caused by pests and diseases [4]. Recently, the use of biological control agents such as parasites, predators, bacteria or fungi has been encouraged and found to be an efficient method to

reducing the harmful effects of pest insects on many crops. Therefore, it considers an effective and do not leave any harmful effect on environment [5, 6].

During the last 20 years, a series of studies about using of fungi as a Plant Growth Promoting Fungus (PGPF) such as species belonging to the genera *Trichoderma*, *Fusarium*, *Penicillium*, and *Phoma*. PGPF are nonpathogenic saprophytes and are reported to suppress fungal, bacterial diseases and insects of a number of crop plants [7]. The PGPF association with roots of various plant species and infection has also been shown to modulate growth, morphology, nitrogen assimilation, resource allocation and mineral uptake of the host plant and also improves host reproductive fitness by enhancing plant growth, increase biomass and grain yield of crop plants i.e. rice, wheat, maize and barley [8, 9, 10, 11]. For biocontrol agent on the two-spotted spider mite and it was observed significantly reduced plant damage. Fungal biocontrol agents are important natural regulators of insect populations which have potential as mycoinsecticide agents against diverse insect pests in agriculture. These fungi infect their hosts by penetrating through the cuticle, gaining access to the hemolymph, producing toxins, and grow by utilizing nutrients present in the haemocoel to avoid insect immune responses. The using of fungi as alternative to insecticide could be very useful for insecticide resistant management [12].

Nubian Watermelon (*Citrullus lanatus*) is one of the cucurbitaceae family which cultivated from early times in Egypt [13] and Egypt is the fifth country worldwide in the production [14]. Nubian Watermelon is a favorite and popular consuming crop and it can be used as animal feeds (green parts) and the seeds are used as snacks, as well as the residues are used as a source of heat energy for cooking [15]. Also, it can be used as a source of protein supplement to ruminant animals and a new source of vegetable oil [16, 17]. [18] Studied the relationship between the colonization of *Lotus japonicus* by plant growth-promoting fungi including *T. koningi*, *Fusarium equiseti*, and *Penicillium simplicissimum* and biosynthesis of the isoflavonoid phytoalexin vestitol and they found that only *T. koningi* colonized the roots long-term, suppresses isoflavonoid phytoalexin vestitol production and increased plant dry weight reached to 126% as compared to other tested fungi. [19] evaluated the role of *Fusarium solani* strain K (FsK) in altering plant responses to the two spotted spider mite *Tetranychus urticae* in tomato plants and they found that spider mite performance was negatively affected on FsK-colonized plants and feeding damage was lower on these compared to control plants. Also, FsK-colonization led to increased plant biomass to both spider mite-infested and un-infested plants and enhance indirect tomato defense as FsK-colonized plants attracted more predators than uncolonized plants.

So, the aim of this study was to determine the efficacy of different concentrations from cultural filtrate of *T. viride* and *T. harzianum* against *T. cucurbitacearum* (*In vitro*), and their efficacy of inoculation for improving vegetative growth and yield of Nubian Watermelon.

84

85

## MATERIALS AND METHODS

### Laboratory study:

#### Organism and culture conditions

*Trichoderma viride* and *Trichoderma harzianum* were provided from Bacteriology Laboratory, Sakha Agricultural Research Station, Kafr El-Sheikh. The fungal strains were cultured on potato dextrose broth (PDB) for 15 days at 25°C. Then, centrifuged at 10.000 rpm for 20 min, and the culture medium was discarded. Next, the supernatant was filtered by passing the culture broth through a sterile membrane filter (0.2 µm) [20].

94

95

96

### Mite culture:

98 Colonies of the two-spotted spider mite, *Tetranychus cucurbitacearum* (Acarina. *Tetranychidae*), were collected from castor bean plants at the experimental farm of Sakha Agricultural Research Station, Kafr El-Sheikh Governorate and reared under laboratory conditions ( $25 \pm 2^\circ \text{C}$  and  $65 \pm 5\%$  RH) on potato leaves away from any contamination with pesticides.

### Effect of cultural filtrates of fungal strains on *T. cucurbitacearum* egg laying zero time pretreated leaf discs during 5 days

105 The effects of tested cultural filtrates of fungal strains (*T. viride* and *T. harzianum*) to the spider mite, *T. cucurbitacearum* were evaluated by the leaf disc dip technique according to [21]. Four concentrations (25, 50, 75 and 100%) of the tested cultural filtrates were prepared. Four discs of castor bean leaves were dipped in each concentration for 5 seconds and left to dry. Ten adult female mites were transferred on each disc after treatment zero time. Accumulative number of eggs laid was assessed 24, 48, 72, 96 and 120 hours later. All treatments were conducted at ( $25 \pm 2^\circ \text{C}$  and  $65 \pm 5\%$  RH). Each treatment was replicated four times.

### Effect of some concentrations of cultural filtrates of fungal strains on hatchability of *T. cucurbitacearum* egg at two ages

115 To get homogenous with the same age, five adult females of *T. cucurbitacearum* were transferred to potato leaves discs on wet cotton pad in petri-dishes to lay eggs for 24 hours, and then removed. Each treatment involved 120 eggs at age of 1 day old and 120 eggs of 2 day old in addition to the control with the same number. All treatments were dipped in four concentrations of each cultural filtrate of fungal isolates for 5 seconds as well as in water as control. The hatchability was assessed five days after treatments and the reduction in hatchability was counted for each cultural filtrate.

### Field experiment:

123 An experiment was carried out at Sakha Agricultural Research Station farm to evaluate the effect of inoculation with *T. viride* and *T. harzianum* for growth promotion of Nubian watermelon plants during summer seasons of 2017 and 2018 and their effects to control motile stages and egg stage of *T. cucurbitacearum* with compare vertimec (Abamectin 1.8 EC) as acaricide compound during summer season of 2018.

128 Seeds of Nubian watermelon var. Colocynthoide were obtained from Horticultural Research Institute, ARC, Egypt. Area of the experimental plot was ( $42 \text{ m}^2$  each), and sown with seeds on 25/4/2017 for first season and 22/4/2018 for second season with spacing 40 cm. The inoculation treatments were prepared as peat-based inoculums, 15 ml of culture per 30 g of sterilized carrier and mixed with the seeds before sowing using a sticking material. Four replicates were used for each treatment in a complete randomized block design and the experimental soil analyses during the two growing seasons were shown in Table 1. Plants were thinned to two plants per hill after three weeks of planting. Treatments were as follows:

- T1: Control (no inoculation).
- T2: Inoculation with *T. viride* ( $1 \times 10^7$  spores  $\text{mL}^{-1}$ ).
- T3: Inoculation with *T. harzianum* ( $1 \times 10^7$  spores  $\text{mL}^{-1}$ ).

139 Acaricide compound (Vertimec) was sprayed using a Knapsack sprayer with one nozzle at the recommended rate (40 cm  $\text{fed}^{-1}$ ). All cultural practices for growing Nubian watermelon plants were performed as recommended by Egyptian Ministry of Agriculture for mineral fertilizers.

### Plant analysis

143 Samples of ten Nubian watermelon leaves were collected at random from each plot before and 3, 7, 14 and 14 days from spraying. The samples were kept in paper bags and transferred to the laboratory

for examination by stereo-microscope binocular. Motile stages and egg stage of *T. cucurbitacearum* were counted and recorded. Percentage of reduction in population was assessed according to [22].

**Table 1: Some physical, chemical and biological characteristics of the experimental soil during 2017 and 2018 growing seasons.**

Season	Clay %		Silt %		Sand %		Texture		pH	EC (dS.m <sup>-1</sup> )	O.M (%)
2017	16.88		37.12		46.00		Loam		7.12	0.210	1.86
2018	16.23		38.77		45.00		Loam		7.22	0.290	1.88
Season	Cations (mq L <sup>-1</sup> )				Anions (mq L <sup>-1</sup> )				Macro-nutrient (ppm)		
	Na <sup>+</sup>	Ca <sup>++</sup>	Mg <sup>++</sup>	K <sup>+</sup>	CO <sub>3</sub> <sup>--</sup>	HCO <sub>3</sub> <sup>-</sup>	Cl <sup>-</sup>	SO <sub>4</sub> <sup>--</sup>	N	P	K
2017	0.11	0.84	0.46	0.58	-	0.98	0.82	0.19	7.32	7.48	340
2018	0.12	0.85	0.46	0.60	-	0.95	0.86	0.22	8.90	7.86	370
Season	Total count of bacteria (CFU g <sup>-1</sup> dry weight)				Total count of fungi (CFU g <sup>-1</sup> dry weight)				Total count of actinomycetes (CFU g <sup>-1</sup> dry weight)		
2017	1.6X10 <sup>9</sup>				5.5X10 <sup>4</sup>				4.3X 10 <sup>6</sup>		
2018	2.1X10 <sup>9</sup>				6.8X10 <sup>4</sup>				4.9X 10 <sup>6</sup>		

Physical, chemical and biological analyses of soil were determined by Department of Soil Chemistry and Department of Agriculture Microbiology, Soils, Water and Environment Research Institute, ARC.

#### Data Recorded

##### Growth characters:

A random sample of three plants from every experimental unit was taken after 30 and 60 days from sowing to investigate the following growth parameters: number of leaves per plant, stem length (cm), number of branches per plant and leaves dry weight per plant.

##### Biochemical analysis

##### Chlorophyll content (SPAD):

It was estimated in the fifth leaf from growing tip after 30 and 60 days from sowing by using chlorophyll meter Model-SPAD. 502 Minolta. Co., Japan [23].

##### Plant mineral content:

After 30 and 60 days from sowing the fifth leaf from the plant growing tip were dried at 70°C then ground in a Willy mill. The dry material was wet digested with sulphuric acid-perchloric mixture as described by [24], as well as nitrogen, phosphorus and potassium percentages were determined according to the methods described by [25, 26, 27], respectively.

##### Microbial estimations:

In the rhizosphere of soil samples, total count of fungi was estimated by martin's medium according to [28] at 30 and 60 days from sowing as well as at harvest.

##### Enzyme activity:

At 30 and 60 days from sowing as well as at harvest, dehydrogenase and urease activities in the soil samples were determined as described by [29] and [30], respectively.

##### Fruit yield:

At the fruit stage, productivity was measured as the following: number of fruits per plant, number of seeds per fruit, fruit weight (g), dry weight of 100 seeds (g), fruit yield per m<sup>2</sup> (kg), seed yield per m<sup>2</sup> (g), and weight of seed yield (ton per fed.).

#### Statistical analysis:

All data obtained were subjected to analysis of variance and significant differences among means were determined at 5% level of significance according to [31].

#### Results and discussion

##### Effect of different concentrations of cultural filtrates of *T. viride* and *T. harzianum* on accumulative eggs deposited by adult females *T. cucurbitacearum* on zero time pretreated leaf discs during five days:

The data in Table (2) indicate the accumulative eggs deposited by adult females of *T. cucurbitacearum* and different concentrations of cultural filtrates of two fungal strains of *T. viride* and *T. harzianum* on zero time pretreated leaf discs during five days.

The data showed that all tested concentrations of fungi *T. viride* and *T. harzianum* were non-effective on egg deposition comparable with control treatment except concentrations 75 and 100 % throughout five days. On other hand the concentrations 75 and 100 % of cultural filtrates of fungi were the highest effect than the other concentrations from the first to third days after treatment, while other tested concentrations were gave the same effect on egg deposition by adult females of spider mite throughout 4<sup>th</sup> and 5<sup>th</sup> days comparable with control. In general the effect of different concentrations of cultural filtrates of two fungi indicated that the tested concentrations of fungi *T. viride* were non-effective on egg deposition by adult females after five days from treatment, while in *T. harzianum*, the concentration 75% was the most effect than the other concentrations. The obtained results agree with those [32] they found the total mortality percentage caused by fungus *Cladosporium cladosporioides* (freese) on bean (*Phaseolus vulgaris* L.) varied from 50.90 to 74.76 % and LT50 values ranged from 2.34 to 3.0 at 90 days . The results revealed that isolates of *C. cladosporioides* were effective against two spotted spider mite *Tetranychus urticae*.

**Table 2. Accumulative eggs deposited by adult females *T. cucurbitacearum* of different concentrations of *T. viride* and *T. harzianum* on zero time pretreated leaf discs during five days.**

Concentrations (%)	Mean No. of eggs deposited (day)				
	<i>T. viride</i>				
	1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>	5 <sup>th</sup>
100	1.75 <sup>b</sup>	3.50 <sup>c</sup>	6.50 <sup>b</sup>	32.75 <sup>a</sup>	34.00 <sup>a</sup>
75	4.5 <sup>a</sup>	12.25 <sup>b</sup>	20.50 <sup>a</sup>	29.00 <sup>a</sup>	39.00 <sup>a</sup>
50	6.75 <sup>a</sup>	14.50 <sup>ab</sup>	23.00 <sup>a</sup>	28.50 <sup>a</sup>	37.50 <sup>a</sup>
25	4.75 <sup>a</sup>	9.25 <sup>bc</sup>	10.25 <sup>b</sup>	27.00 <sup>a</sup>	37.75 <sup>a</sup>
Control	6.75 <sup>a</sup>	20.00 <sup>a</sup>	29.75 <sup>a</sup>	31.00 <sup>a</sup>	43.20 <sup>a</sup>

Concentrations (%)	Mean No. of eggs deposited (day)				
	<i>T. harzianum</i>				
	1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>	5 <sup>th</sup>
100	8.00 <sup>ab</sup>	18.00 <sup>ab</sup>	29.50 <sup>b</sup>	45.25 <sup>a</sup>	49.50 <sup>ab</sup>
75	9.00 <sup>ab</sup>	14.25 <sup>b</sup>	21.50 <sup>bc</sup>	43.50 <sup>a</sup>	41.75 <sup>b</sup>
50	12.75 <sup>a</sup>	22.00 <sup>ab</sup>	44.00 <sup>a</sup>	57.75 <sup>a</sup>	71.75 <sup>a</sup>
25	11.50 <sup>ab</sup>	27.50 <sup>a</sup>	40.50 <sup>a</sup>	51.75 <sup>a</sup>	54.33 <sup>ab</sup>

Control	6.75 <sup>b</sup>	26.00 <sup>a</sup>	19.75 <sup>c</sup>	51.00 <sup>a</sup>	53.25 <sup>ab</sup>
---------	-------------------	--------------------	--------------------	--------------------	---------------------

In 201 column means followed by a common letter are not significantly different at 1% level by DMRT.

## Effect of different concentrations of cultural filtrates of *T. viride* and *T. harzianum* on hatchability of *T. cucurbitacearum* egg at two ages:

214 Effect of four concentrations of cultural filtrates of two fungal strains of *T. viride* and *T. harzianum* on hatchability of *T. cucurbitacearum* egg at two ages (1 –and 2 day old) is shown in Table 3. It is apparent that egg hatchability decreased with increased of two egg age at treatment by concentrations 25 and 50% of both fungal , while other tested concentrations (75 and 100%) gave the same effect where the reduction in egg hatchability was 100% at two egg age . In these results may be considered the two fungal *T. viride* and *T. harzianum* as good ovicides or biocides to control of spider mite *T. cucurbitacearum*. [33] they used different concentrations of successive extracts of brown alga (*pettonia fascia*) (Muller) against adult females and egg stage of *T. urticae* (Koch) for 7 days which they found 0.1 g / ml concentration gave 100 % mortality in all extracts, the egg stage was more susceptible to diethyl ether extract and using lower concentration of different extracts, the number of eggs laid by the females was drastically decreased especially when using diethyl ether extract.

225  
Table 3. Effect of some concentration of cultural filtrates of fungal strains of *T. viride* and *T. harzianum* on hatchability and reduction (%) of *T. cucurbitacearum* egg at two ages

Concentrations (%)	Egg age (day)							
	<i>T. viride</i>				<i>T. harzianum</i>			
	1 – day old (%)		2 – day old (%)		1 – day old (%)		2 – day old (%)	
	H.	R.	H.	R.	H.	R.	H.	R.
100	0.0	100	0.0	100	0.0	100	0.0	100
75	0.0	100	0.0	100	0.0	100	0.0	100
50	6.98	93.03	0.0	100	8.70	91.31	0.0	100
25	8.70	91.31	0.0	100	9.75	90.25	0.0	100
Control	100	0.0	100	0.0	100	0.0	100	0.0

H. = Hatchability (%); R. = Reduction (%).

## Effect of vertemic on population density of the spider mite on Nubian watermelon leaves as affected by inoculation with fungal strains of *T. viride* and *T. harzianum* in the field experiment:

232 Intensity of infestation with the spider mite *T. cucurbitacearum* presented in Table (4) as counted through (30) days beginning from the 30 up to 60 days after sowing this period which are expected to intensity infestation with the mites .

235 The data indicated that the highest reduction of motile stages and egg stage was recorded after three days from application of *T. harzianum* after that all treatments recorded reduction 100 % of both motile stages and egg stage after 7 and 14 days from application. The average percent reduction ranged between (77.88 – 93.94 %) of motile stage while egg stage recorded 100% reduction of all treatments. The obtained results agree with those [34] found that the Biafly exhibited mean reduction

76.44% in motile stages of spider mite *T. urticae* and 81.55 % mean reduction in egg stage on soybean plants in the field conditions .

**Table 4. Population density of the spider mite *T. cucurbitacearum* on Nubian watermelon leaves as affected by inoculation with *T. viride* and *T. harzianum* in the field experiment during 2018 season**

Treatment	Reduction at indicated days (%)								Mean reduction (%)	
	before treatment		3		7		14			
	Mite	Eggs	Mite	Eggs	Mite	Eggs	Mite	Eggs	Mite	Eggs
<i>T. viride</i>	44.67	37.67	33.64	100	100	100	100	100	77.88	100
<i>T. harzianum</i>	55.33	46.0	81.82	100	100	100	100	100	93.94	100
Control	50.67	57.0	45.33*	52.0*	25.33*	30.33*	13.67*	23.33*	28.11*	35.22*

\* Mean number of *T. cucurbitacearum* motile and egg stages in control through experiment period.

#### **Vegetative growth:**

Growth parameters included number of leaves, Leaf dry weight, stem length and numbers of branches were estimated at 30 and 60 days from sowing during 2017 and 2018 seasons.

Data of Table 5 indicate that in most cases, plants inoculated with *T. viride* and *T. harzianum* had the highest values of leaves number / plant during both growth periods (30 and 60 days after sowing), compared to un-inoculated plants (control). The maximum values were obtained from plants inoculated with *T. harzianum* which recorded 27.66 and 30.33 for the first season and 206.66 and 214.06 for the second season. The differences were significant in both seasons.

Concerning leaf dry weight and stem length, data of Table 5 showed that increasing effect on the leaf dry weight and stem length during different growth periods of both seasons. The increased rate was obtained from plants inoculated with *T. harzianum* which recorded 31% at 30 days and 33% at 60 days from sowing compared to control treatment during 2017 season. The same trend was observed at 2018 season. For number of branches / plant, all inoculation treatments with studied fungi (*T. viride* and *T. harzianum*) showed significantly higher number of branches / plant during both seasons and different growth periods reached to 10.66 and 12.33 at 60 days from sowing for inoculation treatment with *T. harzianum* during 2017 and 2018 seasons, respectively.

Effectiveness of inoculation with *Trichoderma* treatments in improvement of growth were evident from the initial stages itself wherein number of leaves, Leaf dry weight, stem length and number of branches were improved over the control during the two growing seasons. These results suggest that *Trichoderma* strains have a strong capacity to mobilize and increase plants nutrient uptake, thus making it more efficient and competitive than many other soil microbes to promote growth and development [35]. A number of mechanisms for plant growth promotion by *Trichoderma* which lead to the increase in nutrient availability and its uptake, resulting in the efficient nutrient absorption and thereby promoting plant growth have been reported by [36]. These traits were considered as the basis

for 173 *Trichoderma* exerted beneficial effects on plant growth and development. However, now it is  
 174 becoming increasingly clear that certain strains have substantial direct influence on plant growth and  
 175 development and on crop productivity with multiple mode of action [37]. Similar improvement of  
 176 different vegetative growth by strains of *T. harzianum* has been reported earlier in sunflower [38], bean  
 177 [39], cotton [40], tomato [41] and chilli [42] sunflower [43].

278

**Table 5. Effect of inoculation with PGPF on number of leaves, Leaf dry weight (g), stem length (cm) and numbers of branches of Nubian watermelon plants at 30 and 60 days from sowing during 2017 and 2018 summer seasons**

Treatments	No. leaves plant <sup>-1</sup>		Leaf dry weight (g)		Stem length (cm)		No. branches	
	2017							
	30	60	30	60	30	60	30	60
T1	17.33 c	151.33 c	0.044 b	0.093 c	42.33 c	196.00 b	3.33 b	6.33 c
T2	24.00 b	167.66 b	0.054 a	0.109 b	55.00 b	231.66 a	4.33 ab	8.33 b
T3	27.66 a	206.66 a	0.058 a	0.124 a	64.00 a	216.00 b	5.33 a	10.66 a
LSD 0.05	1.48	13.10	0.004	0.006	4.66	9.32	1.15	1.63
Treatments	No. leaves plant <sup>-1</sup>		Leaf dry weight (g)		Stem length (cm)		No. branches	
	2018							
	30	60	30	60	30	60	30	60
T1	10.33 c	153.66 c	0.047 b	0.100 c	45.33 c	201.33 c	3.66 b	6.66 c
T2	25.66 b	173.66 b	0.057 a	0.110 b	58.33 b	219.00 b	4.66 ab	8.33 b
T3	30.33 a	214.00 a	0.060 a	0.130 a	65.66 a	236.00 a	5.33 a	12.33 a
LSD 0.05	1.63	8.04	0.003	0.005	2.57	7.14	1.15	1.15

In a column means followed by a common letter are not significantly different at 5% level by DMRT. T<sub>1</sub>: Control; T<sub>2</sub>: inoculation with *T. viride* and T<sub>3</sub>: inoculation with *T. harzianum*.

282

### Biochemical analysis of leaves:

284 Biochemical analysis of leaves was determined as the total chlorophyll content (SPAD) and the  
 285 content of nitrogen (N), phosphorus (P) and potassium (K) as percentage of leaves dry weight.

286 Data in Figure 1 cleared that Nubian watermelon plants inoculated with *T. harzianum* had the  
 287 highest chlorophyll content followed by that inoculated with *T. viride* in both seasons, while the lowest  
 288 values were obtained from non-inoculated plants (control). The application of the different fungi strains  
 289 revealed that there is an increase in nitrogen, phosphorus and potassium percentage. The highest N  
 290 content was found in treatments 3, where it recorded 1.75 and 2.92% followed by 1.68 and 2.88% for  
 291 treatment 2 at 30 and 60 days from sowing compared to control treatment during 2017 season,  
 292 respectively. The same trend was observed in 2018 season (Fig. 1). In case of P content (%), the results  
 293 showed that the influence of the studied bio-inoculants on the leaves P content had a similar trend. T3  
 294 treatment gave, 0.22 and 0.23% with regard to control (T1) which exhibited 0.16 and 0.17% at 60 days  
 295 of 2017 and 2018 seasons, respectively. For K content (%), the results showed an increase of K content  
 296 of leaves recorded 1.39 and 2.48% for T3 (inoculated with *T. harzianum*), followed 1.25 and 2.39% for  
 297 T2 (inoculated with *T. viride*), compared to 1.11 and 2.18% for control treatment (T1) at 30 and 60  
 298 days during the first growing season. The differences were significant in both seasons (Fig. 1).



299 These results may be refer to that inoculation treatments influences absorption and translocation  
of magnesium [44], which plants contain more Mg in their leaves than the non-inoculated ones and that  
may interpret their higher chlorophyll content. Also, the increase in the chlorophyll content attributed  
can be ascribed to the presence of rhizomicrobes in the rhizosphere influencing the crop roots to secrete  
growth promoting substances.

304 The increase in N% in Nubian watermelon plants at 30 and 60 days from sowing lead to the  
increase in nutrient availability and its uptake and these increased due to single or combined  
inoculation of microorganisms which has been documented on different crops such as sunflower [43]  
soybean [45] common bean [46] and cowpea [47]. For phosphorus, mineralization process through  
microorganisms as well as plants have many potential mechanisms to increase P uptake from soil  
including regulation of phosphate membrane transport systems, the increased growth of root hairs, the  
release of phosphatases, changes in root architecture and the release of organic acids which all of them  
due to the availability of nutrient phosphate in the soil that eventually will be uptake by plants.  
Similarly, fungal strains (*T. viride* and *T. harzianum*) are known to produce some organic products  
such as citric acid, oxalic acid that are mainly known to decompose or solubilize natural silicates and  
help removal of metal ions from the rocks and soils and this process was more effective in the  
alkaline soils and lead to increase K uptake to plant [48, 49].

316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337




Fig. 1. Effect of inoculation with PGPF on chlorophyll content and percentages of NPK of Nubian watermelon plants at 30 and 60 days from sowing during 2017 and 2018 summer seasons

## Enzyme activity:

Data of Table 6 revealed an increase in dehydrogenase and urease activities with the application of different inoculation treatments. The dehydrogenase activity was noted to increase then decrease in the harvest stage. In general, the treatment T2 (inoculated with *T. viride*) recorded the highest values at all growth stages, which recorded 155.77, 257.29 and 114.62 mg TPF g<sup>-1</sup> soil day<sup>-1</sup> at 30, 60 and harvest during 2017 growing seasons, respectively. The same trend was observed in the second growing season (2018).

On the other hand, urease activity was shown to rise in 60 days from sowing due to inoculation treatments compared to the respective control. Also, it is clearly showed that urease activity levels decreased with the increase in plant age (harvest). T2 treatment showed significant maximum urease activity (201.03 and 211.95 mg NH<sub>4</sub><sup>+</sup>- N g<sup>-1</sup> soil d<sup>-1</sup>) followed by T3 (188.59 and 201.22 mg NH<sub>4</sub><sup>+</sup>- N g<sup>-1</sup> soil d<sup>-1</sup>) compared to control (156.27 and 164.18 mg NH<sub>4</sub><sup>+</sup>- N g<sup>-1</sup> soil d<sup>-1</sup>) at 60 days from sowing during 2017 and 2018 growing seasons, respectively .

It is well known that enzymes play a key role in the transformation, recycling and availability of plant nutrients in soil. They are likely to be influenced by fertilizers and manures. [50] Showed that the increase in dehydrogenase activity was mainly due to the higher microbial population and the earlier studies revealed that the enzyme activities are often used as indices of microbial growth rather than the microbial number, which further may reflect the microbial respiration and the potential capacity of soil to perform biological transformations of importance to soil fertility. Also, the variation in the urease was little influenced by different inoculation treatments. More than the microbial population, the enzyme activities are regulated by the soil characters like organic carbon, pH and nutrient status [51] The observations are in accordance with [45, 46, 49, 52].

**Table 16. Effect of inoculation with PGPF on dehydrogenase and urease activities at different growth stages of nubian watermelon plants during 2017 and 2018 summer seasons**

Treatments	Dehydrogenase activity (mg TPF g <sup>-1</sup> soil d <sup>-1</sup> )					
	2017			2018		
	30	60	harvest	30	60	harvest
T1	106.86 c	140.45 c	92.37 c	113.74 c	129.10 c	86.22 c
T2	155.77 a	257.29 a	114.62 a	159.84 a	248.46 a	121.15 a
T3	127.53 b	234.36 b	106.10 b	144.74 b	237.47 b	106.21 b
LSD 0.05	7.55	9.32	6.44	7.29	5.18	3.95
Treatments	Urease activity (mg NH <sub>4</sub> <sup>+</sup> - N g <sup>-1</sup> soil d <sup>-1</sup> )					
	2017			2018		
	30	60	harvest	30	60	harvest
T1	91.85 c	156.27 c	105.81 c	101.47 c	164.18 c	107.34 c
T2	113.79 a	201.03 a	115.24 b	122.02 a	211.95 a	137.15 a
T3	108.18 c	188.59 b	124.84 a	111.39 b	201.22 b	132.01 b
LSD 0.05	2.79	3.90	3.49	3.64	4.48	3.77

In a column means followed by a common letter are not significantly different at 5% level by DMRT. T<sub>1</sub>: Control; T<sub>2</sub>: inoculation with *T. viride* and T<sub>3</sub>: inoculation with *T. harzianum*.

## Fungal estimations:

Response of Nubian watermelon plants to the inoculation with *T. viride* and *T. harzianum* strains and their activities in the rhizosphere of soil samples with different times after application during 2017 and 2018 growing seasons are presented in Figure 2. The differences were significant in both seasons.

Concerning untreated control, results showed few increasing in counts of fungi with different days after treatment. However, the treated soil showed an increasing trend of log number from 30 day (6.49 and 6.26), to 60 day (7.36 and 7.09), after treatment then gradually decreases and found to be minimum on harvest (5.71 and 5.39), for inoculation with *T. viride* and *T. harzianum* treatments at 2017 season, respectively. Similar trend was also exhibited in treated soil at 2018 season.

The application of inoculation with plant growth promoting fungi *T. viride* and *T. harzianum* may lead to significant changes in microbial populations and activities influencing microbial ecological balance affecting soil fertility. Also, the gradual increase in fungi counts may be attributed to the ability to temporarily mineralize and use soil organic carbon as energy source and this greater count of fungi in the rhizosphere at 30 and 60 days from sowing leading to greater release of plant nutrients in soil for enhancement the growth and yield of crops [53, 54, 55].




Fig. 2 Effect of inoculation with PGPF on total count of fungi (CFU log<sub>10</sub> g<sup>-1</sup>) at different growth stages of nubian watermelon plants during 2017 and 2018 summer seasons

### Fruit yield:

392 Fruit yield was determined as number of fruits per plant, number of seeds per fruit, , fruit  
weight (g), dry weight of 100 seeds (g), fruit yield per m<sup>2</sup> (kg), seed yield per m<sup>2</sup> (g), and weight of  
seed yield (ton fed.<sup>-1</sup>).

395 Data in Table 7 indicated that plants treated with *T. viride* (T2) had significantly the highest  
number of fruits per plant, number of seeds per fruit, fruit weight (g) and dry weight of 100 seeds (g)  
which recorded 1.92, 273.07, 1126 g and 16.29 g as compared to untreated control treatment, which  
attained 1.21, 185.08, 526.66 g and 14.41 g at the first growing season (2017), respectively. The same  
trend was observed in the second growing season (2018).

400 Data presented in Table 8 cleared that the highest fruit yield per m<sup>2</sup> was found in treatments 2,  
where it recorded 7.33 and 7.01 followed by 4.71 and 5.22 for treatment 3 compared to control  
treatment during 2017 and 2018 growing seasons, respectively. In case of seed yield per m<sup>2</sup> (g), the  
results showed that the influence of the studied bio-inoculants had a similar trend. T2 treatment gave,  
2224.00 and 226.00 g with regard to control (T1) which exhibited 107.33 and 121.66 at 2017 and 2018  
seasons, respectively (Table 8). For weight of seed yield (ton fed.<sup>-1</sup>) the results showed an increase of  
seed yield of Nubian watermelon plants recorded 0.93 and 0.95 ton fed.<sup>-1</sup> for T2 (inoculated with *T.*  
*viride*), followed 0.76 and 0.76 ton fed.<sup>-1</sup> for T3 (inoculated with *T. harzianum*), compared to 0.45 and  
0.51 ton fed.<sup>-1</sup> for control treatment (T1) at the first and second growing seasons, respectively. The  
differences were significant in both seasons (Table 8).

410 It is clear from the above mentioned data that plants inoculated with PGPF had higher values  
for most fruit yield parameters compared to control [56, 57]. Also, the root system of Nubian  
watermelon plants mostly had very strong root system [58] it is often capable of absorbing water and  
nutrients more efficiently than non-inoculated plants and may serve as a good supplier of endogenous  
plant hormones [59, 60]. These results are in harmony with those obtained by several researches on  
cucumber [61, 62], watermelon [63, 64], Nubian watermelon [65].

416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429

**Table 7. Effect of inoculation with PGPF on number of fruits / plant, number of seeds / fruit, fruit weight (g) and dry weight of 100 seeds (g) of nubian watermelon plants during 2017 and 2018 summer seasons**

Treatments	number of fruits / plant	number of seeds / fruit	fruit weight (g)	dry weight of 100 seeds (g)
<b>2017</b>				
<b>T1</b>	1.21 c	185.08 c	526.66 c	14.41 b
<b>T2</b>	1.92 a	273.07 a	1126 a	16.29 a
<b>T3</b>	1.54 b	233.00 b	939.66 b	14.29 b
<b>LSD 0.05</b>	0.06	8.19	55.30	0.89
<b>2018</b>				
<b>T1</b>	1.24 c	193.61 c	764.66 c	13.16 b
<b>T2</b>	1.99 a	298.16 a	1132 a	14.11 a
<b>T3</b>	1.47 b	270.83 b	1009.33 c	13.06 b
<b>LSD 0.05</b>	0.09	13.38	33.36	0.21

In a column means followed by a common letter are not significantly different at 5% level by DMRT. T<sub>1</sub>: Control; T<sub>2</sub>: inoculation with *T. viride* and T<sub>3</sub>: inoculation with *T. harzianum*.

**Table 8. Effect of inoculation with PGPF on fruit yield (kg/m<sup>2</sup>), seed yield (g/m<sup>2</sup>), and weight of seed yield (ton fed.<sup>-1</sup>) of nubian watermelon plants during 2017 and 2018 summer seasons**

Treatments	fruit yield (kg/m <sup>2</sup> )	seed yield (g/m <sup>2</sup> )	Weight of seed yield (ton fed. <sup>-1</sup> )
<b>2017</b>			
<b>T1</b>	2.22 c	107.33 c	0.45 c
<b>T2</b>	7.33 a	222.00 a	0.93 a
<b>T3</b>	4.71 b	181.00 b	0.76 b
<b>LSD 0.05</b>	0.48	8.42	0.03
<b>2018</b>			
<b>T1</b>	3.56 c	121.66 c	0.51 c
<b>T2</b>	7.01 a	226.66 a	0.95 a
<b>T3</b>	5.22 b	183.00 b	0.76 b
<b>LSD 0.05</b>	0.15	13.18	0.03

In a column means followed by a common letter are not significantly different at 5% level by DMRT. T<sub>1</sub>: Control; T<sub>2</sub>: inoculation with *T. viride* and T<sub>3</sub>: inoculation with *T. harzianum*.

## CONCLUSION

The results of the present investigation confirmed that application of inoculation with PGPF (*T. viride* and *T. harzianum*) at the time of planting could be recommended for controlling the two-spotted spider mite, *Tetranychus cucurbitacearum* of nubian watermelon plants as well as increased the activities of most soil enzymes, especially dehydrogenase and urease and enhancement the vegetative growth and seed yield.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Muthu KB, Priji P, Unni KN, Sajiths S, Anithak N, Rosana B, Benkamins MB. New *Bacillus thuringiensis* strain isolated from the gut of Malabari goat is effective against *Tetranychus macfarlanei*. Journal of applied entomology, 2015, 140 (3) : 187 – 198 . DoI : <https://doi.org/10.1111/jen.12235>.
2. Park YL, Lee JH. Leaf cell and tissue damage of cucumber caused by two spotted spider mite (Acari: Tetranychidae) . Journal of Economic Entomology, 2002, 95 (5) : 952 – 957 .
3. Velleu T, Vontas J, Tsagkarakou A, Dermauw W, Tirry L. Acaricide resistance mechanisms in the two spotted spider mite *Tetranychus urticae* and other important Acari: a review. Insect Biochemistry and Molecular Biology, 2010, 40 (8): 563–572. DOI:<https://doi.org/10.1016/j.ibmb.2010.05.008>
4. Fiedler Z. Interaction between beneficial organisms in control of spider mite *Tetranychus urticae* (Koch.). Journal of Plant Protection Research. 2012, 52 (2): 226–229. DOI: <https://doi.org/10.2478/v10045-012-0035-2>
5. Arznlou M, Mousavi S, Bakhshi M, Khakvar R, Bandehagh A. Inhibitory effects of antagonistic bacteria inhabiting the rhizosphere of the sugarbeet plants , on *cercospora beticola* sacc, the causal agent of cercospora leaf spot disease on sugarbeet . Journal of plant protection Research, 2016, 56 (1) : 6 – 14 . DoI : <https://doi.org/10.1515/jppr-2016-0002>
6. Hamza A, Mohamed A, Derbalah A. Unconventional alternatives for control of tomato root caused by *Rhizoctonia solani* under greenhouse conditions . Journal of plant protection research, 2016, 56 (3): 298 – 305. DoI :<https://doi.org/10.1515/jppr-2016-0046> .
7. Chandanie WA, Kubota M, Hyakamachi M. Interaction between arbuscular mycorrhizal fungus *Glomus mosseae* and plant growth promoting fungus *Phoma* sp. on their root colonization and growth promotion of cucumber (*Cucumis sativus*). Mycoscience. 2006, 46:201-204.
8. Chapplick GP. Recovery from drought stress in *Lolium perenne* (Poaceae): are fungal endophytes detrimental? American J. of Bot. 2004, 91:1960-1968.
9. Harrison MJ. Signalling in the arbuscular mycorrhizal symbiosis. Annu. Rev. Microbiol. 2005, 59:19-48042.
10. Dashmukh S, Ckellhoven RH, Schafer P, Imani J, Sharma M, Weiss M, Waller F, Kogel KH. The root endophytic fungus *Piriformospora indica* requires host cell death for proliferation during mutualistic symbiosis with barley. PNAS. 2006, 103(49):18450-18457.
11. Murali M, Amruthesh KN, Sudisha J, Niranjana SR, Shetty HS. Screening for plant growth promoting fungi and their ability for growth promotion and induction of resistance in pearl millet against downy mildew disease. J. of Phytopathology, 2012, 4(5): 30-36
12. Hek AE, Leger RJ. Interactions between fungal pathogens and insect hosts. Annu. Rev. Entomol. 1994, 39:293–322. doi:10.1146/annurev.en.39.010194.001453
13. Menniche L. An ancient Egyptian herb. Third university of Texas press printing . Great Britain, 1989, pp. 92.
14. FAO. The state of food insecurity in the world 2008.
15. Mariod AA, Ahmed YM, Matthaus B, Khaled G, Siddig A, Gaber AM, Abdelwahab SI. A comparative study of the properties of six Sudanese cucurbit seeds and seed oils. An oil chem. Soc. 86 (12): 1181 – 1181.
16. Ibrahim MO, Ahmed FA, Sulieman YR. The Nutritive value of Gurum seed cakes (*Citrullus lanatus* var. *colocynthis*) as feed for ruminants . Sudan Ancient production. 2002, 15 : 71 – 78 .

17. 478 Syada AK, Elhussin SA. Physical and chemicals .characteristics of *citrulluslanatus* var.  
499 *colocynthoide* seed oil . physical. Sci. 2008, 19 (2) : 69 – 75 .
18. 500 Osunaka A, Hyakumachi M, Takenaka S. (2011). Plant Growth–Promoting Fungus, *Trichoderma*  
501 *koningi* Suppresses Isoflavonoid Phytoalexin Vestitol Production for Colonization on/in the  
502 Roots of Lotus japonicas. Microbes Environ. 2011, 26, 2, 128–134,  
503 doi:10.1264/jsme2.ME10176
19. 504 Pappas ML, Liapoura M, Papantoniou D, Avramidou M, Kavroulakis N, Weinhold A, Broufas GD,  
505 Papadopoulou KK. The Beneficial Endophytic Fungus *Fusarium solani* Strain K Alters Tomato  
506 Responses Against Spider Mites to the Benefit of the Plant. Front. Plant Sci. 2018, 9:1603. doi:  
507 10.3389/fpls.2018.01603  
508
20. 509 Bogdady MME. Integrated Postharvest Diseases Management of Certain Pome Fruits. Ph.D.  
510 thesis Fac. Agric. 1993, Al-Azhar Univ., 55 pp.
21. 511 Egler EH. Leaf dis technique for laboratory tests of acaricidesj. Econ. Entomod. 1947, 40: 441 –  
512 442 .
22. 513 Henderson CF, Tilton EW. Test with acaricides against the brown wheat mite .J .Econ. Entomol.  
514 1955, 48 : 157 – 161 .
23. 515 Silva UL. A rapid and non-destructive method to determine chlorophyll in intact leaves. Hort. Sci.  
516 1986, 21: 1449.
24. 517 Allen SG, Girmshaw HM, Parkinson JA, Quarmby C. Chemical analysis of ecological materials.  
518 1974, Black Well Sci. Publ. Oxford, London. 565pp
25. 519 Pregl F. Quantitative organic microanalysis 4<sup>th</sup> Edit. J. A. Churchill Ltd. London; 1945.
26. 520 ough E, Mager AH. Improvement in deiness colorimetric method for phosphorus and arsenic.  
521 Ind. En. Chemical Anal. Ed. 1939;1:136-139.
27. 522 Brown JD, Lilliland O. Rapid determination of potassium and sodium in plant material and soils  
523 extracts by flam-photometry. Proc. Amer. Soc. Hort. Sci. 1946;48:341- 346.
28. 524 Allen ON. Experiments in Soil Bacteriology. 1959, Burgess pub. Co., Ninn, Minnesota
29. 525 Sida LE, Klein DA and Snatoro T. Soil dehydrogenase activity. Soil Sci. 1964, 98, 371-376.
30. 526 nicholy SK, Rice EL. Soil enzymes in relation to old succession : Amylase, invertase, cellulose,  
527 dehydrogenase and urease. Soil Sci. Soc. American Proc.,1973, 37, 47-50.
31. 528 Duncan BD. Multiple range and multiple test. Biometrics. 1955, (11) : 1 – 42 .
32. 529 Ben C, Hayat R. Preliminary evaluation of *Cladosporium cladosporioides* (fresen) de vries in  
530 laboratory conditions, as a potential candidate for biocontrol of Tetranychus urticae (Koch)  
531 department of plant . World J. Microbiol. Biotechnal. 2009, 25: 489 – 492.
33. 532 er SAA, Dimetry NZ, Mahran GH, Saleh MM, El-sherei MM, Motawe HM, M . H Grace MH.  
533 A caricid al and Biological properties of Brown marine Alga Petalomia fascia (Muller Kuntze)  
534 against the two – spotted spider mite *Tetranychus urticae* (koch) – Bull .zool , soc . 1994,  
535 Egypt, 42 (91 – 101).
34. 536 Agouz R. Integrated spider mite management. 2003, Ph.D. Agric. (pesticides). Tanta Univ. pp.  
537 253 .
35. 538 itez T, Rincón AMM, Limón C, Codón AC. Biocontrol mechanisms of *Trichoderma* strains.  
539 Int. Microbiol. 2004, 7:249-260
36. 540 rman GE, Howell CR, Vitrebo A, Chet I, Lorito M. *Trichoderma* species opportunistic, arivalent  
541 plant symbionts. Nat. Rev. Microbiol. 2004, 2:43-56



37. ~~542~~ Herman GE. Overview of mechanisms and uses of *Trichoderma* spp. Phytopathol. 2006, 96:190-543 194
38. ~~544~~ Dawar S, Hayat S, Anis M, Zaki MJ. Effect of seed coating material in the efficacy of microbial 545 antagonistic for the control of root rot fungi on okra and sunflower. Pak. J. Bot. 2008, 546 40(3):1269-1278
39. ~~547~~ Oyos-Carvajal L, Ordua S, Bissett J. Growth stimulation in bean (*Phaseolus vulgaris* L.) by 548 *Trichoderma*. Biol. Cont. 2009, 51:409-416
40. ~~549~~ Annmugaiah V, Balasubramanian N, Gomathinayagam S, Monoharan PT, Rajendran A. Effect of 550 single application of *Trichoderma viride* and *Pseudomonas fluorescences* on growth promotion in 551 cotton plants. Afr J Agric Res., 2009, 4(11):1220-1225
41. ~~552~~ Armi R, Hajieghrari B, Gigloul A. Effect of *Trichoderma* isolates on tomato seedling growth 553 response and nutrient uptake. Afr. J. Biotech..2011, 10:5850-5855
42. ~~554~~ Alam MS, Rahman MA, Bulbul SH, Alam MF. Effect of *Trichoderma* on seed germination and 555 seedling parameters in chilli. Int. J. Expt. Agric., 2011, 2(1):21-26
43. ~~556~~ Nagaraju A, Sudisha J, Mahadeva MS, Shin-ichi I. Seed priming with *Trichoderma harzianum* 557 isolates enhances plant growth and induces resistance against *Plasmopara halstedii*, an incitant 558 of sunflower downy mildew disease. Australasian Plant Pathol., 41:609–620. DOI 559 10.1007/s13313-012-0165-z
44. ~~560~~ Iz JM, Belakbir A, Lopez-Cantarero A, Romero L. Leaf macronutrient content and yield in 561 grafted melon plants: a model to evaluate the influence of rootstock genotype. Sci. Hort. 1997, 562 71:227-234.
45. ~~563~~ Omara A, Hauka F, Afify A, Nour El-Din M, Kassem M. The Role of Some PGPR Strains to 564 Biocontrol Rhizoctonia Solani in Soybean and Enhancement The Growth Dynamics and Seed 565 Yield. Environ. Biodiv. & Soil Security, 2017, 1, 47 – 59.
46. ~~566~~ Nahrawy S, Omara A. Effectiveness of Co-inoculation with *Pseudomonas koreensis* and 567 Rhizobia on Growth, Nodulation and Yield of Common Bean (*Phaseolus vulgaris* L.). 568 Microbiology Research Journal International, 2017, 21(6): 1-15, 2017; Article no.MRJI.37422
47. ~~569~~ Omara A, El-Gaafarey T. Alleviation of Salinity Stress Effects in Forage Cowpea (*Vigna* 570 *unguiculata* L.) by *Bradyrhizobium* sp. Inoculation. Microbiology Research Journal 571 International, 2018, 23(3): 1-16, 2018; Article no.MRJI.40727
48. ~~572~~ Omara A, Azza AG, El-Akhdar IA. Isolation and Identification of Zinc Dissolving Bacteria and 573 Their Potential on Growth of Zea mays. Egypt. J. Microbiol. 2016, 51, 29 – 43.
49. ~~574~~ Nour El-Din M, Omara A, Elbagory MH. Effect of Bulking Agent Levels (Rice Straw) on Compost 575 Quality and Net Return. International Journal of Plant & Soil Science, 2016, 19(3): 1-12; 576 Article no.IJPSS.36518
50. ~~577~~ Indrayan K, Adhya JK, Sethunathan N. Dehydrogenase and invertase activities of flooded soils. 578 Soil Biol. Biochem. 1980 12, 271-273.
51. ~~579~~ Nagaraja MS, Ramakrishna VR, Siddaramappa R. Effect of atrazine urea-N-mineralization and 580 activity of some soil enzymes. Indian J. Agron., 1998, 46: 189-192.
52. ~~581~~ Akhdar I, Omara A, Abdel-Rahman MA. Intergradation of Different Fertilizers for Sustainable 582 Agriculture Enhanced Growth and Yield of Wheat (*Triticum aestivum* L.). Env. Biodiv. Soil 583 Security. 2018, 2, 11- 24.
53. ~~584~~ Das AC, Debnath A. Effect of systemic herbicides on N<sub>2</sub>-fixing and phosphate solubilizing 585 microorganisms in relation to availability of nitrogen and phosphorus in paddy soils of West 586 Bengal. Chemosphere, 2006, 65: 1082–1086. doi: 10.1016/j.chemosphere.2006.02.063

54. 587 Chen Z, Chen H, Zou Y, Qiu J, Wen Y, Xu D. Are nutrient stresses associated with  
588 enantioselectivity of the chiral herbicide imazethapyr in *Arabidopsis thaliana*? J. Agric. Food  
589 Chem. 2015, 63, 10209–10217. doi: 10.1021/acs.jafc.5b04495
55. 590 Anara A, El-Ghandor A. Effect of Thiobencarb and Penoxsulam Herbicides on Soil Microbial  
591 Population and Weed Control in Transplanted Rice. Global Advanced Research Journal of  
592 Microbiology, 2018, 7(5), 084-094.
56. 593 Hebarria PH, Castro R, Vivaracho SM, Vallejo AD. Influence of rootstock and soil treatments on  
594 the yield and quality of greenhouse-grown cucumber in Spain. Acta. Hort. 2002, 633: 403-408.
57. 595 Nemr M. Effect of mulch types on soil environmental conditions and their effect on the growth  
596 and yield of cucumber plants. App. Sci. Res. 2006, 22: 67-73.
58. 597 H YC, Lee JM, Lee SG. The current situation and prospect of cucurbits grafting in Korea.  
598 Türkiye Ulusal V. Bah. Bit. Kong. 2007, 37-39. Erzurum, Turkey.
59. 599 ni S, Gruidina R, Cadinu M, Madeddu B, Carlett MG. The influence of four rootstocks on some  
600 melon hybrids and a cultivar in greenhouse. Acta. Hort., 1990, 287: 127-134.
60. 601 JM (1994). Cultivation vegetable I. Current status, grafting methods, and benefits. Hort., 1994,  
602 29(4): 235-239.
61. 603 eel SMAM. Effect of some Cucurbits root stocks on growth, yield and resistance to some soil  
604 pests of cucumber under plastic houses. 1999, Ph. D. Thesis. Zagazig Univ. Moshtohor. Egypt
62. 605 d Alla AM. Effect of soil solarization, fertilizer sort and grafting on growth and productivity of  
606 cucumber crop (*Cucumis sativus*, L.) 2002, Ph. D. Thesis. Fac. Agric., Kafr El-Sheikh. Tanta  
607 Univ., Egypt.
63. 608 med ESM. Effect of grafting on growth and yield of watermelon plants under low plastic tunnels  
609 in Baltiem district. 2005, Ph. D. Thesis. Fac. Agric., Kafr El-Sheikh. Tanta Univ., Egypt.
64. 610 isir H, Sari N. Effect of different rootstock on plant growth, yield and quality of watermelon.  
611 Aust. Exper. Agri. 2003, 43(10): 1269 - 1274.
65. 612 Tanbashawy H. Increase the Productivity of Nubian Watermelon Plant by Grafting Technique.  
613 2012, M.sc., Thesis, Faculty of Science, Tanta University

614

615

616

617

618