

2  
3 **Planting Spacing of Cultivated Soybean**  
4 **Intercropped With Cover Plants**  
5  
6  
7  
8  
9

10 **ABSTRACT**  
11

**Aims:** The objective of this work was to evaluate the influence of planting spacing in soybean intercropped with covering species in the Roraima savanna.

**Study design:** The experimental design was a randomized complete block design with four replications.

**Place and Duration of Study:** The experiments were conducted at Embrapa Roraima, in Campo Experimental Água Boa, municipality of Boa Vista - Roraima state, in 2015 and 2016.

**Methodology:** Plots consisted in the spacing (0.45, 0.55 and 0.65 m) and the subplots were constituted by the cover plant species *Urocloa brizantha*, *Urocloa ruziensi*, *Panicum maximum* and the treatment without intercropping. The used soybean cultivar was BRS Tracajá in two crops. The following variables had evaluated: plant height, number of grains per pod, number of pods per plant, 100-grain weight, plant dry matter, insertion of the first pod, grain yield, dry matter of the covering species e of spontaneous vegetation.

**Results:** Cover plants affected the plant height, number of pods per plant, insertion of the first pod, dry matter of cover species and yield of grains in soybean. The spacing did not influence the growth and production of the soybean crop, except positively in the number of pods per plant with the increased of spacing. The interaction of cover plants and spacing affected the weight of 100 grains, the insertion of the first pod and the dry mass of the cover species. Number of grains per pod and the dry mass of the soybean plants were not affected by the cover plants and by the spacing.

**Conclusion:** The *U. brizantha* species provides the highest production of dry matter intercropped with soybean, however, the yield of the crop decreases. The *U. ruziensi* species is the most suitable for the cultivation intercropped with the crop. The used spacings do not influence the productivity.

12 *Keywords: competition; intercropped cultivation; forages; Glycine max; grain production.*  
13  
14

15 **1. INTRODUCTION**  
16

17 Brazil is the second largest soybean (*Glycine max*) producer in the world, following only the  
18 United States. In the 2015/2016 growing season, this crop occupied an area of 33.17 million  
19 hectares, totaling a production of 95.63 million tons; the average yield of soybean in Brazil  
20 was 2,882 kg ha<sup>-1</sup>. In the same growing season, Roraima occupied an area of 24 thousand  
21 hectares, with a production of 79.2 thousand tons and totaling a productivity of 3,300 kg ha<sup>-1</sup>  
22 [1].  
23

24 Integrated systems can contribute to the production of soybean and to sustainability in  
25 different regions of Brazil, becoming an option to increase and diversify the income of  
26 producers, as well as for future improvements of no-till systems [2, 3].

27

28 The intercropping with forage species is a long-term method and consists of the cultivation of  
29 two or more crops in the same place with the objective of maximizing the productivity and  
30 quality of the obtained production [4, 5].

31

32 Intercropped cultivations with forage species from the genus *Urochloa* have been proving to  
33 be profitable and compatible, aiming at both straw and grain production. However, one of the  
34 limitations faced by producers for the adoption of the no-tillage system for soybean in the  
35 Cerrado of Roraima is related to the difficulty of establishing these plant species after  
36 harvesting commercial crops, due to the marked water deficit occurring from October to  
37 March.

38

39 In order to maximize the yield of a crop, the use of spacing and the used cultivar contribute  
40 most of the time to soybean yield. In this context, it is important to emphasize the spacing  
41 between rows to be used while sowing. According to Tourino (2002) [6], Procópio et al.  
42 (2014) [7], and Balena et al. (2016) [8], spacing can be managed in order to define a more  
43 suitable arrangement to obtain higher yields and the adaptation to harvesting fabaceae by  
44 machines. Also, by defining an adequate spacing, it is possible to provide good productivity  
45 and weed management, thus contributing to soil sustainability.

46

47 The objective of this work was to evaluate the influence of spacing and intercropping with  
48 cover crop species on the performance of soybean in the cerrado of Roraima.

49

## 50 **2. MATERIAL AND METHODS**

51


52 The experiment was conducted in the experimental field of Brazilian Agricultural Research  
53 Corporation (EMBRAPA), Água Boa - CEAB, in the municipality of Boa Vista - Roraima  
54 state; located at the geographical coordinates of reference: 02°49'11"N, 60°40'24"W and 85  
55 m of altitude, in a soil classified as Yellow Latosol, whose analysis of properties was the  
56 following: pH (H<sub>2</sub>O) = 5.4; Ca<sup>2+</sup> = 1.28 cmolc.dm<sup>3</sup>; Mg<sup>2+</sup>cmolc.dm<sup>3</sup> = 0.2 cmolc.dm<sup>3</sup>; K+  
57 cmolc.dm<sup>3</sup>; = 0.19 cmolc.dm<sup>3</sup>; Al<sup>3+</sup> = 0.1; cmolc.dm<sup>3</sup> (H + Al) = 2.62 cmolc.dm<sup>3</sup>; P<sub>2</sub>O<sub>5</sub> =  
58 14.18 mg.dm<sup>3</sup>; SB= 1.67 cmolc.dm<sup>3</sup>; T = 4.29 cmolc.dm<sup>3</sup>; t = 1.77 cmolc.dm<sup>3</sup>; V= 39% and  
59 m=6%, clay =136 g kg<sup>-1</sup>; silt = 29.1 g kg<sup>-1</sup> and sand = 834.7 g kg<sup>-1</sup>.

60

61 The climate of the region, according to the classification of Köppen, is Aw type, tropical  
62 rainy, with an average annual precipitation of approximately 1,700 mm and relative air  
63 humidity around 70% [9].

64

65 The climatic data referring to maximum and minimum temperatures, and rainfall occurred  
66 during the experimental period are described in Figure 1.



**Fig 1. Means of rainfall and maximum and minimum temperatures.**

67

68

69

70 The experimental design was a randomized complete block design in subdivided plots with  
 71 four replications. Plots consisted in the spacing (0.45, 0.55 and 0.65 m) and the subplots  
 72 were constituted by the cover plant species *Urocloa brizantha*, *Urocloa ruziziensis*, *Panicum*  
 73 *maximum* and the treatment without intercropping. The used soybean cultivar was BRS  
 74 Tracajá in two crops, from June to September 2015, and from May to September 2016.

75

76 The plant stand was the same for all treatments, varying only as for the number of plants per  
 77 linear meter, which were adjusted to the different spacing. The subplots occupied areas of  
 78 18.9 m<sup>2</sup> for the 0.45 m spacing; 23.1 m<sup>2</sup> for the 0.55 m spacing, and of 27.3 m<sup>2</sup> for the 0.65  
 79 m spacing. The useful area of each subplot consisted of 5.0 x 2.25 m (11.3 m<sup>2</sup>) for the 0.45  
 80 m spacing; 5.0 m x 2.2 m (11 m<sup>2</sup>) for the 0.55 m spacing, and 5.0 x 1.95 m (9.8 m<sup>2</sup>) for the  
 81 0.65 m spacing, consisting of five, four and three rows of soybean plants, respectively, in  
 82 which 0.50 m at each end of the subplots were excluded, for the realization of the  
 83 evaluations, corresponding to the useful area.

84

85 Before the sowing of soybean in 2015, the area was prepared with two disk plowing and one  
 86 with a leveler to revolve the soil, since it remained for six years without any cultivation.  
 87 Fertilization consisted of 100 kg ha<sup>-1</sup> of P<sub>2</sub>O<sub>5</sub>, in the source of simple superphosphate + 50  
 88 kg ha<sup>-1</sup> of FTE BR 12 + 10 kg ha<sup>-1</sup> of N (urea source) applied in the planting grooves, and  
 89 120 Kg ha<sup>-1</sup> of K<sub>2</sub>O in the source of potassium chloride, with 50% applied during planting and  
 90 50% during coverage, 30 days after emergence (DAE), together with seeds of the cover  
 91 species.

92

93 Soybean sowing was performed in open grooves with a mechanized ridger during the first  
 94 year of cultivation. Therefore, seeds were inoculated with *Bradyrhizobium japonicum*.  
 95 Subsequently, it was sown manually, using densities of 280,000 ha<sup>-1</sup> seeds, held in June  
 96 2015. In the second year of cultivation (2016), sowing was performed mechanically in May in  
 97 no-tillage, using a SEMEATO SAN 200 planter, over the straw formed by the cover species  
 98 from the previous year.

99

100 Covering species were sown 30 days after the emergence of the soybean seedlings (DAE),  
101 using 30 kg ha<sup>-1</sup> of seeds for the species *Urocloa brizantha* and *U. ruziziensis*, and 10 kg ha<sup>-1</sup>  
102 for cv. Massai, mixed with 60 kg ha<sup>-1</sup> of K<sub>2</sub>O planted between the rows of soybean plants.  
103 Weed control was performed at 25 DAE, at stage V4, using the herbicides Flex (Fomesafen)  
104 and Fusilade (Fluazifop-p-butyl), at doses recommended by the manufacturers.

105  
106 In the second cultivation year (2016), according to the covering obtained from the previous  
107 planting, forage was dried with Glyphosate + Flumyazin (Flumioxazin), then soybean was  
108 planted, and after 20 (DAE), Flex (Fomesafen) + Verdict (Haloxifop-Methyl) was applied.  
109

110 During the development of the crop and after the harvesting of soybean, the following  
111 agronomic characteristics were evaluated: plant height, evaluating ten random plants in the  
112 useful area, measuring them from the neck of the plant until the end of the main stem;  
113 number of grains per pod - the total number of grains from ten plants was counted, and the  
114 result was divided by the total number of pods; number of pods per plant - ten random plants  
115 were collected in the useful area of the sub-plot, obtained by counting the total number of  
116 pods and calculating the average; 100-grain weight, determined by weighing one hundred  
117 grains from the useful area, later corrected to 13% moisture; plant dry matter - ten plants  
118 were randomly collected, dried in an oven until constant weight and weighed on a precision  
119 scale; insertion of the first pod, determined from the collection of ten random plants in the  
120 useful area of each subplot, measuring from the neck of the plant until the insertion of the  
121 first pod; grain yield - the grains harvested from the useful area of each plot were weighed,  
122 estimating the production for one hectare, and correcting grain moisture to 13%.  
123

124 One-hundred twenty days after the harvest of soybean, the dry matter of the cover crop area  
125 and the spontaneous vegetation contained in the treatments without intercropping were  
126 evaluated. To determine the dry matter of the covering species, samples were collected  
127 using a 0.50 x 0.50 m square iron, according to the Braun-Blanquet methodology (1950)  
128 [10]. After that, they were taken to the laboratory in order to determine the dry matter of  
129 plants, through oven drying until constant weight at a temperature of 65°C, and then they  
130 were weighed on a precision scale.  
131

132 Data on the production components of soybean and the dry matter of forage species and  
133 spontaneous vegetation were submitted to analysis of variance using the F test. These data  
134 refer to the average of two cultivation years (2015 and 2016). For the comparison between  
135 the means, the Tukey's test was carried out at 5% probability, with the help of the SISVAR  
136 computational application. The variable about shoot dry matter of covering species and  
137 spontaneous plants was transformed into kg ha<sup>-1</sup> to discuss data.  
138

### 139 **3. RESULTS AND DISCUSSION**

140  
141 The height of plants was influenced by the covering species intercropped with soybean,  
142 whose means are presented in Table 1.  
143

144 The greatest PH of soybean intercropped with *P. maximum* cv capim massim may be related  
145 to the characteristic of the species. The forage plant cv. massai presents a smaller size,  
146 forming clumps with a mean height of 0.60 m, and presenting fine leaves, measuring 1 cm in  
147 width [11]. Possibly, these characteristics may have contributed to a smaller competition with  
148 the intercropped species, since soybean reaches a greater height.  
149

150 For the intercrop with the species *U. ruziziensis*, due to a slower initial growth, soybean  
151 probably showed greater vigor in the initial development of plants, but did not differ in height  
152 from *P. maximum*, as well as the low spontaneous vegetation in the area of treatments

153 without intercropping, which were basically composed of lower plants where there was  
154 greater competition of the culture.

155

156 **Table 1. Average plant height (PH), number of pod per plant (NPP) and yield of**  
157 **soybean crop cv. BRS Tracajá, intercropped with covering plants in Boa Vista -**  
158 **Roraima state, 2017**

159

Covering plants	PH (cm)	NPP	Yield (kg ha <sup>-1</sup> )
<i>Urocloa brizantha</i>	0.83 b*	59.0b	2631.1 b
<i>Urocloa ruziziensis</i>	0.87 ab	59.2ab	2880.8 ab
<i>Panicum maximum</i>	0.89 a	62.9ab	2713.9 ab
Spontaneous vegetation	0.85 ab	67.1a	2920.4 a
VC%	3.6	11.3	9.1

160

\*Means followed by the same lowercase letter in the column do not differ by Tukey's test, at 5% probability.

161

162

163 The lower PH found for the intercrop with *U. brizantha* can be justified by the characteristics  
164 regarding the forage cultivar introduced in the intercrop with BRS Tracajá soybean. It is  
165 possible to state that, under these conditions, the intra-species competition was significant,  
166 but with an acceptable height of soybean plants.

167

168 The number of pods per plant (NPP) was influenced by the spacing (Table 2) and also by  
169 the intercrop with covering species (Table 1). A significant difference between spacing was  
170 also verified by Silva et al. (2013) [12], in which there was a higher NPP in a spacing of 0.50  
171 m.

172

173 **Table 2. Average number of pods per plant intercropped with three types of spacing**  
174 **(cm) in between rows of soybean cv. BRS Tracajá, in Boa Vista - Roraima state, 2017**

175

Spacing (m)	Number of pods per plant
0.45	59.5 b*
0.55	58.4 b
0.65	68.2 a
VC %	11.1

176

\*Means followed by the same lowercase letter in the column do not differ by Tukey's test, at 5% probability.

177

178

179 As for the different covering species used in the intercrop with soybean, it is possible to  
180 observe that the spontaneous vegetation, *U. ruziziensis* and *P. maximum*, provided soybean  
181 with the highest NPP and the last two species did not differ from *U. brizantha* (Table 1).  
182 Among the elements used in the production factor, NPP is the characteristic that most  
183 contributes to the grain yield in the soybean crop, since it presents a higher correlation with  
184 production [13].

185

186 There was an interaction between the used spacing and the covering plants for the 100-  
187 grain weight (W100G). When the spacing was split within each covering, it was possible to  
188 observe that *U. ruziziensis* and the spontaneous vegetation influenced the W100G of the  
189 culture (Table 3).

190

191 In the intercrop with *U. ruziziensis*, soybean reached a higher W100G at the spacing of 0.45  
192 and 0.55 m. Possibly, a smaller spacing allowed lower weed interference in the soybean  
193 crop, due to the closing of the crop canopy. As for *U. ruziziensis*, there was a smaller initial

194 development at these spacing. As for the outcome of the covering species within each  
195 spacing level, no significant difference was observed (Table 3).

196  
197  
198  
199  
200

**Table 3. Averages of the 100-grain weight obtained according to the interaction between spacing and covering plant intercropped with soybean cv. BRS Tracajá under different spacings in Boa Vista - Roraima, 2017**

Covering plants	100-grain weight (g)		
	45 cm	55 cm	65 cm
<i>Urocloa brizantha</i>	11.8 aA*	12.1 aA	11.8 aA
<i>Urocloa ruziziensis</i>	13.3 aA	11.3 abA	10.8 bA
<i>Panicum maximum</i>	11.9 aA	12.3 aA	12.8 aA
<i>Spontaneous vegetation</i>	11.7 abA	13.5 aA	11.3 bA
VC1%		7.88	
VC2%		9.23	

201 \*Means followed by the same lowercase letter on the line and uppercase in the column do not differ by  
202 Tukey's test, at 5% probability.

203  
204  
205  
206  
207  
208  
209  
210

Komatsu et al. (2010) [14], while studying the effect of plant spacing on the behavior of specific growth soybean cultivars, observed a greater grain weight when the 0.45 m spacing was used, highlighting this effect among the characteristics of long-cycle soybean cultivars. According to Bianchi et al. (2010) [15], crops with good potential for production cause greater reduction of environmental resources, reducing their availability to other competing species and thereby becoming more competitive with weeds.

211  
212  
213  
214  
215  
216

No significant differences were found as for the W100G intercropped with the species *U. brizantha* and *P. maximum* cv. massai (Table 3). This result may be related to the genetic limit of the forage cultivar and/or species. In a study conducted by Castagnara et al. (2014) [16], it was also not possible to find differences in terms of W100G in the joint sowing of soybean and *U. brizantha*.

217  
218  
219  
220  
221  
222  
223

Grain yield was influenced by the covering crops; the cultivation without intercropping was the best treatment, followed by the species *U. ruziziensis* and *P. maximum* (Table 1). Productivity is closely linked to the production components of soybean and depends directly on the interaction of the genotype with the environment [17]. According to Albuquerque et al. (2012) [18], Castagnara et al. (2014) [16], Albuquerque et al. (2015) [19] and Werner et al. (2017) [3], large crops show higher yields in single crops.

224  
225  
226

Table 4 presents the values about the first pod insertion (FPI) characteristic in the soybean crop, for the interaction between spacing and covering.

227  
228  
229  
230  
231  
232  
233  
234  
235

As for the spacing within each covering level, it was observed that the spacing of 0.65 m influenced the intercrop when the *P. maximum* species was used, decreasing the height of the FPI. The spacing with the highest FPI height was 0.45 and 0.55 m. As for the other covering species, no significant differences were observed (Table 4). A greater spacing allowed lower plants, compared to those of the 0.55 and 0.45 m spacing; thus, there was a small variation in FPI. According to Cruz et al. (2016) [20], the importance of evaluating this variable informs if the minimum height may or may not provide losses during the harvesting process by the cutting bar of the harvester.

236  
237  
238

In the 0.55 m spacing, *U. brizantha* negatively influenced the FPI, resulting in the lowest height, but with similar values to the other treatments (Table 4). This effect may be related to the competition of the intercrop and the variation in the environment, modifying the height

239 of plants. Torres et al. (2015) [21] state that the environmental factors that interfere in the  
 240 FPI are the same that can influence the height of plants, so it is possible that the height of  
 241 the first pod has undergone a variation according to the height of soybean plants.

242  
 243  
 244  
 245  
 246  
 247

**Table 4. Averages of the first pod insertion (FPI) obtained according to the interaction between spacing and covering plant intercropped with soybean cv. BRS Tracajá in three spacings between rows, in Boa Vista - Roraima state, 2017**

Covering plants	First pod insertion		
	45 cm	55 cm	65 cm
<i>Urocloa brizantha</i>	15.9 aB*	16.6 aB	16.3 aB
<i>Urocloa ruzizensis</i>	17.9 aA	17.5 aAB	18.1 aA
<i>Panicum maximum</i>	18.4 aA	18.4 aA	16.5 bB
<i>Spontaneous vegetation</i>	17.1 aAB	17.2 aAB	16.4 aB
VC1%		5.78	
VC2%		4.54	

248 \*Means followed by the same lowercase letter on the line and uppercase in the column do not differ by  
 249 Tukey's test, at 5% probability.

250

251 The *U. brizantha* species, in general, was the one that influenced in terms of lower height in  
 252 the FPI, mainly due to the intense competition that occurs with the culture. *U. brizantha* is  
 253 more demanding for light, thus becoming more competitive for the solar radiation that  
 254 reaches the soil for germination and vegetative development, and the FPI has a direct  
 255 correlation with the use of light in the lower part of the canopy; thus, the more light reaches  
 256 the lower part of the canopy of the soybean crop, the lower the node of the first pod and,  
 257 consequently, the height of the insertion of the first pod.

258

259 A study by Pereira et al. (2011) [22] showed a negative influence on the intercropping with  
 260 *U. decumbens* species, causing a significant effect, and reducing the height of the first pod  
 261 to 11.1 cm, when this forage was sown in the soybean rows, 25 days after sowing.

262 The shoot dry matter yield of covering plants had a significant effect for the interaction  
 263 between spacing and covering plants (Table 5).

264

**Table 5. Averages of the dry matter of covering species (kg ha<sup>-1</sup>), obtained according to the interaction between spacing and covering plant intercropped with soybean cv. BRS Tracajá, in three spacings between rows, in the experimental field of Embrapa, in Boa Vista - Roraima state, 2017**

265  
 266  
 267  
 268  
 269

Covering plant	Dry matter (kg ha <sup>-1</sup> )		
	45 cm	55 cm	65 cm
<i>Urocloa brizantha</i>	74.99 bA*	83.56 abA	92.44 aA
<i>Urocloa ruzizensis</i>	54.60 aB	63.74 aB	68.08 aB
<i>Panicum maximum</i>	69.58 aAB	64.95 aB	70.21 aB
<i>Vegetação espontânea</i>	17.75aC	16.70 aC	15.03 aC
VC1%		14.7	
VC2%		15.1	

270 \*Means followed by the same lowercase letter on the line and uppercase in the column do not differ by  
 271 Tukey's test, at 5% probability.

272

273 Opposite results were obtained by Mata et al. (2012) [23] with lower values for the same  
 274 variable, which can be explained by the smaller spacing between soybean rows (0.40 m)

275 used by these authors, and the 20 to 30-day sowing gap period, which favored the  
276 development to the detriment of forage.

277

278 The *Urocloa* species show greater root growth, which may result in better development  
279 conditions during the dry season [24].

280

281 *U. ruziziensis* becomes promising in the production of straw when intercropped with soybean  
282 in the no-tillage system. Pacheco et al. (2011) [25] mention that out of the species used to  
283 form straw in the off-season, *U. Ruziziensis* is important; even with a low initial development,  
284 it has good regrowth capacity and dry matter gains, thus being an alternative to intercropping  
285 and no-till systems.

286

#### 287 **4. CONCLUSION**

288

289 Cultivar BRS Tracajá presents better grain yield in the single crop, and intercropped with  
290 *Urocloa ruziziensis* and *Panicum maximum*. The highest dry matter yield occurs for the  
291 *Urocloa brizantha* species; however, it causes the greatest reduction in soybean yield. The  
292 *U. ruziziensis* and *P. maximum* species present the best use potential to establish  
293 themselves in intercrop with soybean, reaching good dry matter productivity and less  
294 interference in soybean production components. The used spacing does not influence the  
295 productivity of cultivar BRS Tracajá.

296

297

298

#### **COMPETING INTERESTS DISCLAIMER:**

299

300

301

302

303

304

305

**Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.**

306

307

308

#### **AUTHORS' CONTRIBUTIONS**

309

310

311

This work was carried out in collaboration between all authors. All authors read and approved the final manuscript.

312

#### **REFERENCES**

313

314

1. National Supply Company (CONAB). Monitoring of Brazilian harvest: grains, fifth survey, Safra 2016/17. 2017; 4 (5): 166.

315

316

317

2. Debiasi H, Franchini JC. Soil physical attributes and soybean yield in a crop-livestock integration system with brachiaria and soybean. Rural Science. 2012; 42 (7): 1180-1186. English.

318

Available: <http://dx.doi.org/10.1590/S0103-84782012000700007>



- 319 3. Werner F, Balbinot Junior AA, Franchini JC, Ferreira AS, Silva MAA. Agronomic  
320 performance of soybean cultivars in an agroforestry system. *Tropical Agriculture Research*.  
321 2017; 47 (3): 279-285.
- 322 Available: <http://dx.doi.org/10.1590/1983-40632016v4745937>
- 323 4. Lithourgidis AS, Dordas CA, Damalas CA, Vlachostergios D. Annual intercrops: An  
324 alternative pathway for sustainable agriculture. *Australian Journal of Crop Science*. 2011; 5:  
325 396-410.
- 326 5. Queiroz RF, Chioderoli CA, Furlani CEA, Netherlands HV, Zerbato C. Corn intercropping  
327 with *Urochloa ruziziensis* under no-tillage. *Tropical Agriculture Research*. 2016; 46 (3): 238-  
328 244. English.
- 329 6. Tourino MCC, Rezende PM, Salvador N. Spacing, density and uniformity of sowing on  
330 yield and agronomic characteristics of soybean. *Pesquisa Agropecuária Brasileira*. 2002; 37  
331 (8): 1071-1077. English.
- 332 7. Procópio SO, Balbinot Junior AA, Debiasi H, Franchini JC, Panison F. Double-row seeding  
333 and reduced spacing in soybean cultivation. *Revista Agro @ mbiente online*. 2014; 8 (2):  
334 212-221. English.
- 335 8. Balena R, Giacomini CT, Bender AC, Nesi CN. Sowing time and row spacing in soybean  
336 yield. *Unoesc & Science*. 2016; 7 (1): 61-68. English.
- 337 9. Araújo WF, Andrade Junior ASD, Medeiros RD, Sampaio RA. Probable rainfall in Boa Vista,  
338 State of Roraima, Brazil. *Brazilian Journal of Agricultural and Environmental Engineering*.  
339 2001; 5 (3): 563-567. Portuguese
- 340 Available: <http://dx.doi.org/10.1590/S1415-43662001000300032>
- 341 10. Braun-Blanquet J. *Sociologia vegeta: studies of plant communities*. Buenos Aires: Acme  
342 Agency; 1950.
- 343 11. Brazilian Agricultural Research Corporation (EMBRAPA). *Capim Massai (Panicum*  
344 *maximum Jacq.): New forage for pasture diversification in Acre*. 2001; 41: 16.
- 345 12. Silva WB, Lima LB, Petter FA, Andrade FR. Initial development of *Urochloa ruziziensis*  
346 and agronomic performance of soybean in different spatial arrangements in the cerrado  
347 Mato-Grossense. *Bragantia*. 2013; 72 (2): 146-153. English.
- 348 Available: <http://dx.doi.org/10.1590/S0006-87052013000200006>

- 349 13. Nogueira APO, Sedyama T, Sousa LB, Hamawaki OT, CD Cross, Pereira DG, Matsuo, É.  
350 Track analysis and correlations between characters in soybean cultivated at two sowing  
351 times. *Bioscience Journal*. 2012; 28 (6): 877-888. English.
- 352 14. Komatsu RA, Guadagnin DD, Borgo MA. Effect of plant spacing on the behavior of  
353 determined growth soybean cultivars. *Digit @ I field*. 2010; 5 (1): 50-55. English.
- 354 15. Bianchi MA, Fleck NG, Lamego FP, Agostinetto D. Plant arrangement papers and  
355 soybean cultivar as a result of interference with competing plants. *Weed plant*. 2010; 28  
356 (4): 979-991. English.
- 357 Available: <http://dx.doi.org/10.1590/S0100-83582010000500006>
- 358 16. Castagnara DD, Bulegon LG, Zoz T, Rossol CD, Berte LN, Oliveira PSR, Neres MA.  
359 Consortium soybean cultivation with brachiaria. *Bioscience Journal*. 2014; 30: 168-177.  
360 English.
- 361 17. Cruz TV, Peixoto CP, Martins MC, Pinto MDFS. Components of soybean production in  
362 different sowing times in western Bahia. *Bioscience Journal*. 2010; 26 (5): 709-716. English.
- 363 18. Albuquerque JAA, Sedyama T, Alves JMA, Silva AA, Uchôa SCP. Cultivation of cassava  
364 and beans in intercropping systems held in Coimbra, Minas Gerais, Brazil. *Agronomic  
365 Science Journal*. 2012; 3 (3): 532-538. English.
- 366 19. Albuquerque JAA, Oliva LSDC, Alves JMA, Uchôa SCP, Melo DAD. Cultivation of cassava  
367 and cowpea in intercropping systems held in Roraima's savannah, Brazil. *Agronomic  
368 Science Journal*. 2015; 46 (2): 388-395. English.
- 369 Available: <http://dx.doi.org/10.5935/1806-6690.20150018>
- 370 20. Cross SCS, Sena Junior DG, Santos DMA, Lunezzo, LO, Machado CG. Soybean cultivation  
371 under different sowing densities and spatial arrangements. *Journal of Neotropical  
372 Agriculture*. 2016; 3 (1): 1-6. English.
- 373 21. Torres FE, David GV, Teodoro PE, Ribeiro LP, Correa CG, Luz Júnior RA. Agronomic  
374 performance and genetic dissimilarity among soybean genotypes. *Journal of Agricultural  
375 Sciences*. 2015; 38 (1): 111-117. English.
- 376 22. Pereira RG, Albuquerque AW, Souza ROS, Silva AD, Santos JPA, Barros ES, Medeiros  
377 PVQ. Soil management systems: soybean [*Glycine max* (L.)] intercropped with *Bachiaria  
378 decumbens* (STAPP). *Tropical Agriculture Research*. 2011; 41 (1): 44-51. Portuguese

379 23. Mata JF, Erasmo EAL, Siebeneichler SC, Saraiva AS, Gonçalves RC. Analysis of Brachiaria  
380 growth in different densities and sowing times in consortium with the soybean crop.  
381 Applied Research & Agrotechnology. 2012; 4 (3): 7-30. English.

382 Available: <https://doi.org/10.5777/paet.v4i3.1494>

383 24. Pacheco LP, Pires FR, Monteiro FP, Procópio SO, Assis RD, Petter FA. Depth of sowing  
384 and initial growth of forage species used to cover the soil. Science and Agrotechnology.  
385 2010; 34 (5): 1211-1218. English.

386 Available: <http://dx.doi.org/10.1590/S1413-70542010000500019>

387 25. Pacheco LP, Barbosa JM, Leandro WM, Machado PLOA, Assis RL, Madari BE, Petter FA.  
388 Production and nutrient cycling by cover crops in upland rice and soybean crops. Brazilian  
389 Journal of Soil Science. 2011; 35: 1787-1799. English.

390

UNDER PEER REVIEW