

**INFLUENCE OF DIFFERENT BANANA 'PRATA-ANÃ' BUNCH AGES ON
POST-HARVEST QUALITY**

ABSTRACT

Objective: Thus, the objective was to verify the ideal harvest season of banana 'Prata-Anã' grown in northern Minas Gerais. Bananas were harvested at different ages – 16, 17, 18, 19 and 20 weeks after inflorescence emission –and stored in cold room for 25 days

Study Design: The employed experimental design was the completely randomized design was used in a 5x5 factorial scheme, with five bunch ages and five assessment days.

Study Location and Duration: The experiment was run in an area with banana trees planted 20 months beforehand, located at Unimontes's Experimental Farm, at 530m of altitude, with coordinates being -15°43'46.99" south latitude and -43°19'17.61"west longitude, between April and November 2017.

Methodology: The bananas bunches the were marked weekly from April 14 to May 12, and week days were standardized for each marking. Five bunch ages were defined – 16, 17, 18, 19 and 20 weeks after inflorescence emission – for harvest. For differentiation of emerged bunches, tapes of different color were used. When the bunches marked in the first week completed 20 weeks, all bunches were harvested, which happened on September 1. After harvested the fruits were subjected to storage in refrigerated chamber at 10°C ± 1°C and relative humidity of 90% +5% for 25 days. After being stored for 25 days, the bananas were taken out of the chamber and exposed to a room temperature of 25oC, which they physical and chemical analyses were performed for 9 days, with a two-day interval in between, simulating the marketing period.

Results: The Fruits from 16-week bunches were superior in physical and chemical characteristics compared to other ages.

Conclusion: Bunch harvest age had a direct influence on post-harvest quality of bananas

'Prata-Anã'. Fruits from 16-week bunches were superior in physical and chemical characteristics compared to other ages, meaning a longer post-harvest life.

Keywords: storage, *musa* ssp, maturation stage

1. INTRODUCTION

Banana trees (*Musa* spp.) are the most relevant fruitful trees worldwide and its production is mostly concentrated in tropical countries, being the second most produced fruit in Brazil [1]. According to [2], in 2016, Brazil was the third among countries with the highest banana production, behind India and China only; besides, this fruit appears among the three most produced tropical fruits, alongside orange and pineapple.

The southwest region is the second greatest banana producer, with the north of Minas Gerais being a major producing pole in Brazil, with a high social and economic importance for the region. Banana 'Prata-Anã' (AAB) and its different clones are the most prevalent in cultivation, with good market acceptance due to their excellent quality attributes, being considered elementary in nutrition.

For being climacteric fruits classified as perishable, bananas require techniques that slow down their rapid ripening, preventing post-harvest losses, especially while being transported to more distant consuming markets.

Harvesting fruits at proper maturation stages is determinant to maintaining post-harvest quality. Maturation point is the ideal harvest moment without the occurrence of damages, which provides fruits with a longer preservation period. This point is usually reached when the fruit becomes physiologically mature, which corresponds to its maximum size and weight, but does not have desirable characteristics for marketing and consumption. Later, the fruit continues to go through transformations, ripens naturally and becomes suitable for consumption. However, the ideal harvest point depends on correlations between physiological characteristics inherent of each variety, the ideal maturation stage, and post-harvest preservation technologies applied. In banana trees, fruits harvested prematurely may not be physiologically developed, which hinders their ripening process and final quality [3]. Nevertheless, harvesting overripe fruits leads to rapid quality loss, reducing their marketing period.

In addition to defining the best harvest stage, another way of reducing damages and prolonging storage period is to keep fruits refrigerated. Refrigeration is considered

37 one of the most efficient methods for fruit preservation, maintaining the fruit's desirable
38 characteristics, similar to those of its early stage, due to a delayed maturation process.

39 A fruit's external characteristics – which relate to its appearance as well as size,
40 shape, color, lightness, absence of imperfections – and internal characteristics –
41 perceived in how it tastes, smells and feels – are the main attributes evaluated by
42 consumers, who demand for quality during purchase [3].

43 Therefore, determining the ideal harvest point of bunches is imperative, when
44 the fruit reaches its physiological maturation, that is, its maximum size and weight,
45 which will influence its post-harvest quality and resistance for a longer preservation
46 period.

47 The present study aimed to determine the ideal harvest season of banana 'Prata-
48 Anã' bunches by means of physical and chemical analyses on fruits maturing under
49 cultivation conditions in the north of Minas Gerais, allowing for their maximum
50 utilization in order to provide the final consumer with quality, in accordance with their
51 demands and preferences.

52 **2. MATERIAL AND METHODS**

53 The experiment was run in an area with banana trees planted 20 months
54 beforehand, located at Unimontes's Experimental Farm, at 530m of altitude, with
55 coordinates being -15°43'46.99" south latitude and -43°19'17.61" west longitude. After
56 inflorescence emission, the banana trees were randomly selected and marked through
57 criterion proposed by [4] for sourcing of bananas at different bunch ages. They were
58 marked weekly from April 14 to May 12, and week days were standardized for each
59 marking. Five bunch ages were defined – 16, 17, 18, 19 and 20 weeks after
60 inflorescence emission – for harvest. For differentiation of emerged bunches, tapes of
61 different color were used. When the bunches marked in the first week completed 20
62 weeks, all bunches were harvested, which happened on September 1. After harvest, the
63 bunches were separated into bouquets with four fruits each and washed in water and
64 neutral detergent at 0.2% for latex coagulation and superficial cleaning. The bouquets
65 were then immersed in a solution of Imazalil fungicide, at a dose of 2mL.100mL⁻¹ of
66 water at room temperature, and dried outdoors. Each bouquet was stored in low-density
67 polyethylene packs measuring 25µm in thickness and put inside a standard cardboard
68 box for export. The fruits were subjected to storage in refrigerated chamber at 10°C ±
69 1°C and relative humidity of 90% ±5% for 25 days. After being stored for 25 days, the

70 bananas were taken out of the chamber and packs and exposed to a room temperature of
71 25°C, after which they were analyzed for 9 days, with a two-day interval in between,
72 simulating the marketing period. The following analyses were then conducted:

73 Peel and pulp firmness: determined by maximum penetration strength with a flat
74 tip measuring 4 mm in diameter, placed 10mm away from the fruit, with the aid of a
75 Brookfield digital penetrometer, model CT3 10 KG; measures were taken from the
76 medium area of the fruit with and without peel, and results were expressed as Newton
77 (N).

78 Peel color: analyzed on Color Flex digital colorimeter, model CT3 10 KG,
79 which expresses color using three parameters: L*(lightness), which ranges from 0
80 (black) to 100 (white); a* (transition from green (-a*) to red (+a*)) and b* (transition
81 from blue (-b*) to yellow (+b)). Based on L*, a* and b* values, hue angle (°h) and
82 chroma saturation index (C*) were calculated.

83 Soluble solids: analysis performed by means of Reichert digital refractometer,
84 using banana pulp mashed in a food processor, with results expressed as °Brix.

85 pH: determined by electrometric method in potentiometer, using 10g of mashed
86 sample with 90mL of distilled water, in accordance with [5].

87 Titratable acidity (TA): determined using analyte (10g of pulp homogenized and
88 diluted in 100mL), titrated with sodium hydroxide standard solution (NaOH) at 0.1N,
89 having phenolphthalein as indicator. Results were expressed as malic acid percentage.
90 All methodology used complies with [5].

91 Amide: chemically extracted and spectrophotometrically determined according
92 to a chemical method by [6]. It was determined at 510nm and results were expressed as
93 percentage.

94 Total sugars (TS): extracted with ethyl alcohol and determined by the Antrona
95 method [7]. The sample was subjected to reading on spectrophotometer at 620nm, and
96 results were expressed as percentage.

97 Reducing sugars (RS): determined by Nelson's methodology [6]. Reducing
98 sugar content was calculated by spectrophotometry at 510nm, and results were
99 expressed as percentage.

100 Non-reducing sugars (NRS): obtained by differences between total sugars and
101 reducing sugars, as per the formula below:

102 Non-reducing sugars = Total sugars – Reducing sugars x 0.95.

103 Electrolyte extravasation: it was determined according to [8]; a peel disc was
104 removed per damaged area from each fruit, measuring 1cm in diameter, with the aid of
105 a metal punch. This section was washed in distilled water and superficially dried on
106 absorbent paper, then incubated for 2 hours in a capped test tube containing 18mL of
107 distilled water, under ambient conditions. After this period, electrical conductivity was
108 measured on a SCHOT conductivity meter, model CG 853. Later, the tubes containing
109 the peel samples were autoclaved at 121°C and 1.5atm for 30 minutes. After
110 autoclaving, electrical conductivity was read again. Results were expressed as the ratio
111 between values obtained in the first and second measurements multiplied by 100.


112 The experimental design employed was the completely randomized type (CRD),
113 in a 5x5 factorial scheme, with five bunch ages (16, 17, 18, 19 and 20 weeks after
114 inflorescence emission) and 5 assessment periods (1, 3, 5, 7 and 9 days after storage).
115 Four repeats were used, and the experimental unit was composed of four fruits. Data on
116 the variables were subjected to tests for analysis of homogeneity of variance [9], residue
117 normality by the Shapiro-Wilk test [10] and model non-additivity [11]. Results were
118 then subjected to analysis of variance (ANOVA), considering as sources of variation
119 bunch ages, assessment days after storage, and interaction between bunch ages and days
120 after storage, tested at 5% probability. Interaction was sliced or not, depending on
121 significance; regression analysis was conducted, and models were chosen based on
122 significance, coefficient of determination and potential to explain the biological
123 phenomenon. The variables were studied using statistical program SISVAR.

124

125 **3. RESULTS AND DISCUSSION**

126 Analyzing peel firmness, significant interaction was observed between bunch
127 age and assessment day, factors that simulate the marketing period of fruits; fruits from
128 bunches younger than 19 weeks were firmer on the 9th day after storage, presenting
129 values of 17.39N, 21.36N, 20.86N and 15.77N, for bunch ages corresponding to 16, 17,
130 18 and 19 weeks, respectively; bunches with 17 weeks were superior to the other
131 treatments, with longer shelf life (Figure 1). On the other hand, 20-week bunches were
132 less firm, reaching 6.15N on the last assessment day. [12], while working with banana
133 ‘BRS Tropical’, found that fruits with greater development at the harvest point had

134 reduced firmness. According to [13] and [14], firmness reduction is related to amide
135 hydrolysis and solubilization of pectic substances, as well as to water loss.




136

137 **Figure 1:** Peel firmness of banana ‘Prata-Anã’ harvested at different bunch ages against days after
138 storage

139

140

141 As for pulp firmness, significant interaction was observed between bunch ages
142 and storage days for this variable, which reduced as bunch age increased; on the first
143 assessment day, for the ages of 16, 17, 18, 19 and 20 weeks, the values found were
144 19.36N, 18.27N, 17.18N, 16.10N and 15.01N, respectively (Figure 2). It was possible to
145 observe on the days after storage a sharp firmness reduction in fruits harvested with 18,
146 19 and 20 weeks, which showed respective values of 2.83N, 1.74N and 0.70N,
147 compared to those harvested with 16 and 17 weeks, which presented values of 5.00N
148 and 3.92N, respectively; fruits from the bunch with 16 weeks were firmer than the
149 others. The results of this experiment corroborate with study by [15], which related
150 firmness loss of banana ‘Prata-Anã’ pulp to older bunch harvest age, stressing that high
151 storage temperatures also contribute to firmness loss, and it is possible to observe that
152 not even temperatures under 10°C were enough to prevent softening in fruits from 20-
153 week bunches, that is, these fruits had a greater firmness loss.




154

155

156 **Figure 2:** Pulp firmness of bananas ‘Prata-Anã’ harvested at different bunch ages against days after
 157 storage

158

159 For color-describing variables, the hue angle parameter defines the basic color of
 160 samples and represents the average hue of the banana samples; results were significant
 161 for interaction. The hue angle values found in the banana peels at different bunch ages
 162 (16, 17, 18, 19 and 20 weeks) dropped from 106.2°, 104.7°, 103.2°, 101.7° and 100.2°
 163 to 85.3°, 83.8°, 82.4°, 80.9° and 79.4°, respectively, with this drop occurring from the
 164 1st to the 9th day after storage, which varied by treatment (Figure 3). This behavior is
 165 expected because hue angle values close to 100° presented a greenish color, and as
 166 values move further or closer to 80° the color of the fruit turns yellowish, evidencing
 167 ripening. Fruits harvested with a bunch age of 16 weeks had hue angle values higher
 168 compared to other ages on the last assessment day, indicating their preservation and
 169 allowing for them to be marketed for a longer period. According to [16], fruit color is an
 170 important parameter to track the ripening process, which, in the case of bananas,
 171 corresponds to yellow, due to chlorophyll degradation and carotenoid synthesis, besides
 172 being a criterion used to characterize maturation stages.




173

174 **Figure 3:** Hue angle (a) of bananas ‘Prata-Anã’ harvested at different bunch ages

175

176 Figure 4 displays chromaticity values, which express color intensity, that is,
 177 saturation in terms of pigment. Significant difference is observed between bunch
 178 ages; values stood at 37.64, 38.84, 40.04, 41.25 and 42.45 with 16, 17, 18, 19 and 20
 179 weeks of age, respectively. These results are higher than those found by [3] while
 180 analyzing chromaticity in peels of banana ‘Prata-Anã’ stored at 10°C, with 18, 19
 181 and 20 weeks of development, finding estimated mean values of 33.67, 33.87 and
 182 34.61, respectively.

183




184

185 **Figure 4:** Chromaticity of bananas 'Prata-Anã' harvested at different bunch ages

186

187 The soluble solids variable presented significant interaction, being influenced by
 188 harvest ages and days after harvest. Figure 5 shows the behavior of soluble solid mean
 189 values, and it is possible to observe an increase in soluble solid content as fruits ripen;
 190 from the 1st to the 9th assessment day for all treatments (16, 17, 18, 19 and 20 weeks of
 191 bunch age), the values found were 4.08-16.73, 5.33-17.98, 6.88-19.53, 8.73-21.38, and
 192 10.88-23.53° Brix, respectively. Bananas have a high amide content when green and, as
 193 they ripe, amide is hydrolyzed into simple sugars for it to be used in fruit respiration,
 194 thus raising soluble solid content during maturation. The results observed on the last
 195 assessment day in the present experiment are similar to those found by [17] while
 196 working with fruits of banana trees 'Maravilha' and 'Preciosa', which showed values
 197 between 18.85 and 23.31 after cultivation during the 1st and 2nd production cycles, in the
 198 Submédio São Francisco River.




199

200 **Figure 5:** Soluble solid content in bananas ‘Prata-Anã’ harvested at different bunch ages against days
 201 after storage

202

203 There was significant interaction for pH between bunch ages and assessment
 204 days after storage. Figure 6 shows the behavior of values obtained for banana ‘Prata-
 205 Anã’ pH throughout the assessment days in relation to bunch ages. For all treatments, it
 206 is possible to observe a rapid decline in values obtained 9 days after storage, from 6.04
 207 to 5.10, 6.07 to 5.14, 6.04 to 5.11, 5.94 to 5.01 and 5.78 to 4.84 for bunches with 16, 17,
 208 18, 19 and 20 weeks, respectively. According to [18], while working with banana tree
 209 fruits, found pH values between 5.28 and 5.60, close to those found in the present study.
 210 [19], while working with bananas ‘Prata-Anã’ stored for 14 days under different
 211 controlled atmosphere conditions, found a mean pulp pH value of 4.25 after 3 days after
 212 atmosphere removal and maintenance in ambient atmosphere. According to [16], during
 213 the maturation phase of fruits there is an accumulation of soluble sugars, precursors of
 214 organic acids, with predominance of malic acid, which leads to a pH reduction
 215 throughout ripening.




216

217 **Figure 6:** pH values in bananas ‘Prata-Anã’ harvested at different bunch ages against days after storage

218

219 Titratable acidity showed significant interaction and associated with increased
 220 malic acid concentration. The graph reveals an increase in malic acid content in all
 221 fruits as they ripen and the bunches becomes older. For the bunch ages of 16 and 20
 222 weeks, 0.17g and 0.26g of malic acid were found per 100g of pulp, respectively, on day
 223 1. On the 9th assessment day, which simulates the marketing period, it is possible to
 224 observe that the bunch age of 16 weeks presented a value of 0.40g, whereas the bunch
 225 age of 20 weeks presented a value of 0.49g of malic acid per 100g of pulp, superior to
 226 the other bunch ages, evidencing faster ripening and shorter shelf life (Figure 7). The
 227 mean titratable acidity values observed in the present study 9 days after storage were
 228 similar to those found by [20] while working with ‘Prata-Anã’ in two production
 229 cycles. These results are in line with [21] in that unripe bananas have low acidity, and
 230 during ripening this acidity slowly increases until reaching a maximum value when the
 231 fruit is ripe and later falls with its senescence. Organic acid decline has been attributed
 232 to respiration or sugar conversion that occurs when banana tree fruits are ripening.
 233 These acids provide a sugar-acid balance, which results in a more tasteful fruit when
 234 ripe [22].




235

236 **Figure 7:** Titratable acidity content in bananas ‘Prata-Anã’ harvested at different bunch ages against days
 237 after storage

238

239 As for amide content, there was significant interaction as it was influenced by
 240 harvest seasons and days after storage in cold chamber (Figure 8). From the values
 241 obtained, a slight drop was seen in the amide content of the fruits as bunch age
 242 increased, that is, bunches harvested later, because, as fruits ripen, amide rapidly
 243 degrades to be converted into sugars and may vary depending on bunch harvest season.
 244 According to [23], banana is a fruit with high amide content when unripe, and as it
 245 ripens, amide is broken into sugars for it to be used in the fruit’s respiration, raising the
 246 content of soluble solids. [17], while studying physical and chemical characteristics of
 247 different banana tree fruits, observed differences between AAB group fruits; unripe
 248 bananas ‘Prata-Anã’ showed a value of 29.68%, close to that found in fruits with bunch
 249 age of 16 weeks and superior to the other ages.




250

251 **Figure 8:** Amide content in bananas 'Prata-Anã' at different bunch ages against days after storage

252


253 For total sugar, significant difference was found between bunch ages and
 254 assessment days. Total sugar percentages were 6.60%, 7.95%, 9.29%, 10.64% and
 255 11.98% for the ages of 16, 17, 18, 19 and 20 weeks, respectively (Figure 9). Bunch
 256 harvest age had a significant influence on amide and sugar content during storage.
 257 Concerning different assessment seasons, on days 1, 3, 5, 7 and 9 days after storage,
 258 there is a linear increase in sugar percentage, with values at 1.61%, 5.45%, 9.29%,
 259 13.14% and 16.98% (Figure 10). According to [17], while amide is hydrolyzed, there is
 260 an increase in total sugar content, which makes fruits ripe and sweet. The main sugars
 261 found in ripe banana pulp are glucose, fructose and sucrose.



262

263 **Figure 9:** Total sugar in bananas 'Prata-Anã' harvested at different bunch ages

264




265

266 **Figure 10:** Total sugar in bananas 'Prata-Anã' on days after storage

267

268 Significant results were observed in interaction for the reducing sugar variable.
 269 As of the first assessment, it is possible to observe a slight percentage increase in
 270 reducing sugar for all treatments; however, concerning fruits from bunches with 16
 271 weeks, they reached a lower value than the other treatments – 9.62%. Treatment with the
 272 ages of 17, 18, 19 and 20 weeks presented higher values – 10.13%; 10.63%; 11.15%
 273 and 11.65%, respectively –; age increase resulted in sugar increase, and lower sugar
 274 percentage in the fruits was found in 16-week bunches (Figure 11). [24], while working

275 with climatization of banana 'Prata-Anã', also found increases in reducing sugar content
276 during ripening, arguing that such increase was due to insoluble
277 moleculeinterconversion, such as non-reducing sugars into depolymerized sugars and
278 then soluble sugars.




279

280 **Figure 11:** Variation of reducing sugar content in bananas 'Prata-Anã' harvested at different bunch ages
281 against days after storage

282

283 For non-reducing sugar, significant difference was found between assessment
284 periods, that is, 1, 3, 5, 7 and 9 days after storage, with values being 1.02%, 2.27%,
285 3.52%, 4.77% and 6.02%, respectively (Figure 12). According to [25], while working
286 with 10 banana genotypes (Pacovan, PV 04-44, PV 03-76, 'Prata-Anã', 'Fhia-18',
287 'Pioneira', 'Prata graúda', 'Caipira', 'Nanica', 'Thap maeo'), found non-reducing sugar
288 values for banana 'Prata-Anã' of $1.3 \pm 0.21\%$, which are lower than those found in the
289 present study.

290




291

292

293 **Figure 12:** Variation of non-reducing sugar content on days after storage

294


295 The fruits' electrolyte extravasation percentage had significant difference
 296 between bunch ages and assessment days. There was an increase on the days after
 297 storage, reaching values close to 37% on the 9th assessment day (Figure 13). Thus, fruits
 298 at a more advanced maturation stage tend to lose membrane integrity and have a faster
 299 electrolyte extravasation compared to those at an earlier maturation stage [26].
 300 Significant difference was found between treatments; although variation between some
 301 treatments was small, fruits from 16-week bunches presented lower electrolyte
 302 extravasation percentage compared to the other treatments, with a value of 17.72%
 303 (Figure 14). As shown by the results of this experiment, the fruits' greater resistance is
 304 due to lower cell membrane degradation, estimated by electrolyte extravasation
 305 percentage.



306

307 **Figure 13:**Electrolyte extravasation of bananas ‘Prata-Anã’ against bunch age

308



309

310 **Figure 14:**Electrolyte extravasation of bananas ‘Prata-Anã’ against days after storage

311

312 **4. CONCLUSIONS**

313 Bunch harvest age had a direct influence on post-harvest quality of bananas ‘Prata-
314 Anã’.

315 Fruits from 16-week bunches were superior in physical and chemical characteristics
316 compared to other ages, meaning a longer post-harvest life.

317

318 **5. REFERENCES**

319

320 1. Severo, F. E. ; Matos, M. C. P. ; Clauzet, M. (2017) Banana production in the coastal
321 region of São Paulo: the discourse of sustainability versus the management of solid
322 residues of agrochemicals. *UnisantaBioScience*, 5 (5), 395-407. Retrieved from:
323 <http://wiki.unisanta.br/index.php/bio/article/view/783>

324 2. Faostat - Food and Agriculture Organization of the United Nations. *Crops* (2008).
325 Retract: <http://www.fao.org/faostat/en/#data/QC>. (2008, March 23).

326 3. Ribeiro, D. M (2006). Evolution of Physical and Chemical Properties during the
327 Ripening of Banana "Prata-Anã. (Thesis Doctorado, Federal University of Viçosa,
328 Minas Gerais, Brazil). Retrieved from: <http://locus.ufv.br/handle/123456789/781>

329 4. Fortescue, J.A., & Turner, D. W. (2005). The association between low temperatures
330 and anatomical changes in preanthetic ovules of *Musa* (Musaceae). *Scientia*
331 *horticulturae*, 104 (4), 433-444. <https://doi.org/10.1016/j.scienta.2005.01.005>Get rights
332 and content

333 5. Adolfo Lutz Institute (2008). *Physical Chemistry Method for Food Analysis*. 6ª Ed,
334 1st Digital Edition, São Paulo, p.1020. Retrieved from:
335 http://www.ial.sp.gov.br/resources/editorinplace/ial/2016_3_19/analisedealimentosial_2
336 008.pdf

337 6. Nelson, N. (1944). A photometric adaptation of the Somogyi method for the
338 determination of glucose. *J. biol. Chem*, 153(2), 375-380. Retirado de:
339 <https://pdfs.semanticscholar.org/131b/e395d4d24726d051472489140cd543aa839e.pdf>

340 7. Dische, Z. (1955). New color reactions for determination of sugars in
341 polysaccharides. *Methods of biochemical analysis*, 313-358.
342 <https://doi.org/10.1002/9780470110188.ch11>

343 8. Whilton, T. H.; Bassuk, N. L.; Ranney, T. G.; Reichert, D. L. (1992). An improved
344 method for using electrolyte leakage to assess membrane competence in plant tissues.
345 *Plant physiology*, 98(1), 198-205. <https://doi.org/10.1104/pp.98.1.198>

346 9. Burr, I. W., & Foster, L. A. (1972). A test for equality of variances. *West Lafayette:*
347 *University of Purdue*, 26.

- 348 10. Shapiro, S. S., & Wilk, M. B. (1965). An analysis of variance test for normality
349 (complete samples). *Biometrika*, 52(3/4), 591-611. DOI: 10.2307/2333709
350 <https://www.jstor.org/stable/2333709>
- 351 11. Tukey, J. W. (1949). One degree of freedom for non-additivity. *Biometrics*, 5(3),
352 232-242. DOI: 10.2307/3001938 <https://www.jstor.org/stable/3001938>
- 353 12. Castricini, A., Dias, M. S. C., & Rodrigues, M. G. V. (2016). Post-harvest of banana
354 'Tropical BRS' in the function of technical management. *Caderno de Ciências Agrárias*,
355 8 (3), 52-62. Retrieved from
356 <https://seer.ufmg.br/index.php/ccaufmg/article/view/3123/2802>
- 357 13. Vilas Boas, E. V. B. ; Rodrigues, L. J. ; De Paula, N. R. F. (2003). Physical,
358 chemical-chemical and chemical modifications of the banana 'Maçã' during maturation.
359 *Latin American Symposium on Food Science and Technology*, 5.
- 360 14. Silva, E. A. ; Boliani, A.C. ; Souza Corrêa, L.S. (2006). Evaluation of banana
361 cultivars (*Musa* sp) in the region of Selvíria-MS. *Revista Brasileira de Fruticultura*, 101-
362 103. <http://dx.doi.org/10.1590/S0100-29452006000100028>
- 363 15. Martins, R. N., Dias, M. S. C., Boas, V., de Barros, E. V., & Santos, L. O. (2007).
364 Banana-stored store 'Ananas' comes from bunches at 16, 18 and 20 weeks. *Science and*
365 *Agrotechnology*, 1423-1429. <http://dx.doi.org/10.1590/S1413-70542007000500023>
- 366 16. Birth Junior, B. B. do; Ozorio, L. P. ; Rezende, C. M., Soares, A. G. ; Fonseca, M. J.
367 de O. (2008). Differences between bananas of cultivars Prata and Nanicãoalongo of
368 ripening: physical-chemical characteristics and volatile compounds. *Food Science and*
369 *Technology*, 28 (3) Retrieved from:
370 <http://www.redalyc.org/articulo.oa?id=39594008802>
- 371 17. Silva, M.J. R. da; Jesus, Fr. R. R. de; Angels, J. M. C. dos; Machado, M., Ribeiro,
372 V. G. (2016). Adrenal and post-harvest characterization of 'Maravilha'e' Preciosa'
373 bananas in the São Francisco Valley. *Ceres*, 63 (1) .[http://doc.doi.org/10.1590/0034-](http://doc.doi.org/10.1590/0034-737X201663010007)
374 [737X201663010007](http://doc.doi.org/10.1590/0034-737X201663010007)
- 375 18. Souza, M. E. de; Leonel, S. ; Martins, R. L. ; Segtowick, E. C. dos S. (2013).
376 Physical-chemical characterization and sensorial evaluation of banana fruits. *Native*, 13-
377 17. <http://dx.doi.org/10.14583/2318-7670.v01n01a03>
- 378 19. Siqueira, M. de S. B. 2014. Characterization of the quality of silver-banana stored in
379 different atmospheric controlled conditions. (Master's Dissertation, Universidade

380 Estadual do Norte Fluminense Darcy Ribeiro, Rio de Janeiro, Brazil).
381 <http://locus.ufv.br/handle/123456789/781>

382 20. Silva, M.J. R. da; Angels, J. M. C. dos; Jesus, Fr. R. R. de; Santos, G. S. dos; Lima,
383 F. B. F. ; Ribeiro, V. G. (2015). Production and characterization of banana 'PrataAnã'
384 (AAB) in production cycles (Juazeiro, Bahia). *Ceres*, 60 (1).
385 <http://dx.doi.org/10.1590/S0034-737X2013000100017>

386 21. Aquino, C.F., Solomon, C.C., C.Cecon, P. R., Siqueira, D.L., Ribeiro, S., & Rocha,
387 M. (2017). Physical, chemical and morphological characteristics of banana cultivars
388 depending on maturation stages. *Revista Caatinga*, 30 (1), 87-96.<http://dx.doi.org/10.1590/1983-21252017v30n110rc>

390 22. Sakyi-Dawson, E., Asamoah-Bonti, P., & Amponsah Annor, G. (2008).
391 Biochemical changes in new plantain and cooking banana hybrids at various stages of
392 ripening. *Journal of the Science of Food and Agriculture*, 88 (15), 2724-2729
393 <https://doi.org/10.1002/jsfa.3399>

394 23. Pimentel, R. D. A. ; Guimarães, F. N. ; Santos, V. D. ; Resende, J. D. (2010). Post-
395 harvest quality of banana genotypes PA42-44 and Prata-Anãcultivados in northern
396 Minas Gerais. *Journal of Fruit Research*, 32 (2), 407-412.

397 24. Prill, M.A. de S. ; Neves, L. C. ; Silva, S. ; Grigio, M.L., Chagas, E. A. ; Campos,
398 A.J. (2011). Banana-based 'Silver-Banana' banana: methods and times for greening after
399 refrigerated storage. *Journal of On-line Mentoring*, 5 (2), 134-142.
400 <http://dx.doi.org/10.18227/1982-8470ragro.v5i2.500>

401 25. Jesus, S. C. D., Matsuura, F. C. A. U., Cardoso, R. L., & Folegatti, M. I. D. S.
402 (2004). Physical and chemical characterization of fruits of different genotypes of banana
403 tree. *EmbrapaMeioAmbiente-Elaborated newspaper article (ALICE)* .<http://dx.doi.org/10.1590/S0006-87052004000300001>

405 26. Nolasco, C.D.A., Solomon, C.C., C.Cecon, P. R., Bruckner, C.H., & Rocha, A.
406 (2008). Postharvest quality of 'Prata' banana as affected by hot water treatment. *Science*
407 *and Agrotechnology*, 32 (5), 1575-1581. [http://dx.doi.org/10.1590/S1413-](http://dx.doi.org/10.1590/S1413-70542008000500033)
408 [70542008000500033](http://dx.doi.org/10.1590/S1413-70542008000500033)

409