

2
3 **Impacts of meteorological attributes on**
4 **agronomic characteristics of sunflower**
5 **cultivated in the Cerrado**

6
7
8
9
10
11 **ABSTRACT**
12

The sunflower has adaptation for cultivation in Brazilian conditions and the main product is the oil extracted from the seeds. This study aimed to verify the impacts of meteorological attributes on agronomic characteristics of sunflower genotypes cultivated as a second crop, in MatoGrosso. The experimental design was random blocks, cultivating the genotypes M734 and Hélio 358, in three years. Were evaluate the agronomic characteristics: plant height, capitulum diameter, weight of a thousand achenes, yield of achenes, oil content and oil yield. Then, water deficit and the maximum crop yield are estimated for correlation analysis between meteorological attributes and agronomic characteristics. In both genotypes studied there was a very strong correlation between plant height with evapotranspiration ($r=0.99^*$) and very strong negative correlation ($r=-0.99^*$) with water deficit. For mass of a thousand achenes, there was a very strong negative correlation with the maximum ($r=-0.98^*$) and minimum ($r=-0.96^*$) temperatures, and with evapotranspiration ($r=-0.98^*$) in the Hélio 358. In addition, effect of the water deficit on the oil yield ($r=-0.98^*$) was verified for the same genotype. Very strong negative correlation was found between maximum temperature and oil content ($r=-0.96^*$) in genotype M734. In sunflower cultivation in MatoGrosso, water deficit reduces plant height, capitulum diameter and oil yield. The oil content of the achenes reduces when the maximum temperature increases, during the cultivation.

13
14 *Keywords: Climatic factors; Correlation; Helianthus annuus L.; Yield of achenes.*
15

16 **1. INTRODUCTION**
17

18 Sunflower is an efficient oilseed in nutrient cycling, drought tolerant and adapts to different
19 soil and climate conditions. Due to these characteristics, the diversity of use and the growing
20 demand of the commercial sector for sunflower products, an increase in cultivated area is
21 expected, especially in the central of Brazil [1, 2]. In this region it is common to seed a
22 second crop in February or March, once the conditions are favorable to a second crop.

23 Due to the low photoperiod sensitivity of the sunflower plant, the cultivation period in Brazil
24 can be carried out all year round. However, despite being a rustic plant, the sunflower has
25 low efficiency of water use and the occurrence of high air temperatures in the periods of
26 flowering, filling of achenes and harvesting has been one of the major aspects that can
27 determine the success of the production [3, 4].

28 In this context, there is the importance of solar radiation in the stage of filling achenes, the
29 availability of water throughout the crop cycle, as well as the air temperature in the duration

30 of development stages and total cycle of the culture [5, 6]. The temperature during sunflower
31 development in MatoGrosso is quite different between the two sowing periods used in the
32 region (crop and second crop), which implies in the occurrence of anthesis at different times
33 [7]. In a study with sunflower, it was observed that the diameter of chapters, the mass of
34 achenes and the oil content of the achenes are affected by the availability of water [8].

35 Environmental conditions may also alter the oil content and fatty acid profile of sunflower
36 achenes[7, 9]. In addition, the genotype x environment interaction, which may differ in the
37 plant performance according to the cultivation region and between different cultivars [10, 11].

38 Due to these factors, the meteorological conditions that occur in the period of second crop in
39 the sunflower cultivation in the Cerrado can cause substantial differences in the
40 morphological characteristics, the yield and the oil content of achenes. Through the
41 determination of the main meteorological attributes that affect the productive characteristics
42 of the sunflower cultivated in the region, it becomes possible to plan management strategies
43 that reduce the losses caused by environmental conditions. Thus, it was aimed to verify the
44 impacts of meteorological attributes on agronomic characteristics of sunflower genotypes
45 cultivated as a second crop, in the State of MatoGrosso.
46

47 **2. MATERIAL AND METHODS**

48
49 The experiments were carried out at Santa Luzia Farm, in Campo Verde, MatoGrosso
50 (latitude 15°45'12"S e longitude 55°22'44"W). The soil of the experimental area is classified
51 as Red-Yellow Latosol, where soybeans preceded the sunflower.

52 The experimental design was random blocks with four replicates. It was evaluated different
53 sunflower genotypes, being selected for the present research the hybrids M734 and Hélio
54 358, cultivated in the years 2009, 2011 and 2012. The plots were constituted by four rows of
55 6.00 m in the spacing of 0.90 x 0.25 m and, having as a useful area, the two central lines,
56 eliminating 0.5 m from the margins.

57 In the three years, the sowing was carried out at the beginning of March by manually placing
58 three seeds per hill. Seven to ten days after the emergency the thinning was done leaving in
59 each hole the most vigorous plant. The fertilizer used was 30-80-80 kg ha⁻¹ of NPK and 2.0
60 kg ha⁻¹ of boron on the sowing hill and 30 kg ha⁻¹ of N top-dressing, at 30 days after sowing.

61 The crop area was kept free from weed interference and the necessary phytosanitary
62 treatments were carried out. At the beginning of flowering, the capitulum diameter and plant
63 height measurements of ten random plants of the area of each plot were made, and then the
64 mean values for these characteristics were calculated. The capitulum diameter was obtained
65 by measuring from one edge to another in the center of the capitulum and the plant height
66 from the insertion of the capitulum on the stem to ground level. When the plants were in
67 stage R7 (first phase of development of achenes), the capitulum were covered with bags to
68 avoid damages by birds.

69 After harvesting, manual threshing of the capitulum was performed and the impurities were
70 separated. Then, the mass of one thousand achenes was determined by weighing eight
71 replicates of 100 achenes, in analytical balance. The yield of achenes was obtained by
72 weighing all the achenes harvested from the useful area of each plot, with subsequent
73 moisture correction to 11%.

74 From each plot a sample of approximately 200 g of achenes was separated to determine the
 75 oil content. After this, the oil yield was calculated by multiplying the oil content by the
 76 achenes yield.

77 The calculation of climatological water balance was performed according to the method
 78 proposed by the authors cited in the reference [12], with precipitation and temperature data
 79 grouped in in periods of ten days. The precipitation values were collected at the experiment
 80 installation site and the temperature (minimum, maximum and mean compensated) obtained
 81 from the Poxoréo Conventional Surface Weather Observation Station, located in Poxoréo-
 82 MT, near the experiment site, based on the National Institute of Meteorology (INMET)
 83 database. The maximum yield was estimated using the Agroecological Zone Method (FAO)
 84 described by the authors cited in the reference [13], adopting a leaf area index for the crop of
 85 15, according to the author cited in the reference [14].

86 After obtaining the agronomic characteristics of the sunflower (Table 1) and the
 87 meteorological attributes (Table 2), Pearson correlation analyzes were performed. The
 88 interpretation was made according to the author cited in the reference [15]: $r = 0.10$ to 0.19
 89 (very weak); $r = 0.20$ to 0.39 (weak); $r = 0.40$ to 0.69 (moderate); $r = 0.70$ to 0.89 (strong); $r =$
 90 0.90 to 1 (very strong), considering the significance level of 5%.

91
92
93
94
95

Table 1. Averages of plant height (PH), capitulum diameter (CD), mass of a thousand achenes (MTA), oil content (OC), oil yield (OY) and achenes yield (AY) of sunflower genotypes grown in MatoGrosso, in 2009, 2011 and 2012

Year	Genotype	PH (cm)	DC (cm)	MTA (g)	OC (%)	OY (kg ha ⁻¹)	AY (kg ha ⁻¹)
2009	M734	138	17	70	38.0	1089	2854
	Hélio 358	114	17	63	47.0	1069	2270
2011	M734	148	17	70	38.8	1292	3311
	Hélio 358	123	18	54	44.9	1048	2328
2012	M734	178	19	70	39.1	814	2082
	Hélio 358	147	18	52	44.0	827	1878

96
97
98
99
100
101

Table 2. Real estimated yield (REY), maximum temperature (T_{MAX}), average temperature (T_{MED}), minimum temperature (T_{MIN}), water deficit (DEF), crop evapotranspiration (ETC) and real evapotranspiration (ETR) of sunflower genotypes grown in MatoGrosso, in 2009, 2011 and 2012

Year	REY (kg ha ⁻¹)	T _{MAX} (°C)	T _{MED} (°C)	T _{MIN} (°C)	DEF (mm)	ETC (mm)	ETR (mm)
2009	1473	31.59	23.38	18.07	204.52	378.53	174.00
2011	1910	32.03	24.05	18.41	159.44	389.29	229.86
2012	2753	31.98	23.94	18.58	18.12	369.17	351.05

Temperature data: Poxoréo Conventional Surface Weather Observation Station (2009, 2011 and 2012).

102
103
104
105

106 3. RESULTS AND DISCUSSION

107
108
109
110
111

For plant height, there was a very strong positive correlation with real evapotranspiration ($r = 0.99^*$) and very strong negative correlation ($r = -0.99^*$) with water deficit (Tables 3 and 4), corroborating with results obtained by the authors cited in the reference [16], who verified that the increase in soil water level provides an increase in height of the sunflower plant.

112
113

Table 3. Correlation coefficient (r) between meteorological attributes and agronomic characteristics of the genotype M734 cultivated in MatoGrosso

	PH	CD	MTA	OC	OY	MRY	REY
T_{MAX}	0.61	0.40	-0.80	-0.96*	0.01	-0.04	0.68
T_{MED}	0.57	0.36	-0.77	-0.91	0.06	0.01	0.65
T_{MIN}	0.89	0.76	-0.98	-0.97	-0.41	-0.46	0.93
DEF	-0.99*	-0.97*	0.95	-0.85	-0.78	-0.81	-0.99*
ETC	-0.69	-0.84	0.46	-0.22	0.99	0.98	-0.61
ETR	0.99*	0.95*	-0.97	0.89	-0.97*	-0.77	0.99*

114
115
116
117
118
119
120

T_{MAX} : maximum temperature (°C), T_{MED} : average temperature (°C), T_{MIN} : minimum temperature (°C), DEF: water deficit (mm), ETC: crop evapotranspiration (mm day⁻¹), ETR: real evapotranspiration (mm day⁻¹), PH: plant height (cm), CD: capitulum diameter (cm), MTA: mass of a thousand achenes (g), OC: oil content (%), OY: oil yield (kg ha⁻¹), MRY: measured real yield (kg ha⁻¹), REY: real estimated yield (kg ha⁻¹).

*significant at 5% probability.

121
122
123

Table 4. Correlation coefficient (r) between meteorological attributes and agronomic characteristics of the genotype Hélio 358 cultivated in MatoGrosso

	HP	CD	MTA	OC	OY	MRY	REY
T_{MAX}	0.63	0.99*	-0.98*	-0.92	-0.47	-0.29	0.68
T_{MED}	0.59	0.98*	-0.94	-0.90	-0.43	-0.24	0.65
T_{MIN}	0.90	0.94	-0.96*	-0.99	-0.80	-0.67	0.93
DEF	-0.99*	-0.68	0.80	0.86	-0.98*	-0.93	-0.99*
ETC	-0.67	0.04	0.13	0.25	0.80	0.90	-0.61
ETR	0.99*	0.74	-0.98*	-0.90	-0.97*	-0.90	0.99*

124
125
126
127
128
129
130

T_{MAX} : maximum temperature (°C), T_{MED} : average temperature (°C), T_{MIN} : minimum temperature (°C), DEF: water deficit (mm), ETC: crop evapotranspiration (mm day⁻¹), ETR: real evapotranspiration (mm day⁻¹), PH: plant height (cm), CD: capitulum diameter (cm), MTA: mass of a thousand achenes (g), OC: oil content (%), OY: oil yield (kg ha⁻¹), MRY: measured real yield (kg ha⁻¹), REY: real estimated yield (kg ha⁻¹).

*significant at 5% probability.

131 Plant height reflects the nutritional conditions in the stem elongation period, and the
132 response of the cultivars can show the efficiency of the plant to the edaphoclimatic
133 conditions of cultivation [17], including water availability. In low water conditions, the
134 decrease in height of sunflower plants possibly occur due to the increase of abscisic acid
135 level that slows plant growth, the stomatal closure as a mechanism to reduce water loss, as
136 well as by changes in hormonal activity [18].

137 For the M734 genotype, there was a very strong positive correlation of the capitulum
138 diameter with the real evapotranspiration ($r = 0.95^*$) and very strong negative correlation ($r =$
139 -0.97^*) with the water deficit (Table 3), agreeing with results obtained by the authors cited in
140 the reference [8] who verified greater capitulum diameter with the increase of water
141 availability.

142 Water deficit in sunflower may cause floral abortion and reduction of the capitulum diameter
143 resulting in decrease of achenes production [3]. The influence of water deficit on the
144 capitulum diameter was observed in Hélio 358 (Table 4), a result that reinforces that
145 tolerance to low water availability depends not only on the species, but also to the cultivar
146 [19].

147 In the Hélio 358, there was a very strong positive correlation of the capitulum diameter with
148 the maximum ($r = 0.99^*$) and mean ($r = 0.98^*$) temperature (Table 4). For sunflower, low
149 temperatures generate an increase in the plant cycle, causing losses in productivity,
150 especially when it occurs at the beginning of flowering [20], which is related to the size of the
151 capitulum to be formed. The capitulum diameter depends to the environmental conditions,
152 management and genotype. This characteristic is relevant because capitulum with larger
153 diameters provide higher yields of achenes[21].

154 For mass of a thousand achenes, there was a very strong negative correlation with the
155 maximum ($r = -0.98^*$) and minimum ($r = -0.96^*$) temperatures, and with evapotranspiration (r
156 $= -0.98^*$) in the Hélio 358 (Table 4). In sunflower, the increase in temperature decreases the
157 accumulation of intercepted radiation, which may cause losses in mass of achenes[9].

158 Similarly, reductions in the production of dry mass of achenes and oil content occur due to
159 the lower availability of water at the beginning of flowering or in the filling of achenes, which
160 reflects the climatic conditions in the second crop in the region [22, 23]. The translocation of
161 assimilates in the plant is very related to evapotranspiration, and the need of water increases
162 with the development of the plant. In sunflower, this demand generally ranges from 0.5 to 10
163 mm per day depending on the phenological stage [20].

164 There was a very strong negative correlation of the maximum temperature with the oil
165 content ($r = -0.96^*$) in the M734 (Table 3). Generally, raising the maximum temperature
166 causes damage to the production of oil in the plant. The authors cited in the reference [24]
167 studying the influence of air temperature on the oil content of sunflower seeds found that
168 there was a decrease in the total content at higher temperature.

169 In addition to the loss of oil content, the temperature affects the fatty acid composition of
170 sunflower oil [7]. According to the authors cited in the reference [20], a mild temperature
171 during the synthesis of oil benefits the concentration of linoleic acid, while the occurrence of
172 temperature above 35°C between flowering and physiological maturation causes the
173 irreversible increase of the content of oleic acid.

174 For oil yield, there was a very strong negative correlation between real evapotranspiration in
175 the genotypes M734 and Hélio 358 ($r = -0.97^*$), and with the water deficit in Hélio 358 ($r = -$
176 0.98^*) (Tables 3 and 4). The measured yield of achenes showed no correlation with the
177 meteorological attributes, but for estimated yield there was a very strong positive correlation
178 with real evapotranspiration ($r = 0.99^*$) and very strong negative correlation ($r = -0.99^*$) with
179 water deficit in both genotypes.

180 In cultivation, the productivity and the quality are strongly influenced by genotype,
181 environment and their interaction [25]. The water deficit may prolong the total sunflower
182 growth period [26], increasing the risk of losses in the final yield of the crop. The lack of
183 water in the vegetal tissues implies less leaf expansion, as a mechanism to minimize the
184 loss by evapotranspiration. When water stress occurs in later stages of plant development,
185 the reduction of leaf expansion in the anthesis period causes a lower yield of achenes, in
186 addition to damage the absorption of important nutrients to the plant, such as boron [20].

187 In sunflower, the authors cited in the references [27, 28]found higher yields of achenes due
188 to the increase of water availability through irrigation. In the culture, the most sensitive
189 development phases to water deficiency are from the beginning of capitulum formation to the
190 beginning of flowering and the formation and filling of achenes. About 500 mm of water, well
191 distributed throughout the cycle, results in yields close to the maximum potential for the crop
192 [8, 29].

193

194 4. CONCLUSION

195

196 The water deficit decreases plant height, capitulum diameter and yield of oil in sunflower
197 cultivated as a second crop in MatoGrosso.

198

199 Increases in maximum temperature during sunflower cultivation reduce the oil content of
achenes in the region studied.

200

201 The impacts of meteorological attributes on some agronomic characteristics of sunflower
depend on the genotype.

202

203

204

205 REFERENCES

206

207 1. Grunvald AK, Carvalho CGP, Oliveira ACB, Andrade CAB. Adaptabilidade e estabilidade
208 de genótipos de girassol no Brasil Central.
209 Pesquisa Agropecuária Brasileira. 2008;43(11):1483-1493. Portuguese. DOI: 10.1590/S0100-
210 204X2008001100006

211

212 2. Souza FR, Silva IM, Pellin DMP, Bergamin AC, Silva RP. Características agrônomicas do
213 cultivo de girassol consorciado com *Brachiaria ruziziensis*.
214 Revista Ciência Agronômica. 2015;46(1):110-116. Portuguese. DOI: 10.1590/S1806-
215 66902015000100013

216

217 3. Aleman CC, Bertipaglia R. Influência da lâmina de irrigação no cultivo de girassol.
218 Colloquium Agrariae. 2015;11(2):24-30. Portuguese. DOI: 10.5747/ca.2015.v11.n2.a123

219

220 4. Leite RMVBC, Castro C, Brighenti AM, Oliveira FA, Carvalho CGP, Oliveira ACB.
221 Indicações para o cultivo de girassol nos Estados do Rio Grande do Sul, Paraná, Mato
222 Grosso do Sul, Mato Grosso, Goiás e Roraima. Comunicado Técnico 78, Embrapa
223 Soja. Londrina. 2007;1-4. Portuguese. Accessed 02 Feb 2019.

224

225 Available: <https://ainfo.cnptia.embrapa.br/digital/bitstream/item/18674/1/comtec78.pdf>

226

227 5. Izquierdo NG, Dosio GAA, Cantarero M, Luján J, Aguirrezábal LAN. Weight per grain, oil
228 concentration, and solar radiation intercepted during grain filling in black hull and striped hull
229 sunflower hybrids. Crop Science. 2008;48(2):688-699. DOI: 10.2135/cropsci2007.06.0339

230

231 6. Thomaz GL, Zagonel J, Colasante LO, Nogueira RR. Produção do girassol e teor de óleo
232 nos aquênios em função da temperatura do ar, precipitação pluvial e radiação solar. Ciência
233 Rural. 2012;42(8):1380-1385. Portuguese.

233

234 7. Regitano Neto A., Miguel AMRO, Mourad AL, Henriques EA, Alves RMV. Efeito da
235 Temperatura no Perfil de Ácidos Graxos do Óleo de Girassol. Documentos 262, Embrapa
236 Semiárido. Petrolina. 2015;1-7. Portuguese. Accessed 15 Feb 2019.

237

238 Available: <https://www.alice.cnptia.embrapa.br/alice/bitstream/doc/1017979/1/691.pdf>

239

240 8. Silva ARA, Bezerra FML, Sousa CCM, Pereira Filho JV, Freitas CAS. Desempenho de
241 cultivares de girassol sob diferentes lâminas de irrigação no Vale do Curu, CE.
Revista Ciência Agronômica. 2011;42(1):57-64. Portuguese. DOI: 10.1590/S1806-
66902011000100008

- 242 9. Aguirrezábal LAN, Lavaud Y, Dosio AA, Izquierdo NG, Andrade FH, González LM.
243 Intercepted Solar Radiation during Seed Filling Determines Sunflower Weight per Seed and
244 Oil Concentration. *Crop Science*.2003;43(1):152-161. DOI: 10.2135/cropsci2003.0152
- 245 10. Gomes AHS, Chaves LHG, Guerra HOC. Drip irrigated sunflower intercropping.
246 *American Journal of Plant Sciences*.2015;6(11):1816-1821.DOI: 10.4236/ajps.2015.611182
- 247 11. Porto WS, Carvalho CGP, Pinto RJB, Oliveira MF, Barneche ACO. Evaluation of
248 sunflower cultivars for central Brazil.*Scientia Agricola*. 2008;65(2):139-144. DOI:
249 10.1590/S0103-90162008000200005
- 250 12. Thornthwaite CW, Mather JR. The water balance.Centerton: Drexel Institute of
251 Technology; 1955.
- 252 13. Doorenbos J,Kassam AH. Efeito da água no rendimento das culturas. Campina Grande:
253 UFPB; 1994.Portuguese.
- 254 14. Carvalho DB. Análise de crescimento de girassol em sistema de semeadura direta.
255 *Revista Acadêmica: Ciências Agrárias e Ambientais*. 2004;2(4):63-70.Portuguese. DOI:
256 10.7213/cienciaanimal.v2i4.15135
- 257 15. Shimakura SE. Correlação. In: CE003 - Estatística II. Paraná: Dep. de Estatística da
258 Universidade Federal do Paraná; 2006.Portuguese.
- 259 16. Paiva Sobrinho S,Tieppo RC, Silva TJA. Desenvolvimento inicial de plantas de girassol
260 em condições de estresse hídrico. *Enciclopédia Biosfera*. 2011;7(12):1-12.Portuguese.
- 261 17. Ivanoff MEA, Uchôa SCP, Alves JMA, Smiderle OJ, Sedyama T. Formas de aplicação
262 de nitrogênio em três cultivares de girassol na savana de Roraima. *Revista Ciência*
263 *Agronômica*. 2010;41(3):319-325.Portuguese. DOI: 10.1590/S1806-66902010000300001
- 264 18. Dutra CC, Prado EAF, Paim LR, Scalon SPQ. Desenvolvimento de plantas de girassol
265 sob diferentes condições de fornecimento de água. *Semina: Ciências Agrárias*.
266 2012;33(1):2657-2668.Portuguese.DOI: 10.5433/1679-0359.2012v33Supl1p2657
- 267 19. Bezerra FTC, Dutra AS, Bezerra MAF, Oliveira FILHO AFD, Barros GDL.
268 Comportamento vegetativo e produtividade de girassol em função do arranjo espacial das
269 plantas. *Revista Ciência Agronômica*. 2014;45(2):335-343.Portuguese. DOI:
270 10.1590/S1806-66902014000200015
- 271 20. Gazzolaet al. A cultura do girassol. Piracicaba: Escola Superior de Agricultura Luiz de
272 Queiroz; 2012.
- 273 21. Amorim EP, Ramos NP, Ungaro MRG, Kiihl TAM. Correlações e análise de trilha em
274 girassol. *Bragantia*. 2008;67(2):307-316.Portuguese. DOI: 10.1590/S0006-
275 87052008000200006
- 276 22. Castro C, Moreira A, Oliveira RF, Dechen AR. Boro e estresse hídrico na produção do
277 girassol. *Ciência e Agrotecnologia*. 2006;30(2):214-220.Portuguese. DOI: 10.1590/S1413-
278 70542006000200004

- 279 23. Silva AG, Pires R, Morães EB, Oliveira ACB, Carvalho CGP. Desempenho de híbridos
280 de girassol em espaçamentos reduzidos. *Semina: Ciências Agrárias*. 2009;30(1):31-
281 38. Portuguese. DOI: 10.5433/1679-0359.2009v30n1p31
- 282 24. Harris HC, McWilliam JR, Mason NK. Influence of temperature on oil content and
283 composition of sunflower seed. *Australian Journal of Agricultural Research*.
284 1978;29(6):1203-1212. DOI: 10.1071/AR9781203
- 285 25. Denčić S, Mladenov N, Kobiljski B. Effects of genotype and environment on breadmaking
286 quality in wheat. *International Journal of Plant Production*. 2011;5(1):71-82. DOI:
287 10.22069/IJPP.2012.721
- 288 26. Ma T, Zeng W, Li Q, Wu J, Huang J. Effects of water, salt and nitrogen stress on
289 sunflower (*Helianthus annuus* L.) at different growth stages. *Journal of Soil Science and*
290 *Plant Nutrition*. 2016;16(4):1024-1037. DOI: 10.4067/S0718-95162016005000075
- 291 27. Anastasi U, Santonoceto C, Giuffre AM, Sortino O, Abbate V. Yield performance and
292 grain lipid composition of standard and oleic sunflower as affected by water supply. *Field*
293 *Crops Research*. 2010;119(1):145-153. DOI: 10.1016/j.fcr.2010.07.001
- 294 28. Gomes EP, Fedri G, Ávila MR, Biscaro GA, Rezende RKS, Jordan RA. Produtividade de
295 grãos, óleo e massa seca de girassol sob diferentes lâminas de irrigação suplementar.
296 *Revista Brasileira de Engenharia Agrícola e Ambiental*. 2012;16(3):237-246. Portuguese.
297 DOI: 10.1590/S1415-43662012000300001
- 298 29. Castro C, Farias JRB. Ecofisiologia do girassol. In: Leite RMVBC, Brighenti AM, Castro
299 C. *Girassol no Brasil*. Londrina: Embrapa Soja; 2005. Portuguese.