

1 **Original Research Article**

2 **BIOMASS AND STOCK OF NUTRIENTS IN DIFFERENT GENOTYPES OF**

3 ***Eucalyptus* IN SOUTHERN BRAZIL**

4
5
6 **Abstract:** The objective of this study was to estimate the biomass, nutrient stocks, and
7 nutrient utilization efficiency of six genotypes of *Eucalyptus*. The experiment was
8 conducted in Eldorado do Sul, Rio Grande do Sul, Brazil. The selected trees were
9 fractionated into leaves, branches, stembark and stemwood. The amount of total biomass
10 ranged from 68.40 to 117.52 Mg ha⁻¹, with the highest production being *E. urolobulus*,
11 and *E. dunnii* the lowest. The canopy (leaves and branches) accumulated between 17% and
12 52% of the total macronutrients in *E. benthamii* (P1) and *E. urolobulus* and from 24% to
13 34% of the total micronutrients in *E. dunnii* and *E. urolobulus*. While the stem (wood and
14 bark) accumulated between 48 to 83% and 66 to 76% of the total macro and micronutrients,
15 respectively. For the stemwood, it was observed that *E. benthamii* (P2) presented the
16 highest values of nutritional efficiency for N, Ca, Cu and Fe, and *E. urolobulus* for P, Mg
17 and B. The different genotypes of *Eucalyptus*, under the same edaphoclimatic conditions,
18 presented different biomass production. Variations in concentration, in the allocation of the
19 amount of nutrients in the different genotypes, and in the different components of the same
20 genotypes were observed.

21 **Keywords:** *Eucalyptus* productivity, Forestry nutrition, Silviculture, Sustainability.

22

23 **1. INTRODUCTION**

24 *Eucalyptus* silviculture has expanded worldwide, mainly because of the increasing
25 demand for wood and the high potential of the genus for biomass production [23]. In
26 Brazil, the expansion of forestry was boosted by a government policy that subsidized
27 reforestation programs from 1967 to 1989, with the aim of developing an internationally
28 competitive logging industry [7].

29 The possibility of using eucalyptus wood for various purposes led both large and
30 small companies to establish eucalyptus plantations for multiple uses [7]. Currently,
31 eucalyptus plantations occupy 5.6 million hectares of the country's forest plantation area,
32 with an annual growth of 2.8% [11]. This rate of increase has been constant for more than
33 40 years [7], with growth rates strongly dependent on the genetic of clones, forestry
34 practices, and climate [4]. Thus, improving the use efficiency of natural resources through
35 the creation of genotypes and using appropriate practices of site management is a
36 fundamental challenge of maintaining or increasing productivity in a sustainable manner
37 [7].

38 Biomass production varies according to the availability of resources at different sites,
39 mainly through influences in the processes of photosynthesis, respiration,
40 compartmentalization of carbon, underground flow, and leaf production, among others [15].
41 The quantification of forest biomass allows the determination of the production potential,
42 or adequacy, of certain species for specific purposes, and the prediction of crop yields, thus
43 helping to assess the loss or accumulation of biomass over time [12].

44 To define management practices in forest plantations, it is important to choose
45 species that achieve maximum biomass production for a given location by maximizing the
46 uptake of nutrients [9]. For this, the prolongation of the harvest cycle is necessary. In order

47 to achieve maximum efficiency during nutrient cycling, it is important to reduce the
48 unnecessary export of nutrients [17]. In this context, the objectives of future studies on
49 forest biomass should reconsider traditional practices and seek new alternatives to maintain
50 an efficiently balanced crop [5].

51 Studies on the biomass production and the nutrient stocks of different
52 species/provenances, planted under the same edaphoclimatic conditions, are key to select
53 genotypes which are able to achieve high productivity in a sustainable way. Therefore, the
54 objective of the present study was to estimate the biomass, nutrient stocks, and nutrient use
55 efficiency in six different genotypes of *Eucalyptus* established in Eldorado do Sul, Rio
56 Grande do Sul (RS), Brazil.


57

58 **2. METHODS**

59 **2.1 Characterization of the site**

60 The experiment was conducted in Eldorado do Sul, Rio Grande do Sul, southern of
61 Brazil, in the Horto Florestal Terra Dura, owned by Celulose Riograndense – CMPC (30°
62 11'30.3"S and 51° 37'47.7"W). The approximate altitude of the place is 158 m.

63 The climate of the region is characterized as subtropical humid (Cfa), according to
64 the climatic classification of Köppen presenting an average temperature of 19 °C. The
65 average annual precipitation reaches 1,400 mm [2]. In the period from 2012 to 2016,
66 the average rainfall was 1283.6 mm per year. The annual mean temperature was
67 approximately 17.6 ° C (Figure 1). The soil in the experimental area is of the type Red-
68 Yellow Argissol. Table 1 presents the clay and chemical attributes of the soil at depths from
69 0 to 130 cm.



71

72 Figure 1 - Climatic diagram of the municipality of Eldorado do Sul, RS, Brazil, during the
73 study period (2012 to 2016).

74

75 Table 1 – Physical and chemical attributes of the soil of the area implanted with different
76 genotypes of *Eucalyptus* at 49-months-old in Eldorado do Sul, RS, Brazil

77

Prof.	pH	Argila	C.O	V	m	Al	T	N	P
	H ₂ O	-----%-----			--cmol _c dm ⁻³ --		%	mg g ⁻¹	
0-30	5,0	17	0,88	35	34	0,9	10,3	0,10	2,0
30-60	4,3	9	0,77	11	71	3,7	14,0	0,09	1,6

60-90	4,4	25	0,66	15	69	4,8	15,3	0,08	1,0
90-100	4,6	4	0,42	17	64	3,6	12,0	0,06	0,7
100-130	4,7	6	0,22	20	61	3,1	10,0	0,04	0,6
Prof.	K	Ca	Mg	S	B	Zn	Mn	Cu	Fe
	-----cmol _c dm ⁻³ -----			-----mg dm ⁻³ -----				g dm ³	
0-30	0,14	3,3	0,9	19,4	0,4	0,5	13	0,8	0,1
30-60	0,14	0,9	0,5	32,5	0,7	0,5	13	1,2	0,1
60-90	0,15	1,0	0,8	61,7	0,5	0,3	7	1,2	0,1
90-100	0,14	1,0	0,9	60,9	0,3	0,3	5	1,0	0,1
100-130	0,12	0,9	0,9	59,0	0,3	0,3	5	0,7	0,1

78 O.C: organic carbom; V = saturation by base; m = saturation by aluminum; T = total cation

79 exchange capacity.

80

81 2.2 Planting of the experimental area

82 The genotypes were planted in April 2012, with spacing each plant in a plot of 3 m
83 x 3 m. Subsoiling was performed at a depth of 60 cm, using a subsoiler with three stems,
84 and a liming treatment was applied consisting of 2 Mg ha⁻¹ of limestone, and 200 kg ha⁻¹ of
85 single superphosphate. Three different fertilizers were applied under different methods:
86 fertilization during planting, coverage fertilization, and maintenance fertilization. The
87 fertilizer used during planting consisted of, 110 g plant⁻¹ of N-P₂O₅-K₂O (06:30:06) + 0.3%
88 Zn and 0.2% Cu. For coverage fertilization 200 kg ha⁻¹ of N-P₂O₅-K₂O (12:00:20) + 0.7%
89 of B were applied, and for the maintenance fertilization, 300 kg ha⁻¹ of N-P₂O₅-K₂O
90 (24:00:26) + 0,5% B were applied.

91 Before planting, a chemical weeding with 2.5 kg ha⁻¹ of glyphosate was carried out.
 92 After planting chemical weeding was carried out at 120 and 300 days, with 1.7 kg ha⁻¹ of
 93 Scout (glyphosate) at the interrow. Also, it was carried out to combat leaf-cutting ants.

94 The following *Eucalyptus* clones were planted: *E. benthamii* (P1), *E. benthamii* (P2),
 95 *E. saligna*, *E. dunnii*, hybrid of *E. urophylla* × *E. globulus* (*E. uroglobulus*), and hybrid of
 96 *E. urophylla* × *E. grandis* (*E. urograndis*). *E. benthamii* (P1) is a provenance originating
 97 from Guarapuava, Paraná, Brazil and *E. benthamii* (P2) is from Telêmaco Borba, Paraná,
 98 Brazil. At the time of data collection, the stands were 49 months old.

99 For each genotype of eucalyptus, a plot of 720 m² was demarcated, where the DBH
 100 (diameter at breast height, measured at 1.30 m above ground level) of all individuals was
 101 measured with diametrical tape. The heights of 20% of the plants were measured using a
 102 Vertex hypsometer; thus, the heights that were not measured in the field were estimated
 103 through hypsometric models. According to Table 2, the mean volume varied from 73.96 to
 104 114.99 m³ ha (*E. dunnii* and *E. benthamii* (P2)). The highest mortality of trees occurred in
 105 the settlement of *E. dunnii* (21%). In contrast, the hybrid *E. urograndis* had a 100%
 106 survival.

107

108 Table 2 - Dendrometric characterization of different genotypes of *Eucalyptus* at 49-month-
 109 old in Eldorado do Sul, RS, Brazil

110

Genotypes of <i>Eucalyptus</i>	N° of individuals per ha	Basal area (m² ha⁻¹)	Volume (m³ ha⁻¹)
---------------------------------------	-------------------------------------	---	---

<i>E. benthamii</i> (P1)	986 (192)**	24,4ab (8,8)	105,19a (51,1)
<i>E. benthamii</i> (P2)	1.000 (216)	22,7b (6,3)	114,99a (48,0)
<i>E. saligna</i>	972 (206)	23,7ab (3,5)	103,63a (29,8)
<i>E. dunnii</i>	875 (195)	16,7c (6,9)	73,96b (40,62)
<i>E. uroglobulus</i>	903 (183)	22,2b (7,3)	100,27a (43,56)
<i>E. urograndis</i>	1.111 (229)	26,4a (4,9)	111,93a (43,24)

111 Mean of each variable in different treatments (genotypes of *Eucalyptus*) followed by equal
 112 letters, do not differ significantly by the Tukey test at the 5% level of error. * Values in
 113 parentheses are the standard deviation of the mean.

114

115 **2.3 Biomass and nutrient stocks**

116 According to the data obtained in the plot inventory, three trees with a mean diameter
 117 were sampled for each genotype of eucalyptus. The selected trees were felled and separated
 118 in the following components: leaves, branches, stembark and stemwood.

119 A sampling of the wood and bark of the stem was done by dividing the trunk into
 120 three sections of equal parts, with the sampling performed on three points in the median

121 position of each section. All biomass samples were weighed in the field with a precision
122 scale to determine the moisture content. Subsequently they were sent to the laboratory and
123 dried in an oven at 70 °C with circulation and air exchange until weight stabilization. Based
124 on the dry biomass of each component and the number of trees per hectare of each genetic
125 material, the total biomass per hectare was estimated.

126 For nutrient determination, the samples were milled with Wiley-type blades, with 30
127 mesh sieves and submitted to chemical analysis to determine N content by the Kjeldahl
128 method; Ca, Mg, Cu, Fe, Mn and Zn content by atomic absorption spectrometry; P and B
129 content by spectrophotometry; K content by flame photometry, and S content by
130 turbidimetry. The nutrients were analyzed according to the methodology of [24,13]. The
131 estimates of the nutrient stock for each component was obtained by multiplying the dried
132 biomass by the concentration of nutrients. The estimate per hectare was performed by
133 extrapolating the stock per individual based on the number of individuals present in each
134 sampling unit.

135

136 **2.4 Nutrient Use Efficiency (NUE)**

137 The values of nutrient use efficiency (NUE) were obtained by dividing the amount
138 of biomass of each component and the amount of nutrient from each biomass component,
139 according to the equation:

$$140 \quad \text{NUE} = \frac{\text{(Amount of biomass)}}{\text{(Amount of nutrient)}}$$

141

142 **2.5 Statistical procedures**

143 Statistical analyses were performed at a 5% error probability level with the statistical
144 software Assistat 7.7 [21]. The biomass and nutrient concentration data were subjected to
145 analysis of variance and Tukey's test for comparison of means between treatments
146 (genotypes of eucalyptus).

147

148 **3. RESULTS AND DISCUSSION**

149 **3.1 Aboveground biomass**

150 The highest total biomass production was observed in *E. uroglobulus* and the lowest
151 in *E. dunnii*, with 117.52 and 68.40 Mg ha⁻¹, respectively (Table 3). Similar values to the
152 genotype of the present study were reported by [19] while evaluating *E. globulus* in a four-
153 year-old plantation in Butiá, RS, Brazil (83.2 Mg ha⁻¹). Lower values were reported by [20]
154 while evaluating *Eucalyptus* spp. in plantations of two and four years of age in Vera Cruz
155 (RS), Brazil (26.70 and 44.55 Mg ha⁻¹); and by [16], studying *E. saligna* at 1.1 years of age
156 in Telêmaco Borba, Paraná (PR), Brazil (37.35 Mg ha⁻¹). In a study conducted by [28], in
157 the Pearl River Delta region of southern China, when grouping species of eucalyptus into
158 three age classes: < 6 years, 6–15 years, and 16 years of age, the authors found a marked
159 increase in the accumulation of biomass with the increase of age with values of 54.63,
160 136.94, and 186.43 Mg ha⁻¹, respectively. This suggests that the production of biomass is
161 influenced by plant age, species specific characteristics and planting location.

162 In relation to the stemwood biomass, the *E. uroglobulus* hybrid produced 25 and
163 44% more than the *E. saligna* and *E. dunnii* clones, respectively. Genetic factors
164 (improvement and provenance), edaphoclimatic conditions, and management practices are
165 directly related to the production capacity of the species [8].

166 Table 3 - Production and partition of biomass for the different components of genotypes

167 *Eucalyptus* at 49-month-old established in Eldorado do Sul, RS, Brazil

168

Genotypes of <i>Eucalyptus</i>	Leaves	Branches	Stembark	Stemwood	Biomass
	Mg ha ⁻¹				
<i>E. benthamii</i> (P1)	4,36b*	7,04a	8,17ab	73,04b	92,19b
	(4,73)**	(7,64)	(8,86)	(79,23)	(100,00)
<i>E. benthamii</i> (P2)	3,92bc	5,08b	8,60a	84,44ab	102,04ab
	(3,84)	(4,98)	(8,43)	(82,75)	(100,00)
<i>E. saligna</i>	3,22c	5,60b	7,92ab	72,50b	89,25bc
	(3,61)	(6,28)	(8,87)	(81,24)	(100,00)
<i>E. dunnii</i>	3,09c	4,59b	6,05c	54,68c	68,40c
	(4,51)	(6,70)	(8,84)	(79,94)	(100,00)
<i>E. urolobulus</i>	6,52a	7,47a	6,69bc	96,84a	117,52a
	(5,55)	(6,36)	(5,69)	(82,40)	(100,00)
<i>E. urograndis</i>	3,05a	7,41a	7,76ab	83,58ab	101,80ab
	(2,99)	(7,28)	(7,62)	(82,10)	(100,00)

169 *Averages of each fraction of biomass in different treatments (genotypes of *Eucalyptus*)

170 followed by equal letters, do not differ significantly by the Tukey test at the 5% level of

171 error. ** Values in parentheses refer to the percentage of each component in relation to the

172 total biomass of each genotype.

173

174 The greatest contribution to total biomass was from the stemwood, followed by the
175 stembark, branches, and leaves, except in the clone *E. uroglobulus*, from which the greatest
176 contribution to total biomass was from the stembark. The relative distribution of biomass,
177 considering the same components, was the same as that found by: [25] while studying *E.*
178 *urophylla* × *E. globulus* at 10 years of age, in Eldorado do Sul, RS, Brazil; by [16] while
179 evaluating *E. saligna* at 6.7 years of age in Telemaco Borba, PR, Brazil; and by [6] while
180 studying the biomass of eucalyptus plantations of different ages in the Central-Eastern
181 Region of the State of Minas Gerais, Brazil. In plantations of *E. nitens* in northern Spain,
182 the distribution trend in terms of total biomass was wood > bark > thick twigs > dried twigs
183 > leaves > fine twigs > twigs [14].

184 By adding the value of the bark to that of the wood, the biomass of the stem
185 represents from 88 to 91% of total aboveground biomass, whose the lowest value was
186 found in *E. benthamii* (P1) and the highest in *E. benthamii* (P2), while the canopy (leaves
187 and branches) represents 9 to 12% of the total aboveground biomass. Some previous
188 studies have reported contrasting results: [8] while evaluating *E. dunnii* at 4 years of age,
189 reported that 81% of the aerial biomass was found in the wood and bark components; and
190 [19], estimating the biomass of *E. globulus*, also at 4 years of age, reported that 77% of the
191 biomass was found in the same components.

192 In an experiment with *E. saligna* and *E. urophylla* × *E. grandis* at 18 months of age,
193 the authors observed that even at an early age the contribution of the wood component to
194 biomass was the largest relative to total aerial biomass, while the contribution from the bark
195 component was the lowest. The average proportions were 41.5 and 37.4% for the wood and
196 7.5 and 7.1% for the bark, for *E. saligna* and *E. urophylla* × *E. grandis*, respectively [27].

197 Before the closure of the canopy, there is a period of intense growth in which most
 198 of the photoassimilates synthesized by the plant are channeled into the canopy and root
 199 systems. In this phase, the roots partially exploit soil volume and trees do not compete with
 200 each other for growth factors (e.g., light, water, and nutrients). After the crowning of the
 201 tree canopy, the accumulation of nutrients in the trunks occurs with more intensity, as the
 202 formation of the canopy reaches a phase of relative stability, due to auto-shading that
 203 imposes a maximum leaf area limit [20].

204

205 3.2 Concentration of nutrients

206 Nutrient concentrations varied between genotypes and between different
 207 components within the same genotype (Table 4). In general, the leaves had the highest
 208 concentrations of nutrients and the wood the lowest concentrations, while the branches and
 209 bark exhibited intermediate values. The tendency for most nutrients to accumulate in the
 210 leaves is because leaves have a higher metabolic activity than other components of the plant
 211 [25].

212

213 Table 4 - Nutrient concentrations in the different biomass components of genotypes of
 214 *Eucalyptus* at 49-month-old established in Eldorado do Sul, RS, Brazil

215

Genotypes of <i>Eucalyptus</i>	Fractions	N	P	K	Ca	Mg	S	B	Cu	Fe	Mn	Zn
		g kg ⁻¹						mg kg ⁻¹				
<i>E. benthamii</i> (P1)	Leaves	22,83a	1,38a	8,24ab	6,07a	2,85ab	1,35a	19,68bc	4,90a	133,63a	460,59a	16,10*
	Branches	1,91a	0,34a	3,34a	5,58a	1,83*	0,31a	6,20ab	3,72b	51,44a	307,19a	10,77*
	Stembark	5,38a	0,58a	5,92a	15,94a	3,69ab	0,32a	13,97a	3,18a	32,33ab	508,08a	15,82*
	Stemwood	0,89a	0,17a	1,97a	0,57ab	0,28bc	0,18a	2,90*	1,08ab	69,03a	20,51a	4,94*

<i>E. benthamii</i> (P2)	Leaves	23,52a	1,27ab	6,63bc	5,19a	2,72ab	1,28a	24,48ab	5,74a	124,37ab	358,99a	13,13ab
	Branches	1,57a	0,26ab	2,90a	2,90a	1,31ab	0,29a	5,20b	2,84b	45,03a	162,36ab	9,42*
	Stembark	5,32a	0,69a	4,91a	8,08b	3,68ab	0,35a	13,14a	2,56a	27,70b	285,22ab	13,29*
	Stemwood	0,45a	0,11b	1,52bc	0,37b	0,20c	0,18a	1,86ab	0,73b	20,26a	14,66a	4,58ab
<i>E. saligna</i>	Leaves	20,61a	1,26ab	8,31a	4,63a	3,11ab	1,26a	28,72a	5,45a	77,23b	179,76a	11,96ab
	Branches	1,57a	0,29ab	3,14a	5,56a	1,93a	0,34a	7,13*	6,55a	39,74a	107,19b	8,32*
	Stembark	2,16bc	0,53a	4,19a	9,10b	3,72a	0,33a	11,25a	3,72a	32,32ab	238,13b	7,26*
	Stemwood	0,74a	0,08b	1,32c	0,47b	0,36b	0,21a	1,81ab	1,10ab	27,69a	6,49a	4,25ab
<i>E. dunnii</i>	Leaves	21,62a	1,23ab	6,08c	6,49a	3,55a	1,26a	19,52bc	6,38a	100,98ab	300,37a	15,17*
	Branches	1,89a	0,25ab	2,44a	4,21a	1,79a	0,31a	6,40ab	4,54ab	47,51a	165,63ab	7,71*
	Stembark	4,08ab	0,42a	5,84a	9,17b	3,23ab	0,25a	12,83a	3,06a	32,14ab	288,47ab	9,61*
	Stemwood	0,80a	0,09b	1,37c	0,70a	0,55a	0,19a	2,75*	0,92ab	23,34a	18,94a	2,84c
<i>E. urolobulus</i>	Leaves	18,46a	0,99b	6,27bc	4,18a	2,13b	1,01b	15,20c	4,27a	83,19ab	222,94a	10,32b
	Branches	1,54a	0,19b	3,34a	3,76a	0,99b	0,26a	6,20ab	3,15b	46,76a	95,45b	7,38*
	Stembark	2,70bc	0,42a	6,45a	7,84b	2,92b	0,35a	14,21a	2,72a	29,16ab	231,58b	8,75*
	Stemwood	0,91a	0,08b	1,60bc	0,44b	0,18c	0,23a	1,30b	1,06ab	65,58a	10,37a	3,41bc
<i>E. urograndis</i>	Leaves	21,10a	1,21ab	7,67abc	5,35a	3,98a	1,29a	30,72a	6,49a	74,51b	196,63a	12,81ab
	Branches	0,81a	0,22ab	3,14a	5,08a	1,86a	0,26a	4,87b	4,58ab	48,48a	172,69ab	8,55*
	Stembark	1,90c	0,59a	4,42a	9,33b	3,36ab	0,34a	8,49*	3,22a	56,56a	284,29ab	6,88*
	Stemwood	0,87a	0,09b	1,71ab	0,55ab	0,27bc	0,18a	3,10*	1,47a	30,55a	10,20a	4,74*

216 Averages of each fraction of biomass in different treatments (genotype of *Eucalyptus*)
 217 followed by equal letters, do not differ significantly by the Tukey test at the 5% level of
 218 error.

219

220 This same trend, i.e., the highest concentration of nutrients in the leaves and the
 221 lowest in the wood, was also reported in populations of *E. urograndis* at 18 months of age
 222 in Piratini, RS, Brazil [25], in an *E. dunnii* stand at four years of age in Alegrete, RS, Brazil
 223 [8], and in *E. globulus* in Chile [1].

224 In relation to the analyzed macronutrients, N, P, K, and S were more concentrated in
 225 the leaves in most genotypes, except for *E. urolobulus*, where K was more concentrated in
 226 the bark. Ca and Mg were found in higher concentrations in the bark in most genotypes,
 227 except for *E. dunnii* and *E. urograndis*, in which the highest Mg content occurred in the

228 leaves. In an *E. dunnii* stand at 9 years of age in Algorta, Uruguay, the highest
 229 concentrations of N, P, and K were found in the leaves, but the bark also accumulated high
 230 concentrations of nutrients, mainly Ca [10].

231 For micronutrients, the highest concentrations occurred in the leaves, except for Cu
 232 in *E. saligna*, which the highest concentration was observed in the branches; and for Mn in
 233 *E. benthamii* (P1), *E. saligna*, *E. urolobulus*, and *E. urograndis*, and Zn in *E. benthamii*
 234 (P2), which the highest concentrations were observed in the bark. This same trend, with
 235 higher content of micronutrients in leaves, was also found by [26] in *Eucalyptus urograndis*
 236 stands at 18 months of age in Piratini-RS municipality.

237

238 3.3 Amount of nutrients

239 Nitrogen occurred in greater quantities in the leaves of most genotypes, with the
 240 exception of *E. urograndis* in which higher concentrations of N were found in the wood. P,
 241 K, and S had greater representation in the wood, and Ca and Mg in the bark in most
 242 genotypes, except in *E. dunnii*, in which the highest amount of Mg was observed in the
 243 wood (Table 5). Micronutrients were stored more in the wood, with the exception of Mn
 244 which accumulated in higher concentrations in the bark.

245

246 Table 5 - Amount of nutrients in the biomass components of different genotypes of
 247 *Eucalyptus* at 49-month-old established in Eldorado do Sul, RS, Brazil

248

Genotypes of <i>Eucalyptus</i>	Fractions	N	P	K	Ca	Mg	S	B	Cu	Fe	Mn	Zn
		kg ha ⁻¹							g ha ⁻¹			
<i>E. benthamii</i> (P1)	Leaves	99,70	6,00	36,00	26,50	12,40	5,90	86,00	21,40	586,30	2.033,50	70,40
	Branches	13,40	2,40	23,50	39,30	12,80	2,20	43,60	26,30	362,90	2.173,40	76,20

	Stembark	44,10	4,80	48,60	129,80	30,20	2,70	115,30	26,10	265,20	4.164,70	130,70
	Stemwood	65,50	12,30	143,50	41,90	20,10	13,20	210,30	78,30	4.956,10	1.515,90	360,90
	Total	222,70	25,60	251,60	237,60	75,50	24,00	455,20	152,20	6.170,60	9.887,50	638,20
	Leaves	92,50	5,00	26,00	20,20	10,70	5,00	95,60	22,40	490,40	1.400,70	51,30
	Branches	7,70	1,30	14,60	14,90	6,70	1,50	26,40	14,20	228,60	835,00	47,00
<i>E. benthamii</i> (P2)	Stembark	45,60	6,00	42,30	70,40	31,80	3,00	113,20	21,90	240,50	2.452,30	113,30
	Stemwood	37,90	9,60	127,60	31,40	16,70	15,50	155,20	61,40	1.688,90	1.230,00	383,60
	Total	183,70	21,80	210,50	136,90	65,80	25,00	390,40	119,70	2.648,50	5.918,00	595,20
	Leaves	66,50	4,00	26,70	14,90	10,00	4,00	92,50	17,60	248,40	574,90	38,60
	Branches	8,70	1,60	17,60	31,30	10,80	1,90	39,80	36,60	221,60	594,70	46,40
<i>E. saligna</i>	Stembark	17,00	4,20	33,10	72,10	29,50	2,60	89,20	29,50	255,20	1.882,40	57,50
	Stemwood	53,60	5,90	95,70	34,50	25,90	15,10	131,20	80,00	1.993,90	466,70	309,80
	Total	145,70	15,70	173,10	152,80	76,10	23,70	352,80	163,80	2.719,10	3.518,70	452,30
	Leaves	66,00	3,80	18,50	19,90	10,80	3,90	59,50	19,50	309,90	941,40	46,70
	Branches	8,20	1,10	10,90	19,30	8,10	1,40	29,30	20,40	217,20	791,90	34,80
<i>E. dunnii</i>	Stembark	24,50	2,50	34,70	55,30	19,40	1,50	77,50	18,30	193,50	1.818,30	57,00
	Stemwood	41,50	4,80	74,10	38,20	30,30	10,40	150,40	51,00	1.304,20	1.070,80	155,20
	Total	140,30	12,20	138,20	132,70	68,60	17,10	316,70	109,20	2.024,80	4.622,30	293,70
	Leaves	120,00	6,40	40,80	27,30	14,00	6,60	99,40	27,60	548,80	1.457,60	67,80
	Branches	11,40	1,40	25,40	28,20	7,40	1,90	46,80	23,50	349,40	729,80	55,40
<i>E. urolobulus</i>	Stembark	17,90	2,80	43,50	51,90	19,70	2,40	95,40	18,40	198,00	1.556,70	59,20
	Stemwood	88,10	7,70	154,80	42,80	17,50	22,10	125,80	103,20	6.782,90	1.018,30	328,50
	Total	237,40	18,40	264,50	150,20	58,70	33,00	367,40	172,70	7.879,00	4.762,30	511,00
	Leaves	64,30	3,70	23,40	16,30	12,10	3,90	93,60	19,80	227,10	599,40	39,10
	Branches	6,00	1,70	23,20	37,70	13,80	1,90	36,10	33,90	359,20	1.279,60	63,30
<i>E. urograndis</i>	Stembark	14,80	4,60	34,30	72,40	26,10	2,70	65,90	25,00	438,90	2.206,10	53,40
	Stemwood	73,00	7,30	143,00	46,10	22,90	14,80	259,50	122,50	2.553,30	852,30	396,40
	Total	158,00	17,20	223,90	172,50	74,90	23,40	455,00	201,10	3.578,50	4.937,40	552,10

249

250 The concentrations of macronutrients in the total biomass followed the order: K >
251 Ca > N > Mg > S > P in most genotypes. In *E. benthamii* (P1), however, P content was
252 higher than S content. In *E. benthamii* (P2) and *E. urolobulus*, N content was higher than
253 Ca content. For *E. dunnii*, the concentrations of macronutrients in the total biomass
254 followed the order: N > K > Ca > Mg > S > P. In stands of *E. urograndis* at 30 and 60
255 months of age in Seropédica, Rio de Janeiro, Brazil, the following order was observed: K >
256 N > Ca > Mg > P [18].

257 For micronutrients, the order of concentrations in most genotypes was: Mn > Fe >
258 Zn > B > Cu, except for *E. dunnii*, whose the amount of B was greater than that of Zn; and
259 for *E. uroglobulus*, whose the amount of Fe was higher than that of Mn.

260 The highest amount of P, Ca, B, Mn, and Zn was found in *E. benthamii* (P1); of N,
261 K, S, and Fe in *E. uroglobulus*; of Mg in *E. saligna*; and of Cu in *E. urograndis*. In *E.*
262 *uroglobulus* was observed with 39 and 41% more than N and 35 and 48% more of K than
263 the *E. saligna* and *E. dunnii* clones, respectively. In *E. benthamii* (P1) P concentrations
264 were found to be 33, 39, and 52% higher compared to *E. urograndis*, *E. saligna*, and *E.*
265 *dunnii*, respectively.

266 The canopy (leaves and branches) accumulated between 17 and 52% of the total
267 macronutrients in *E. benthamii* (P1) and *E. uroglobulus*, and from 24 to 34% of total
268 micronutrients in *E. dunnii* and *E. uroglobulus*. The stem (wood and bark) accumulated
269 between 48 to 83% and 66 to 76% of the total macro and micronutrients, respectively.

270 The distribution and total content of nutrients in the canopy are affected mainly by
271 changes in the amount of biomass and by differences that occur owing to age, both of the
272 tree and the leaves, in their different physiological stages [3].

273

274 **3.4 Nutrient use efficiency**

275 Genotypes and their different components showed variations in nutrient use
276 efficiency (NUE) (Table 6). With the exception of Fe in *E. benthamii* (P1) and *E.*
277 *uroglobulus*, in which NUE was larger in the stembark, and of N in *E. urograndis*, where
278 the branches had the highest concentrations, the stemwood presented the highest values of

279 NUE, which is very relevant to forest companies, because this is the main product taken
 280 from forest plantations.

281

282 Table 6 – Nutrient use efficiency in the biomass components of different genotypes of
 283 *Eucalyptus* at 49-month-old established in Eldorado do Sul, RS, Brazil

284

Genotypes of <i>Eucalyptus</i>	Fractions	N	P	K	Ca	Mg	S	B	Cu	Fe	Mn	Zn
<i>E. benthamii</i> (P1)	Leaves	44	723	121	164	351	740	50.706	203.368	7.437	2.144	61.975
	Branches	526	2.907	300	179	549	3.176	161.441	267.639	19.407	3.241	92.390
	Stembark	185	1.699	168	63	271	3.070	70.875	312.609	30.799	1.961	62.494
	Stemwood	1.116	5.939	509	1.742	3.634	5.514	347.279	933.091	14.737	48.182	202.399
<i>E. benthamii</i> (P2)	Leaves	42	791	151	194	368	784	40.987	175.244	7.992	2.798	76.460
	Branches	660	3.878	347	341	757	3.473	192.308	358.293	22.212	6.080	108.033
	Stembark	189	1.441	203	122	271	2.907	75.968	393.553	35.754	3.507	75.873
	Stemwood	2.228	8.795	662	2.692	5.071	5.437	544.211	1.376.084	49.995	68.651	220.141
<i>E. saligna</i>	Leaves	49	798	121	216	322	797	34.868	182.763	12.979	5.608	83.607
	Branches	645	3.482	319	179	520	2.981	140.612	152.949	25.284	9.421	120.734
	Stembark	466	1.891	239	110	269	2.989	88.700	268.164	31.023	4.206	137.632
	Stemwood	1.353	12.391	758	2.103	2.803	4.807	552.606	905.966	36.364	155.370	234.061
<i>E. dunnii</i>	Leaves	47	817	167	155	285	798	51.850	158.141	9.962	3.279	66.107
	Branches	556	4.047	419	237	564	3.307	156.571	224.899	21.106	5.790	131.794
	Stembark	247	2.379	174	109	312	4.018	78.002	330.696	31.254	3.326	106.134
	Stemwood	1.317	11.439	738	1.431	1.807	5.279	363.615	1.071.303	41.925	51.065	352.245
<i>E. urolobulus</i>	Leaves	54	1.014	160	239	465	988	65.593	236.128	11.878	4.472	96.073
	Branches	657	5.269	294	265	1.009	3.892	159.548	317.854	21.379	10.236	134.777
	Stembark	374	2.356	154	129	339	2.823	70.113	363.003	33.799	4.299	112.953
	Stemwood	1.100	12.617	626	2.265	5.519	4.376	770.081	938.682	14.277	95.103	294.774
<i>E. urograndis</i>	Leaves	47	825	130	187	252	776	32.552	154.162	13.421	5.086	78.057
	Branches	1.241	4.467	319	197	538	3.803	205.480	218.341	20.629	5.791	117.012
	Stembark	525	1.684	227	107	297	2.914	117.801	310.776	17.681	3.518	145.423
	Stemwood	1.145	11.475	584	1.811	3.645	5.634	322.119	682.490	32.734	98.058	210.872

285

286 In general, the highest values of NUE were found in micronutrients, where Cu stood
 287 out in all biomass components but presented greater values in the stemwood. However, Mn

288 had a lower NUE in most components, with the exception of the stemwood, in which the
289 lowest NUE was found for Fe. For this component, nutrient use efficiency decreased in the
290 following order: Cu > B > Zn > Mn > Fe.

291 In relation to macronutrients, P stood out as the most utilized element in the
292 stemwood. In contrast, N presented the least efficiency in the leaves. The NUE of the
293 stemwood for macronutrients decreased in the following order in most genotypes: P > S >
294 Mg > Ca > N > K, with the exception of *E. urolobulus* in which Mg was higher than S.
295 Similar results, although with inversion in the distribution of some nutrients, were reported
296 by [18] while studying *E. urograndis* at the age of five in Seropédica, RS, Brazil (P > Mg >
297 Ca > N > K); by [17] while evaluating the provenance of *E. grandis* and *E. saligna* in forest
298 sites of São Paulo, Brazil (P > Mg > K > N > Ca); and by [5] while studying *E. urograndis*
299 at two years of age in Botucatu, São Paulo, Brazil (P > Mg > S > N > K > Ca). The
300 variation in nutrient use efficiency can occur due to several factors, such as: the intrinsic
301 characteristics of the genotype, the failure to obtain optimal or critical nutritional balance
302 between the soil and the plant and water conditions [17].

303 In general, the lowest NUE values were found in the leaves, with the exception of
304 some elements, in which the lowest coefficients were observed in the stembark, as was the
305 case for Ca, Mg, and Mn in the clones *E. benthamii* (P1) and *E. saligna*; for K, Ca, Mg, and
306 Mn in *E. urolobulus*; for Ca and Mg in *E. benthamii* (P2); for Ca and Mn in *E.*
307 *urograndis*; and for Ca in *E. dunnii*. In this context, the harvesting of the leaves will result
308 in the greatest export of nutrients, especially N and K. In contrast, considering only the
309 harvesting of the stemwood with bark, Ca and Mg are the limiting nutrients in terms of the
310 productivity of the next cycle, but this limitation may be reduced if only the wood is

311 harvested. In relation to the other biomass components, P presented the highest NUE for the
312 leaves and branches in most genotypes, except for *E. benthamii* (P1) where S had the
313 highest value. As for the bark of the shaft, the largest NUE was found for S.

314 Taking into account the greater commercial interest in stemwood, it was observed
315 that the highest biomass yields were accompanied with the highest values of nutritional
316 efficiency for some elements, that is, the highest efficiency values for *E. urolobulus* (P,
317 Mg, and B) and *E. benthamii* (P2) (N, Ca, Cu, and Fe). Regarding the other genotypes, *E.*
318 *saligna* showed higher efficiency for K and Mn, *E. urograndis* for S, and *E. dunnii* for Zn.
319 The high efficiency presented by a species in the use of nutrients implies that it has a lower
320 nutritional requirement, therefore, a parameter of great utility in the selection of species to
321 be used in reforestation, especially in nutrient poor soils [22].

322

323 4. CONCLUSIONS

324 The different genotypes of *Eucalyptus*, under the same edaphoclimatic conditions,
325 present different biomass production.

326 There are a great variation in the concentration and allocation of the amount of
327 nutrients in the different genotypes of *Eucalyptus* and in the different components of the
328 same genotypes.

329 The highest biomass yields were accompanied with the highest values of nutritional
330 efficiency for some elements, that is, the highest efficiency values for *E. urolobulus* (P,
331 Mg, and B) and *E. benthamii* (P2) (N, Ca, Cu, and Fe).

332

333

334

335 **5. REFERENCES**

336 1. Albaugh TJ, Rubilar RA, Maier CA, Acuña EA, Cook RL. Biomass and nutrient mass of
337 *Acacia dealbata* and *Eucalyptus globulus* bioenergy plantations. *Biomass and Bioenergy*.
338 2017; 97: 162-171.

339

340 2. Alvares CA, Stape JL, Sentelhas PC, Gonçalves JLM, Sparovek G. Köppen's climate
341 classification map for Brazil. *Meteorologische Zeitschrift*. 2013; 22 (6): 1-18.

342

343 3. Bellote AFJ, Silva HD. Sampling techniques and nutritional evaluations in eucalypt
344 plantations, In: Gonçalves JLM, Benedetti V. *Forest nutrition and fertilization*. Piracicaba:
345 IPEF; 2004.

346

347 4. Binkley D, Campoe OC, Alvarez C, Carneiro RL, Cegatta I, Stape JL. The interactions
348 of climate, spacing and genetics on clonal *Eucalyptus* plantations across Brazil and
349 Uruguay. *Forest Ecology and Management*. 2017; 405: 271-283.

350

351 5. Eufraide Junior HJ, Melo RX, Sartori MMP, Guerra SPS, Ballarin AW. Sustainable use
352 of eucalypt biomass grown on short rotation coppice for bioenergy. *Biomass and*
353 *Bioenergy*. 2016; 90: 15-21.

354

- 355 6. Gatto A, Barros NF, Novais RF, Silva IR, Leite HC, Villani EMA. Estoque de carbono
356 na biomassa de plantações de eucalipto na região centro-leste do estado de Minas Gerais.
357 Revista Árvore. 2011; 35 (4): 895-905.
358
- 359 7. Gonçalves JLM, Alvares CA, Higa AR, Silva LD, Alfenas AC, Stahl J et al. Integrating
360 genetic and silvicultural strategies to minimize abiotic and biotic constraints in Brazilian
361 eucalypt plantations. Forest Ecology and Management. 2013; 301: 6-27.
362
- 363 8. Guimarães CC, Schumacher MV, Witschorek R, Souza HP, Santos JC. Biomassa e
364 nutrientes em povoamento de *Eucalyptus dunnii* Maiden no Pampa Gaúcho. Revista
365 Árvore. 2015; 39 (5): 873-882.
366
- 367 9. Guo LB, Sims REH, Horne DJ. Biomass production and nutrient cycling
368 in *Eucalyptus* short rotation energy forests in New Zealand.: I: biomass and nutrient
369 accumulation. Bioresource Technology. 2002; 85 (3): 273-283.
370
- 371 10. Hernández J, Pino A, Salvo L, Arrarte S. Nutrient export and harvest residue
372 decomposition patterns of a *Eucalyptus dunnii* Maiden plantation in temperate climate of
373 Uruguay. Forest Ecology and Management. 2009; 258 (2): 92-99.
374
- 375 11. IBÁ, Indústria Brasileira de árvores: ano base 2016/IBÁ. Brasília/DF, 2016.
376

- 377 12. Kuyah S, Dietz J, Muthuri C, Noordwijk M, Neufeldt H. Allometry and partitioning of
378 above- and Bellow-ground biomass in farmed eucalyptus species dominant in Western
379 Kenyan agricultural landscapes. *Biomass and Bioenergy*. 2013; 55 (1): 276-284.
380
- 381 13. Miyazawa M, Pavan MA, Muraoka T. Análises químicas de tecido vegetal. In: Silva,
382 F.C. Manual de análises químicas de solos, plantas e fertilizantes. Brasília: Embrapa
383 Comunicação para Transferência de Tecnologia, p. 171-224; 1999.
384
- 385 14. Pérez-cruzado C, Rodríguez-Soalleiro R. Improvement in accuracy of aboveground
386 biomass estimation in *Eucalyptus nitens* plantations: Effect of bole sampling intensity and
387 explanatory variables. *Forest Ecology and Management*. 2011; 261(1): 2016-2028.
388
- 389 15. Ryan MG, Stape JL, Binkley D, Fonseca S, Loos RA, Takahashi EM et al. Factors
390 controlling *Eucalyptus* productivity: How water availability and stand structure alter
391 production and carbon allocation. *Forest Ecology and Management*. 2010; 259 (9): 1695-
392 1703.
393
- 394 16. Salvador SM, Schumacher MV, Viera M, Stahl J, Consensa CB. Biomassa e estoque de
395 nutrientes em plantios clonais de *Eucalyptus saligna* Smith. em diferentes idades. *Scientia*
396 *Forestalis*. 2016; 44 (110): 311-321.
397

- 398 17. Santana RC, Barros NF, Neves JCL. Eficiência de utilização de nutrientes e
399 sustentabilidade da produção em procedências de *Eucalyptus grandis* e *Eucalyptus saligna*
400 em sítios florestais do estado de São Paulo. Revista Árvore. 2002; 26 (4): 447-457.
401
- 402 18. Santos FM, Chaer GM, Diniz AR, Balieiro FC. Nutrient cycling over five years of
403 mixed-species plantations of *Eucalyptus* and *Acacia* on a sandy tropical soil. Forest
404 Ecology and Management. 2017; 384: 110-121.
405
- 406 19. Schumacher MV, Caldeira MVW. Estimativa da biomassa e do conteúdo de nutrientes
407 de um povoamento de *Eucalyptus globulus* (Labillardière) sub-espécie Maidenii. Ciência
408 Florestal. 2001; 11(1): 45-53.
409
- 410 20. Schumacher MV, Witschoreck R, Calil FN. Biomassa em povoamentos de *Eucalyptus*
411 spp. de pequenas propriedades rurais em Vera Cruz, RS. Ciência Florestal. 2011; 21 (1),
412 17-22.
413
- 414 21. Silva FAZ, Azevedo CAV. Principal components analysis in the software assistat
415 statistical attendance. In: Word Congress on Computers in Agriculture 7, Reno-NV-USA:
416 American Society of Agricultural and Biological Engineers; 2009.
417
- 418 22. Silva HD, Poggiani F, Coelho LC. Eficiência de utilização de nutrientes em cinco
419 espécies de *Eucalyptus*. Boletim de Pesquisa Florestal. 1983; (6/7): 1-8.
420

- 421 23. Silva PHM, Poggiani F, Libaldi PL, Gonçalves AW. Fertilizer management of eucalypt
422 plantations on sandy soil in Brazil: Initial growth and nutrient cycling. *Forest Ecology and*
423 *Management*. 2013; 301: 67-78.
- 424
- 425 24. Tedesco MJ, Gianello C, Bissani CA, Bohnen H, Volkweiss SJ. Análise de solo, plantas
426 e outros materiais. (2.ed.). Porto Alegre, RS: Departamento de Solos, UFRGS; 1995.
- 427
- 428 25. Viera M, Schumacher MV, Trüby P, Araújo EF. Biomassa e nutrientes em um
429 povoamento de *Eucalyptus urophylla* x *Eucalyptus globulus*, em Eldorado do Sul-RS.
430 *Revista Ecologia e Nutrição Florestal*. 2013; 1 (1) :1-13.
- 431
- 432 26. Viera M, Bonacina DM, Schumacher MV, Calil FN, Caldeira MVW, Watzlawick LF.
433 Biomassa e nutrientes em povoamento de *Eucalyptus urograndis* na Serra do Sudeste-RS.
434 *Semina: Ciências Agrárias*. 2012;33 (1): 2481-2490.
- 435
- 436 27. Viera M, Schumacher MV, Bonacina DM, Ramos LOO, Rodríguez-Soalleiro R.
437 Biomass and nutrient allocation to aboveground components in fertilized *Eucalyptus*
438 *saligna* and *E. urograndis* plantations. *New Forests*. 2017; 47: 1-18, 2017.
- 439
- 440 28. Zhang H, Guan D, Song M. Biomass and carbon storage of *Eucalyptus* and *Acacia*
441 plantations in the Pearl River Delta, South China. *Forest Ecology and Management*. 2012;
442 277: 90-97.