

Agronomic efficiency of bone meal under acidification in *Brachiaria ruziziensis* dry matter production in Western Amazon

ABSTRACT

Aims: The objective of this work was to evaluate the agronomic efficiency of the bone meal under acidification in the production of dry matter of *Brachiaria ruziziensis* in relation to a soluble source.

Study Design: The experiment was conducted in a completely cascaded design with seven treatments and three replications: Witness; Single Superphosphate (SS); Bone Meal (BM); BM treated with 10% Oxalic Acids; 10% Acetic Acid and 1% to 0.5% Hydrochloric Acid.

Place and Duration of Study: The experiment was carried out from October 2014 to February 2015, at the Experimental Farm of the Federal University of Rondônia (UNIR), located 15 km from the city of Rolim de Moura, Rondônia, Brazil.

Methodology: The bone meal used in the experiment was produced manually, where bovine bones were collected and burned for carbon removal and particle reduction. The oxalic, acetic and hydrochloric acids were used to increase the solubility of the bone meal and applied to the soil for growth of *B. ruziziensis*. It was evaluated the Agronomic Efficiency Index (AEI), Phosphorus Conversion Efficiency (PCE), Shoot Dry Matter (SDM) and Root Dry Matter (RDM).

Results: AEI was obtained for acid treatments above 60% and PCE satisfactory when compared to the soluble source, except for 0.5% hydrochloric acid and significantly above the BM without acid treatments. There was higher production of SDM and RDM with the soluble source (SS), however the acid treatments promoted dry matter production above the BM without acid treatments.

Conclusions: The application of acids in bone meal promoted satisfactory agronomic efficiency gains for plants of *Brachiaria ruziziensis*.

Keywords: *Phosphor; Forage; Exploitation*

1. INTRODUCTION

The Brazil is the second largest producer of beef in the world, behind only the United States, and according to the Brazilian Institute of Geography and Statistics [1], the herd surpasses 218 million head. Approximately 80% of the Brazilian herd is extensively raised under planted / natural pastures, since it is the most economical method of cattle feeding to cattle ranchers [2]. Of the fodder used for this purpose, those of the genus *Brachiaria* correspond to 70% and 80% of the area planted in Brazil. The *Brachiaria ruziziensis* is one of the most planted because of its high palatability, grazing support and high dry matter production (6 to 15 t ha⁻¹) [3].

It is estimated, however, that about 70 to 80% of the pastures used for this purpose are degraded or in the process of degradation [4]. According to [2], the main causes of this degradation are focused on the adopted management, such as fire use, low quality seeds, high stocking rates and absence of fertilization. Among these, the absence and inefficiency of fertilization has been noted as a more aggravating factor of pasture degradation in Brazil [5,6]. As soil fertility decreases, there is a marked productive loss of fodder and, consequently, decrease in animal production [3].

When considering the Amazon region, the problem tends to be more pronounced, considering the most acidic soil characteristics, low cation exchange capacity, high aluminum saturation and low nutrient reserves, mainly in phosphorus (P) (between 1 and 3 g dm⁻³) [2]. It is suggested, therefore, that fertilization with this nutrient is carried out, in order to guarantee its adequate supply.

Phosphorus maintenance for pasture production has been performed with the use of liming and the use of soluble phosphorus sources, such as single and triple superphosphate [7]. Although widely used, they come from the exploration of phosphate rocks, whose reserves are estimated to deplete in up to 100 years [8 9]. In this sense, it is necessary the study of new sources of phosphorus in order to guarantee the supply of this element [10,6].

Recently, studies by several authors have shown that the use of solid animal waste processed, bone meal (BM), have been pointed as potential source of phosphorus and calcium plants [11,11]. Under heat action, bones can also supply magnesium, potassium and iron from the blood [13]. However, due to the chemical nature of the bones (hydroxyapatite), BM has low solubility of phosphorus in water (0.26% of P₂O₅) despite its high concentration, between 30, 6 and 38.8% P₂O₅ [14]. However, it is necessary to study acidifying agents capable of readily providing the available phosphorus in the bones, as tested for 2% citric acid, which provided 23% solubility to bone meal [15].

Thus, the objective of this study was to evaluate the agronomic efficiency of the bone meal under acidification in the production of dry matter of *Brachiaria ruziziensis* in relation to a soluble source.

2. MATERIAL AND METHODS

The experiment was conducted at the Experimental Sector of Agronomy of the Federal University of Rondônia - UNIR, Campus Rolim de Moura, RO - Brazil, located at coordinates 11° 34 '58.60 "S and 61°46'22.30" W with an altitude of 277 m. The climate of the region, according to the classification of Koppen, is Aw type, defined dry season, mean temperature of 28 ° C, mean annual precipitation of 2,250 mm and relative humidity of 85% [16].

The study was conducted in open-air pots of 3.9 kg in a completely randomized design, with seven treatments and three replicates, totaling 21 experimental units (Table 1).

Table 1. Treatments used in experimentation.

Treatments	Description	Symbol
1	Unmanaged Witness	Wit.
2	P (Simple Superphosphate)	SS
3	P (Bone Meal)	BM
4	(Bone Meal + 10% Oxalic Acid)	Oxa ^{10%}
5	(Bone Meal + 10% Acetic Acid)	Ace ^{10%}
6	(Bone Meal + 1% hydrochloric acid)	Hcl ^{1%}
7	(Bone Meal + 0.5 % hydrochloric acid)	Hcl ^{0,5%}

The conduction of the open - air experiment aimed to study the effect of plants under natural conditions. As a source of phosphate fertilization, bone meal (37.5% P_2O_5) and simple superphosphate (18% P_2O_5) were used. For each source, the equivalent of 100 kg ha^{-1} of P_2O_5 was used, depending on the concentration of each source. For the evaluations under equal conditions, both P sources were subjected to grinding and sieving of 2.00 mm.

As substrate for the plants, the arable layer (0-20 cm) of the experimental area classified as Eutrophic Yellow Red Oxisol [17] was used, with the following chemical and physical characteristics: pH in water = 5.4 ; Organic matter = 3 dag kg^{-1} ; P = 3.7 mg dm^{-3} ; K = 102 mg dm^{-3} ; S- SO_4^{2-} = 3.6 mg dm^{-3} ; Ca = 4.1 cmol_c dm^{-3} ; Mg: 1.4 cmol_c dm^{-3} ; Fe = 88 mg dm^{-3} ; Cu = 1.8 mg dm^{-3} ; Zn = 1.5 mg dm^{-3} ; Mn = 25 mg dm^{-3} ; B = 0.14 mg dm^{-3} ; Potential Acidity= 4.8 cmol_c dm^{-3} ; Al = 0.12 cmol_c dm^{-3} ; Sum of Bases = 5.7 cmol_c dm^{-3} ; Potential cation exchange capacity = 10.5 cmol_c dm^{-3} ; Basis Saturation= 54%; Saturation by aluminum = 2%; Sand = 530 g kg^{-1} ; Silt = 83 g kg^{-1} and Clay = 381 g kg^{-1} .

The bone meal was produced under burn for elimination of organic compounds and for maceration in smaller particles. For this, the bovine bones were arranged inside a masonry barbecue and accommodated in coin type screens, using charcoal for the burning for approximately eight hours. After cooling the material, about 24 hours later, the bones were ground with the use of gral and pistil and sieved in a 2.00 mm mesh. The material was characterized in laboratory for analysis of the contents of P and Ca, presenting 37.5% P_2O_5 and 43.76% CaO.

As acid extractors of phosphorus, the oxalic, acetic and hydrochloric acids (PAs) were used. For Oxalic and Acetic acids, 10 ml dilution in 100 ml of distilled water was used. For hydrochloric acid, dilution of 1 and 0.5 ml in 100 ml of distilled water was used. The solutions were applied to 100 g of bone meal. Subsequently, the material was forced to air drying at 65 ° C for approximately 72 hours. The bone meal was homogenized to the soil and arranged in the experimental units.

The Brachiaria ruziziensis sprouts collected in the experimental area and standardized (12 cm in height and mean mass of 4.52 g) were used as indicator of the treatments. Three shoots of *B. ruziziensis* were arranged in each experimental unit. In addition to the treatments, complementary fertilization with macro and micronutrients was applied as recommended for the forage described by [3]. The water supply was performed by daily manual irrigation, applying 300 ml / pot / day. Cultural treatments were carried out regarding the manual removal of invasive plants from the soil seed bank.

The plants were cultivated for 120 days, with Shoot Dry Matter (SDM) at 60, 90 and 120 days. The forage was cut at 20 cm from the soil level and the SDM determined from drying in a forced circulation oven at 65 °C for approximately 72 hours. At the end of the periods the sum of the SDM produced in the three periods was calculated for the calculation of the Agronomic Efficiency Index (AEI) and Phosphorus Conversion Efficiency (PCE).

The AEI was calculated from equation 1:

$$AEI (\%) = [(Y_2 - Y_1) / (Y_3 - Y_1)] \times 100 \text{ (Equation 1)}$$

Where: AEI = Agronomic Efficiency Index, in%; Y_1 = SDM obtained in the control, in g; Y_2 = SDM obtained with the treatments, in g; Y_3 = SDM obtained from the reference source (SS), in g.

The PCE was calculated from equation 2:

$$PCE (mg\ mg^{-1}) = (SDM\ Treatments - SDM\ Witness) / APA\ (Equation\ 2)$$

Where: PCE = Phosphorus Conversion Efficiency, mg mg⁻¹; SDM Treatments = Dry Matter of the Aerial Part Total obtained in each treatment, in mg; SDM Witness = Dry matter of the Total Aerial Part obtained in the control; APA = Amount of Phosphorus Added, in mg / pot.

The dry matter of the roots (RDM) was obtained after 120 days of treatments. For this, the roots were removed from the experimental units, washed and subjected to drying in forced circulation oven at 65 °C for approximately 72 hours. SDM and RDM were determined using a precision digital scale. The obtained data were submitted to the analysis of variance and the contrast of the means realized by the test of Tukey to 5% of probability. For the analyzes, the statistical program ASSISTAT 7.7 was used.

3. RESULTS AND DISCUSSION

Significant results were observed in all studied variables (P <0.01). When analyzing table 1, it is noticed that the Total Shoot Dry Matter (SDM) was significantly higher in the commercial soluble source (SS) application. However, acidification of bone meal (BM) with 10% acetic acid, 10% Oxalic acid and 1% Chloride promoted an increase of 224%, 231% and 249% in relation to treatment that did not receive acidification (BM). In relation to the control, the application of BM without acidification promoted increments of only 56.1%.

With the exception of the treatment with 0.5% hydrochloric acid, the other acid treatments promoted a significantly similar effect on the Agronomic Efficiency Index (AEI), with the lowest AEI observed, confirming its natural phosphate characteristics. Oxa 10%, Ace 10% and Hcl 1% acidifying treatments promoted I AEI on average 31% lower, taxing much lower by observing the AEI without acidification of FO. Owing to the efficiency scale, it can be noted that the AEI decreased in the following order: SS> HCl1%> Ace10%> Oxa10%> HCl 0.5%> BM (Table 2).

Phosphorus Conversion Efficiency (PCE) was significantly higher in the standard treatment (SS), however high PCE values were obtained with acid treatments, when compared to those that did not receive acidification (BM). Among the acid treatments, HCl0.5% was the one with the lowest efficiency. The PCE obeyed the following order: SS> HCl1%> Ac10%> Oxa10%> HCl0.5%> BM (Table 2).

Table 2. Values of Shoot Dry Matter (SDM) of *Brachiaria ruziziensis*, Agronomic Efficiency Index (AEI) and Phosphorus Conversion Efficiency (PCE) of bone meal (BM) under acidification.

Treatments	SDM g	AEI %	PCE mg mg ⁻¹
Witness	4,1 d	----	----
SS	23,1 a	100 a	223,13 a
BM	7,3 cd	16,84 d	37,58 c
BM+ Oxa ^{10%}	16,4 abc	64,73 b	144,45 abc
BM+ Ace ^{10%}	16,9 ab	67,36 b	150,32 abc
BM+ HCl ^{1%}	18,1 ab	73,68 b	164,41 ab
BM+ HCl ^{0,5%}	10,3 bcd	32,63 c	72,81 bc
C.V	8,28	7,03	23,09

F	182,7	166,8	7,50
p	0,001	0,001	0,002

C.V: Coefficient of Variation; BM: Bone Meal; SS: Simple Superphosphate; Oxa10%: Bone Meal + 10% Oxalic Acid; Ace10%: Bone Meal + 10% Acetic Acid; HCl1%: Bone Meal + 1% hydrochloric acid; HCl 0,5%: Bone Meal + 0.5 % hydrochloric acid

186

187

188 Due to the high solubility of simple superphosphate (SS) in water [18], shoot dry matter
189 production (SDM) was substantially higher, given the high availability of phosphorus (P) in
190 solution. The same behavior was observed by [19] when using SS compared to bone meal
191 (BM) in Tifton grass. However, these same authors observed satisfactory results with BM.

192

193 In a study using bone meal and meat, [20] observed results of SDM of Tifton grass varying
194 between 65 and 93% in relation to fertilization with SS, a fact not observed in this work,
195 where the application of the alternative source of P resulted in about 30% and the extraction
196 of the acids between 70.9 and 78%. In general terms, the production of dry matter with
197 natural phosphate sources depends on the forage species and the BM production method.

198 The low natural solubility of BM may compromise the short-term response of annual crops,
199 such as *B. ruziziensis*, and is not fully indicated for efficient production of SDM, especially in
200 soils with pH above 5.0, where the reactions with phosphate naturally occur slowly and
201 progressively [15].

202 Agronomic Efficiency Index (AEI) data differ from results obtained in other studies by [15]
203 and [21], which verified high BM efficiencies when compared to the control (soluble
204 phosphate) treatment. Compared with Arad phosphate and triple superphosphate, [15]
205 verified an increase in the availability of P in the soil and in the production of sugarcane in
206 the use of BM. However, in the present study low AEI was observed using this source of P.

207

208 The low efficiency can be related to the pH considered high in the application of bone meal
209 on the substrate, preventing the transformation of the bound Ca + P compounds from BM to
210 available soil phosphorus [14]. Also, according to [6], raising the pH can reduce BM
211 efficiency and increase SS efficiency.

212 Like the soluble sources that undergo acidic attacks in reaction with water, like 2% citric acid
213 on triple superphosphate, the action of the acidic extractors is pronounced in this work. With
214 the exception of 0.5% Hydrochloric Acid, the other extractors showed AEI above 60%,
215 considered high by [22] when compared to the standard source, evidencing its potential use
216 in increasing the solubility of natural phosphate. These same authors, when studying the
217 matter production of pigeon pea (*Cajanus cajan*), obtained similar responses between the
218 use of natural phosphate and triple superphosphate.

219 Due to the high availability of phosphorus from the soluble source (SS), high phosphorus
220 conversion efficiency (PCE) was obtained, a result also observed by [23], which obtained a
221 superiority of 31% of the same source in relation to the use of a natural phosphate
222 (thermophosphate) in millet. However, the application of the acids resulted in satisfactory
223 phosphorus efficiencies in dry matter, when observed the behavior without its presence. The
224 PCE is directly correlated with the capacity of the phosphate source to release the nutrient in
225 the soil solution [23, 6]. Therefore, the low capacity observed by the bone meal
226 compromises the use of the nutrient by the forage.

227 Another fact intrinsic to PCE is the extent and efficiency of root system absorption, which
228 responds to the available P in the soil. The distinction in the morphology and root physiology

229 modifies the uptake of P and other nutrients, compromising PCE [23]. The difference
230 mentioned can be observed in the distribution of roots of *B. ruziziensis* according to the
231 phosphate source and the acid treatment used (Figure 1).
232



233
234 **Figure 1. Root system of *B. ruziziensis* according to the treatments applied.**
235

236 According to [24], the increase in P availability in the soil allows root growth in depth and the
237 emergence of lateral roots in grasses, such as *B. ruziziensis*. Therefore, when the opposite
238 scenario is found, the result responds in the same way, as verified for Dry Matter of Roots
239 (RDM). The observed similarity between the control and the BM evidences the low supply of
240 available phosphorus, resulting in poor root architecture (Figure 1). All acidic treatments
241 provided an increase in weight and volume and in the presence of *B. ruziziensis* root hairs,
242 both related to the increased availability of phosphorus in plants.

243 Observing the behavior of the root: aerial part, the acidification of BM provided a balance
244 between what was produced in the aerial part and in the roots. This fact is reported by [25],
245 arguing that values close to one represent balance between the photosynthetic rate
246 benefited by adequate nutrition and root expansion promoted by the presence of nutrients in
247 the soil.

248 **4. CONCLUSION**

249
250 The Shoot Dry Matter and Root Dry Matter from *B. ruziziensis* were positively influenced by
251 bone meal treatment with acids. Other than acidification with HCl 0.5%, the other treatments
252 provided AEI's above 60%. Acid treatments provided higher PCE's in the absence of
253 acidification.

254 **CONFLICT OF INTERESTS**

255
256
257
258 Authors have declared that no competing interests exist.

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

REFERENCES

1. IBGE. Pesquisa da pecuária municipal; 2018. (Accessed 17 June 2018). Available: <https://sidra.ibge.gov.br/pesquisa/ppm/quadros/brasil/2016>.
2. Dias-Filho MB. Degradação de Pastagens: o que é e como evitar. Belém: Embrapa; 2017.
3. Costa NL, Moraes A, Carvalho PCF, Monteiro ALG, Motta ACV, Oliveira RA. Dinâmica de crescimento e produtividade de forragem de *Trachypogon plumosus* sob níveis de correção da fertilidade do solo e idades de rebrota. *Ciência Animal Brasileira*. 2016;17(2):175-184. Portuguese
4. Ferraz JBS, Felício PED. Production systems - an example from Brazil. *Meat Science*. 2010;84(2):238-242.
5. Ferreira AVL, Ferreira E, Cavali J, Porto MO, Stachiw R. Farinha de Ossos. *Revista Brasileira de Ciência da Amazônia*. 2014;3(1):29-36. Portuguese
6. Damaceno JBD, Ferreira E, Oliveira DM, Guimarães RS, Gama RT, Padilha FJ. Produção de biomassa de *Brachiaria ruziziensis* adubada com farinha de ossos sob tratamentos ácidos. *Revista Agrogeoambiental*. 2018;10(1):83-93. Portuguese
7. Magalhães AF, Pires AJV, Carvalho GGP, Silva FF, Sousa CMV. Influência do nitrogênio e do fósforo na produção do capim-braquiária. *Revista Brasileira de Zootecnia*. 2007;36(5):1240-1246. Portuguese
8. Oelkers EH, Valsami JE. Phosphate mineral reactivity and global sustainability. *Elements*. 2008;4(2):83-87.
9. Lapido-Loureiro FE, Melamed R, Figueiredo Neto J. Fertilizantes: Agroindústria e Sustentabilidade. Rio de Janeiro: Centro de Tecnologia Mineral; 2009.
10. Porto EMV, Alves DD, Vitor CMT, Gomes VM, Silva MF, David AMSS. Rendimento Forrageiro da *Brachiaria brizantha* cv. Marandu submetida a doses crescentes de fósforo. *Scientia Agrária Paranaensis*. 2012;11(3):25-34. Portuguese
11. Avelar AC, Ferreira WM, Brito W, Menezes MABC. Composição mineral de fosfatos, calcário e farinha de ossos usados na Agropecuária Brasileira. *Archivos de Zootecnia*. 2009;58(224):737-740. Portuguese
12. Mattar EPL, Júnior EFF, Oliveira E. Caracterização físico-química de cinza de osso bovino para avaliação do seu potencial uso agrícola. *Pesquisa Agropecuária Tropical*. 2014;1(1):65-70. Portuguese
13. Miyahara RY, Gouvêa D, Toffoli SM. Obtenção e caracterização de cinza de ossos bovinos visando à fabricação de porcelanas de ossos - bone china. *Cerâmica*. 2007;53(327):234-239. Portuguese
14. Caione G, Fernandes FM, Lange, A. Efeito residual de fontes de fósforo nos atributos químicos do solo, nutrição e produtividade de biomassa da cana-de-açúcar. *Revista Brasileira de Ciências Agrárias*. 2013;8(2):189-196. Portuguese

- 303 15. Balbino TGM, Reinicke F, Modes K, Bezerra IL, Dias JRM, Ferreira E. Farinha de ossos
304 na produção capim Tifton. Revista Brasileira de Ciência da Amazônia. 2012;1(1):182-186.
305 Portuguese
- 306 16. Peel MC, Finlayson BL, McMahon TA. Koppen-Geiger, update world map of the Köp-
307 pen-geiger climate classification. Hydrology and Earth System Science. 2007;1(11):1633-
308 1644
- 309 17. Embrapa. Sistema brasileiro de classificação de solo. 2th ed. Brasília: Embrapa; 2013.
- 310 18. Prado GR, Silva LS, Prochnow LI, Griebeler G, Pocojeski E, Moro VJ. Comportamento
311 de superfosfato simples contendo fosfato de ferro de baixa solubilidade em água em solos
312 de várzea do Rio Grande do Sul. Revista Brasileira de Ciência do Solo. 2011;35(3):907-914.
313 Portuguese
- 314 19. Simões AC, Cruz IV, Cruz CV, Souza KG, Souza EFM, Dias JRM, Ferreira E. Meat and
315 bone meals in agronomy performance of tifton grass. International Journal of Agriculture and
316 Forestry. 2012;2(2):78-83.
- 317 20. Oliveira SB, Caione G, Camargo MF, Oliveira ANB, Santana L. Fontes de fósforo no
318 estabelecimento e produtividade de forrageiras na região de alta floresta – MT. Global
319 Science and Technology. 2012;5(1):1-10. Portuguese
- 320 21. Farias LN, Silva EMB, Guimarães SL, Souza ACP, Silva TJA, Schlichting AF.
321 Concentration of nutrients and chlorophyll index in pigeon pea fertilized with rock phosphate
322 and liming in cerrado oxisol. African Journal of Agricultural Research. 2015;10(14):1743-
323 1750.
- 324 22. Silva AG, França AFS, Miyagi ES, Dambros CE, Lopes FB. Eficiência da fertilização
325 fosfatada e nitrogenada em cultivares de milheto. Ciência Animal Brasileira. 2014;15(2):119-
326 127. Portuguese
- 327 23. Zambrosi FCB, Mattos Júnior D, Furlani PR, Quaggio JA Boaretto RM. Eficiência de
328 absorção e utilização de fósforo em porta enxertos cítricos. Revista Brasileira de Ciência do
329 Solo. 2012;36(2):435-496. Portuguese
- 330 24. Singh SK, Badgujar G, Reddy VR, Fleisher DH, Bunce JA. Carbon dioxide diffusion
331 across stomata and mesophyll and photobiochemical processes as affected by growth CO₂
332 and phosphorus nutrition in cotton. Journal of Plant Physiology. 2013;170(9):801-813.
- 333 25. Taiz L, Zeiger E. Fisiologia Vegetal. 5th ed. Porto Alegre: Artmed; 2013.
- 334
- 335
- 336
- 337