

1
2
3
4
5
6
7
8
9
10
11

Original Research Article

Liquid *Bacillus subtilis* Formulation in Rice in the Control of *Meloidogyne javanica* and Lettuce Development

ABSTRACT

Aim: To develop a liquid formulation based on *Bacillus subtilis*-34 using rice to evaluate the shelf life under refrigerated and room conditions and to evaluate the effect of different application times of the formulation on the control of *Meloidogyne javanica* and growth of lettuce plants.

Statistical Design: The design was completely randomized, with five treatments and eight replicates. The results were submitted to analysis of variance and the means compared by the Scott Knott test with 5% error probability.

Location and Duration of the experiment: The experiment was set up during the period from 02/13/2018 to 03/20/2018 in greenhouse located at the State University of Montes Claros, municipality of Janaúba, MG, Brazil.

Methodology: Treatments consisted of irrigation in the substrate of tubes at 8 and 15 days; irrigation in the substrate of tubes at 8 and 15 days and in pot at 25 and 35 days; irrigation in pot only at 25 and 35 days and 2 controls (Onix[®] and absolute control). All pots with plants were infested with 5000 nematode eggs. At 45 days of transplanting, the following nematological variables were evaluated: number of galls, number of egg mass, number of eggs per gram and reproduction factor, and agronomic variables fresh and dry biomass.

Results: Applications in the tube and pot and in the pot only were efficient for the reduction in the reproduction of *M. javanica* and for the development of lettuce plants.

Conclusion: *B. subtilis*-34 remains viable until 9 months in formulation stored under refrigerator and up to 7 months under room conditions.

12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29

Keywords: *Lactuca sativa* L; rhizobacteria; root-knot nematode, technology; shelf life

1. INTRODUCTION

Lettuce (*Lactuca sativa* L.) is a vegetable economically important for Brazil, being cultivated in almost all regions of the country [1]. Lettuce is a good source of fiber, vitamins, especially B, A, C and K, low in calories and a rich source of pigments beneficial to human health [2]. In addition to the nutritional aspect, it is also a culture of great importance from the social point of view, being cultivated mainly by family farmers near large urban centers in the so-called "green belts" [3].

Lettuce has numerous phytosanitary problems, among which phytonematodes stand out. Nematodes of the genus *Meloidogyne* are considered as limiting the commercial cultivation of several vegetables, since they have short cycle and are always cultivated in the same area, favoring the increase in the nematode population. Losses caused by phytonematodes

30 in vegetable crops are estimated at 12.3% in developed countries and 14.6% in developing
31 countries [4]. Lettuce plants, attacked by nematodes, are less developed due to dense
32 formation of galls in the root system and their control is a difficult task.

33

34 Several strategies have been used for the control of nematodes, among them: crop rotation,
35 solarization, use of resistant cultivars and chemical control. Most lettuce cultivars present
36 high susceptibility to *Meloidogyne* species [5]. The species of this genus most important to
37 lettuce and other leafy crops are *M. javanica* and *M. incognita* [6]. The damage caused by
38 agrochemicals, crop rotation infeasibility in small areas and the cost of plastic for solarization
39 evidences the necessity of the use of biological control agents, among them rhizobacteria
40 [7].

41

42 Among rhizobacteria, the genus *Bacillus* stands out, which has the capacity to produce
43 antibiotics, enzymes and toxins, which act directly, causing the mortality of juveniles and / or
44 indirectly affecting their behavior, feeding or reproduction. Plant-host recognition, resistance
45 induction and / or plant growth promotion processes can also be performed [8]. In addition,
46 *Bacillus* produces endospores, which are spores that survive under conditions of nutrient
47 deprivation and conditions of high temperature, which favors the maintenance of the viability
48 of formulations [9].

49

50 Promising results have been obtained in the control of *M. javanica* by *B. subtilis* in banana
51 and tomato crops [10]. *Bacillus* has been described for producing hydrolytic enzymes such
52 as lipases, chitinases and proteases capable of degrading structural components of
53 *Meloidogyne* [11, 12, 13].

54

55 The formulation of biocontrol agents such as rhizobacteria or another microorganism is an
56 essential step for their commercial use [14]. Formulations should improve shelf life since
57 biocontrol agents are living organisms. In addition, formulations should be economical and
58 contain sufficient number of viable colony forming units (CFUs) and be easily applied to soil
59 or plants [15]. Several farmers have multiplied bacteria from organic products on their
60 properties. Thus, the development of low-cost formulations for large-scale use is essential.

61

62 In this context, the aims of this work were to evaluate the application of a liquid formulation
63 of *B. subtilis* produced in rice broth and determine its efficiency in the control of *M. javanica*
64 and in the promotion of lettuce growth.

65

66 2. MATERIAL AND METHODS

67

68 2.1 Development of *Bacillus subtilis* formulation in rice broth and 69 establishment of the growth curve

70 *Bacillus subtilis*-34 isolate maintained in mineral water in glass tubes under room conditions
71 was used. From this suspension, a volume of 1.85 ml was collected. This suspension was
72 placed in 1 liter erlenmeyers and the following components were added in g.L⁻¹ of distilled
73 water: 185 g of raw rice, 185 g of sugar, 55.55 g of sodium chloride (NaCl), 46.29 g of
74 phosphate of potassium monobasic (KH₂PO₄). Chemical compounds, sugar, water and rice
75 were autoclaved at 1.0 atm at 120 ° C for 30 minutes. The formulation had final pH of 7 (±
76 0.2).

77

78 For determination of the initial number of CFUs, a one-milliliter aliquot of the liquid
79 formulation was submitted to a 10⁻⁵ dilution and 100 µL were collected for plating in Petri
80 dishes containing Tryptic Soy Agar (TSA). Petri dishes were incubated at 25°C for 24 hours,
81 when the initial number of CFU.mL⁻¹ was evaluated. For the establishment of the growth

82 curve, the formulation was incubated for 44 hours on an orbital shaker at 28°C at 220 rpm
83 and at 4-hour intervals, the same procedure was performed to determine the number of
84 CFUs. The growth curve of the bacteria was also determined in Tryptic Soy Broth (TSB)
85 medium, following the same methodology as that used for the liquid rice formulation.
86

87 **2.2 Evaluation of different application times of the liquid *Bacillus subtilis*** 88 **formulation in rice broth in the control of *Meloidogyne javanica* and in lettuce** 89 **development**

90
91 The experiment was carried out in greenhouse at the State University of Montes Claros,
92 Janaúba, MG, with the following geographical coordinates (43 ° 16'18.2 "W and 15 ° 49'51.5"
93 S) and average altitude of approximately 540 m a.s.l.). For the evaluation of growth
94 promotion and reduction of nematological variables, *B. subtilis*-34 bacterial isolate was used.
95 The formulation was made as reported in item 2.1.
96

97 The isolate growth was interrupted 28 hours after the beginning of incubation. At this time,
98 dilution at 10⁻⁵ and TSA plating was performed to determine the initial number of CFUs per
99 mL. To evaluate the survival period of the bacterium in the liquid formulation, a volume of 50
100 mL was kept at room temperature in the laboratory of phytopathology on the bench with
101 mean temperature of 26.05°C (maximum of 29.1°C and minimum of 23°C) and another 50
102 mL kept in refrigerator at 9°C. At one-month intervals for a period of 10 months, 10⁻⁵ dilution
103 and TSA plating were performed to determine the number of CFUs.
104

105 Lettuce seedlings 'Aurélia' cultivar were obtained from the sowing in styrofoam tubes
106 containing Bioplant[®] substrate. After 17 days, seedlings were transplanted to 3-liter pots
107 containing: substrate composed of soil (heavy clay, 26.6% silt, 60% clay, 13.4% sand, pH =
108 5) in 3:1 proportion, respectively, previously autoclaved at 1 atm and 120°C for 30 minutes,
109 three times at 24-hour intervals. The substrate was fertilized as recommended for the
110 culture. Prior to assay setup, the substrate was submitted to liming and incubated for 30
111 days.
112

113 The experiment was set up in a completely randomized design with five treatments and eight
114 replicates. Treatments consisted of: T1- irrigation of the formulation to the substrate in tubes
115 at 8 and 15 days, T2 - applied via irrigation to the substrate of tubes at 8 and 15 days and in
116 the pot at 25 and 35 days, T3- irrigation of the formulation to the soil of pots at 25 and 35
117 days after transplanting and 2 controls; T4-Onix[®] (Commercial product based on *Bacillus*
118 *methylothrophicus* - Isolated UFPEDA 20) and T5- without bacterium application and without
119 commercial product. In Onix[®] treatment, each plant received 250 ml of the commercial
120 product, previously diluted in water in the proportion of 4 mL.L⁻¹ one day after transplanting.
121 Regarding the rice formulation, the volume used by application in tubes and in pots was two
122 milliliters and 150 mL, respectively. At each application, a new formulation was produced.
123

124 The inoculation of *M. javanica* occurred 24 hours after the transplanting of seedlings to the
125 pot, each one receiving 5 ml of suspension containing 5000 eggs and eventual J2 calibrated
126 in Peters chamber, applied in three holes around the neck of each plant. At 45 days after
127 transplanting, the number of galls per gram of root (NG / g), egg mass per gram of root (MO
128 / g), number of eggs per gram of root (NO / g) were evaluated, as well as the reproductive
129 factor, calculated by the following formula: $FR = Pf. / Pi$, where Pf is the final nematode
130 population and Pi is the initial population applied to the plant [16]. and number of second
131 stage juveniles (J2). To count the number of egg masses, roots were immersed in floxin B
132 solution (150 mg.L⁻¹). The number of eggs was determined after root extraction [17, 18]. For
133 number of J2 in the soil, samples 200cm³ were processed [19]. Eggs and J2 of *M. javanica*
134 were quantified in Peters counting chamber in invert objective microscope.


135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155

The number of leaves, height, head diameter, fresh shoot mass, dry shoot mass and root weight were also evaluated. In order to determine the shoot dry mass, leaves were placed in paper bags, which were placed in a drying oven with forced air circulation at 65°C for 72 hours. Data were submitted to analysis of variance and means were compared by the Scott-Knott test at 5% error probability. Statistical analysis was performed using the "Sisvar" software [20].

3. RESULTS AND DISCUSSION

3.1 Development of *Bacillus subtilis*-34 formulation in rice broth and establishment of the growth curve

Figure 1 and 2 show the growth curves of *B. subtilis*-34 in the liquid rice formulation and TSB, respectively. In the liquid rice formulation, it was observed that the bacteria remained in the adaptation phase up to 20 hours after plating. After 24 hours, the exponential growth phase begins, culminating with higher number of CFU (6.14×10^8) at 32 hours after incubation. From 36 hours, the decline phase begins. The sudden drop in this value is justified by the depletion of nutrients in the culture medium and by the increase of toxic products from bacterial metabolism [21].



156
157
158
159
160
161
162
163
164
165
166

Figure 1. Growth curve of *Bacillus subtilis*-34 in liquid rice formulation.

In TSB, the highest number of CFUs occurred at 40 hours after incubation (6.41×10^7), that is, eight hours after occurrence in the liquid formulation and with a difference in the number of CFUs of 5.5×10^8 in relation to the liquid rice formulation (Figure 2). For the greenhouse experiment, the formulation was incubated for up to 28 hours because it was already in the logarithmic growth phase. It is important to highlight that 28 hours from the start of incubation, the liquid rice formulation provided CFU of 4.72×10^8 , while in the TSB medium, CFU was 3.49×10^7 .




Figure 2. Growth curve of *Bacillus subtilis*-34 in TSB medium.

167
168
169

170
171
172

3.2 Evaluation of different application times of the liquid *Bacillus subtilis* formulation in rice broth in the control of *Meloidogyne javanica* and lettuce development

173
174
175

Lettuce plants that received the liquid *B. subtilis*-34 formulation applied to the pot and tube + pot showed lower number of galls per gram of root and lower number of egg mass per gram of root (Table 1). The number of eggs per gram of root was significantly lower in all treatments that received the liquid *B. subtilis*-34 formulation compared to Onix[®] and control. The reproduction factor of *M. javanica* was lower in treatment irrigation applied to the tube followed by application in the pot and pot + tube. Application in tube reduced the reproduction factor of the nematode by 30.55% and 36.07% in relation to Onix[®] and control, respectively (Table 1). Positive *B. subtilis* results in reducing nematode populations, mainly of *Meloidogyne* species, in crops such as rice, tomato and banana have been observed in other studies [22, 23, 24]. There was no occurrence of juveniles of second stage in the soil in any of treatments, probably the J2 that hatched infected the roots again.

176
177
178
179
180
181
182
183
184
185
186
187
188

Table 1. Number of galls per gram root (NG/g), number egg mass per gram of root (EM / g), number of eggs per gram of root (NE / g) and reproduction factor (FR) of *Meloidogyne javanica* in lettuce submitted to application of *B. subtilis*-34 via liquid formulation at different times.

189
190
191
192
193

Treatments	NG/g	EM/g	NE/g	RF
Pot	8.00a	2.25a	1,686.52a	2.90b
Pot + Tube	8.12a	3.00a	1,562.07a	3.07b
Tube	21.75b	7.75b	2,173.50a	2.41a
Absolute control	36.50c	11.62b	4,784.71b	3.77c

Onix [®]	47.00c	9.62b	5,800.99b	3.47c
Coefficient of variation	41.85	61.54	40.00	15.37

194 Averages followed by the same letter in the column do not differ from each other by the
195 Scott-Knott test at 5% error probability.

196
197 *Bacillus* species interrupt the nematode life cycle through the production of toxic metabolites
198 that restrict their mobility, prevent juvenile hatching and penetration into plant roots [25].
199 Some authors have demonstrated reduction of the *M. incognita* population in tomato
200 inoculated with *Bacillus* species [26], others have observed that metabolites produced by *B.*
201 *subtilis* trigger hypersensitivity reactions in plant cells and affect oviposition, preventing
202 nematode females from obtaining sufficient energy to produce eggs [27].

203
204 The genus *Bacillus* is described as one of the main microbial groups capable of acting in the
205 control of phytopathogens through the synthesis of secondary metabolites, which, in general,
206 present a wide range of inhibition to different phytopathogen species [28]. *Bacillus* secretes
207 many secondary metabolites, including antibiotics, antifungals and siderophores. Metabolites
208 produced by *Bacillus* may also affect the microflora in the rhizosphere, providing an
209 environment antagonistic to pathogens, or may trigger host defense responses [29]. Cry
210 proteins produced by *Bacillus* species are toxic to nematodes, both of free-living and
211 phytoparasites, and the production of proteases by this group of bacteria have been
212 proposed as virulence factors in their pathogenesis against nematodes [30].

213
214 *Bacillus subtilis*-34 applied by irrigation and via pot + tube promoted higher number of
215 leaves, head diameter and dry shoot mass significantly higher than applications in tube,
216 control and Onix[®] commercial product (Table 2). Applications by pot and pot + tube
217 increased the number of leaves compared to control by about 80.02 and 73.56%. In relation
218 to Onix[®], the increase was 83.52 and 76.94%. Variable head diameter increased 81.98 and
219 75.42% when applying the formulation in pot and pot + tube, respectively, in relation to
220 control; when compared with Onix[®], the increase was 94 and 87.73%. In variable dry shoot
221 mass, considering applications in pot and pot + tube, the increase was 21.74 and 17.94%
222 in relation to control, and in relation to Onix[®], the increase was 23.38 and 19.54%.

223
224 The height and fresh shoot mass of lettuce plants that received *B. subtilis*-34 in the pot were
225 significantly higher than the other treatments, with an increase of about 119 and 322.22%,
226 respectively, in relation to control. On the other hand, application via pot and pot + tube
227 provided an increase of 98.61 and 155% of root weight in relation to control and Onix[®]
228 commercial product. The *in situ* effect by exposure of *B. subtilis* living cells may also lead to
229 an increase in plant biometry [31], reflecting productivity gains, with the bacterium being
230 commercially used for both purposes [32, 33].

231
232 **Table 2. Number of leaves (NL), height (H), head diameter (HD), fresh shoot**
233 **matter (FSM) (g), shoot dry matter (SDM) (g) and fresh root matter (FRM) (g) of**
234 **lettuce infected by *Meloidogyne javanica* and submitted to application of**
235 ***Bacillus subtilis*-34 via liquid rice formulation at different times.**

Treatments	NL	H (cm)	HD(cm)	FSM(g)	SDM(g)	FRM(g)
Pot	34.87a	7.12a	41.62a	76.00a	11.87a	8.62b
Pot + Tube	33.62a	6.25b	40.12a	64.12b	11.50a	10.50a
Tube	26.25b	5.00c	28.12b	23.50c	9.87b	6.25c
Absolute control	19.37c	3.25c	22.87c	18.00c	9.75b	4.37d

236

Onix [®]	19.00c	3.62c	21.37c	17.12c	9.62b	3.37d
Coefficient of variation	11.39	15.32	5.53	22.235	9.76	25.41

237 Averages followed by the same letter in the column do not differ from each other by the
 238 Scott-Knott test at 5% error probability.

239

240 *Bacillus subtilis* has been commercially used for the biocontrol of plant diseases and to
 241 increase crop yields [32, 33]. *B. subtilis* (PRBS-1) applied to tomato plants reduced the
 242 reproduction of root-knot nematode and promoted the growth of plants under greenhouse
 243 conditions [34]. Nemathel[®] treated banana seedlings reduced reproduction of *Radopholus*
 244 *similis*, *Meloidogyne* spp., *Pratylenchus* ssp. and *Helicotylenchus* spp. with efficiency similar
 245 to nematicide Carbofuran [35]. Tomato plants that received *B. subtilis* applications showed
 246 higher shoot growth, characterizing the bacterium as a plant growth promoter, and this effect
 247 may be due, in part, to the production of plant phyto regulators in the rhizosphere [36].
 248

249 3.3 Evaluation of the viability of liquid *Bacillus subtilis* rice formulation under 250 room and refrigerator conditions.

251

252 The initial CFU was 4.72×10^8 . Over the storage period under both conditions, oscillations in
 253 the concentration of bacterial cells were observed, sometimes with higher and sometimes
 254 with smaller values (Figure 3). At 4 months, the number of CFU of formulation stored under
 255 room and refrigerator conditions was 12.5×10^8 and 6.3×10^8 , respectively. At 6 months,
 256 the number of viable cells was similar in both storage conditions 7.7×10^8 at room temperature
 257 and 6.7×10^8 at refrigerator temperature. At 7 months, it was verified that the number of
 258 CFUs remains approximately constant 7.8×10^8 under room conditions, while under
 259 refrigeration conditions, reduction to 3.8×10^8 UFC was observed, and this value remains
 260 approximately constant until 9 months.


261

262 From 8 months, the number of CFUs stored under room conditions becomes minimal, while
 263 in refrigerator, the number of CFUs is approximately 4×10^8 at 9 months. The refrigerated
 264 environment extended the "shelf life" of the bacterium in two months. Low temperatures are
 265 generally used to preserve microorganisms by ensuring metabolism at low activity and
 266 preventing contamination with other microorganisms from affecting the stability of the
 267 biological control microorganism [37].
 268

269 A biological control product to be economically viable needs to have minimum concentration
 270 of 1×10^8 CFU / mL with 85% viability [38], which was achieved by the liquid *B. subtilis*-34
 271 formulation stored under room conditions for up to seven months and under refrigerator
 272 conditions for up to nine months.
 273

274

275



276 **Figure 3. Number of CFU of *Bacillus subtilis*-34 in liquid formulation stored under**
277 **room and refrigerator conditions for twelve months.**
278

279 The results verified in the nematode control and in the promotion of lettuce growth
280 demonstrate that the liquid bacterium formulation was effective despite the lower initial
281 number of CFU.mL⁻¹ (4.72x10⁸) compared to Onix, which had 1x10⁹ CFU. It also presented
282 lower cost when compared to TSB synthetic culture medium since US\$ 128.00 are
283 necessary for the production of one liter of TSB, whereas the liquid rice formulation requires
284 only US\$ 11.64.
285

286 **4. CONCLUSION**
287

288 Higher promotion of lettuce growth and control of *Meloidogyne javanica* was obtained by
289 applying the liquid *Bacillus subtilis*-34 formulation twice to the soil in the pot. *Bacillus*
290 *subtilis*-34 remains viable until nine months in formulation stored under refrigerator
291 conditions and up to seven months under room conditions in northern state of Minas Gerais.
292
293

294 **REFERENCES**
295

- 296 1. Carvalho Filho JLS de, Gomes LAA, Maluf WR. Tolerância ao florescimento precoce e
297 características comerciais de progênies F4 de alface do cruzamento Regina 71 x Salinas
298 88. Acta Scientiarum Agronomy.2009; 31: 37-42. Portuguese.
- 299 2. Matraszek R, Nowak BH, Chwil S, Chwil M. Macroelemental composition of cadmium
300 stressed lettuce plants grown under conditions of intensive sulphur nutrition.2016; 180:24-
301 34.
- 302 3. Villas Bôas RL, Passos JC, Fernandes M, Büll LT, Cezar VRS, Goto R. Efeito de doses e
303 tipos de compostos orgânicos na produção de alface em dois solos sob ambiente protegido.
304 Horticultura Brasileira.2004; 22:28-34. Portuguese.
- 305 4. Anwar SA, Mckenry MV. Incidence and population density of plant parasitic nematodes
306 infecting vegetable crops and associated yield losses. Pakistan Journal of Zoology.2012; 44:
307 327-333.
- 308 5. Fiorini CVA, Gomes LAA, Libânio RA, Maluf WR, Campos VP, Licursi V, et al.
309 Identificação de famílias F2:3 de alface homozigotas resistentes aos nematoides das
310 galhas. Horticultura Brasileira. 2007; 25: 509-513. Portuguese.
- 311 6. Pinheiro JB, Pereira RB, Carvalho ADF, Rodrigues CS, Suinada FA. Manejo de
312 nematoides na cultura do alface. 1ed. Embrapa: Brasília .2013. Portuguese.
- 313 7. Killani AS, Abaidoo RC, Akintokun AK, Abiala MA. Antagonistic Effect of Indigenous
314 *Bacillus subtilis* on Root-/Soil-borne Fungal Pathogens of Cowpea. Researcher.2011; 3:11-
315 18.
- 316 8. Tian B, Yang J, Lian L, Wang C, Li N, Zhang KQ. Role of an extracellular neutral protease
317 in infection against nematodes by *Brevibacillus laterosporus* strain G4. Applied Microbiology
318 and Biotechnology.2007; 74: 372-380.
- 319 9. Formstone A, Carballido-Lopez P, Noirot J, Errington D, Scheffers J. Localization and
320 interaction of teichoic acid synthetic enzymes in *Bacillus subtilis*. Journal
321 Bacteriology.2008; 190:1812-1821.
- 322 10. Ribeiro RCF, Campos VP, Xavier AA, Rocha LS, Ribeiro HB, Aguiar FM, et al.
323 Rizobactérias no controle de *Meloidogyne javanica* e mal do Panamá em bananeira.
324 Nematropica. 2012; 42: 218-226. Portuguese.
- 325 11. Sharma A, Meena KR, Kanwar SS. Molecular characterization and bioinformatics studies
326 of a lipase from *Bacillus thermoamylovorans* BHK67. International Journal of Biological
327 Macromolecules. 2018; 107:2131-2140.

- 328 12. Lee YS, Kim KY. Antagonistic potential of *Bacillus pumilus* L1 against root-Knot
329 nematode, *Meloidogyne arenaria*. Journal Phytopathology.2016; 164:29-39.
- 330 13. Zhang L, Yu J, Xie Y, Lin H, Huang Z, Xu L, et al. Biological activity of *Bacillus*
331 *thuringiensis* (Bacillales: Bacillaceae) chitinase against *Caenorhabditis elegans* (Rhabditida:
332 Rhabditidae). Journal of Economic Entomology. 2014; 107:551-558.
- 333 14. Berninger T, López OG, Bejarano A, Preininger C, Sessitsch A. Maintenance and
334 assessment of cell viability in formulation of no-sporulating bacterial inoculants. Microbial
335 Biotechnology. 2018; 11:277–301.
- 336 15. Sharma RR, SINGH D, Singh R. Biological controlo f postharvest diseases of fruits and
337 vegetables by microbial antagonists: A review. Biological Control. 2009; 50:205-211.
- 338 16. Oostenbrink M. Major characteristic of the relation between nematodes and plants.
339 Mededlingen van de Landbouwhogeschool;1966.
- 340 17. Hussey RS, Barker KA. A comparison of methods of collecting inocula for *Meloidogyne*
341 spp. Including a new technique. Plant Disease Reporter. 1973;57:1025-1028.
- 342 18. Boneti JIS, Ferraz S. Modificação do método de Hussey e Barker para extração de ovos
343 de *Meloidogyne exigua* de cafeeiro. Fitopatologia Brasileira.1981; 6: 553. Portuguese.
- 344 19. JENKINS WR A rapid centrifugal-flotation technique for separating nematodes from soil.
345 Plant Disease Reporter.1964; 48: 692.
- 346 20. Ferreira DF. SISVAR: Um programa para análises e ensino de estatística. Revista
347 Symposium. 2008; 6: 36-41.Portuguese.
- 348 21. Instituto Superior de Ciências da Saúde do Sul. Manual Prático de microbiologia. 2018.
349 Acessado em 07 de julho de 2018.
350 Disponível:<http://www.codigopostal.cibeforma.pt/dir/0/instituto-superior-de-ciencias-da-saude-sul/>.
- 351
- 352 22. Ludwigi J, Mourão AB, Gomes CB. Potencial da microbiolização de sementes de arroz
353 com rizobactérias para o biocontrole do nematoide das galhas. Tropical Plant
354 Pathology.2013; 38: 264-268. Portuguese.
- 355 23. Fernandes RH, Vieira BS, Fuga CAG, Lopes EA. *Pochonia chlamydosporia* e *Bacillus*
356 *subtilis* no controle de *Meloidogyne incognita* e *M. javanica* em mudas de tomateiro.
357 Bioscience Journal. 2014; 30:194-200. Portuguese.
- 358 24. Araújo JJ S, Muniz MFS, Filho GM, Rocha FS, Castro JMC. *Bacillus subtilis* no
359 tratamento de mudas de bananeira infectadas por fitonematoides. Revista Ceres. 2018; 65:
360 99-103. 2018. Portuguese.
- 361 25. Kavitha J, Jonathan EI, Umamaheswari R. Field application of *Pseudomonas*
362 *fluorescens*, *Bacillus subtilis* and *Trichoderma viride* for the control of *Meloidogyne incognita*
363 (Kofoid and White) Chitwood on sugarbeet. Journal of Biological Control. 2007; 21:211-215.
- 364 26. Silva JO, Santana MV, Freire LL, Ferreira BS, Rocha MR. Biocontrol agents in the
365 management of *Meloidogyne incognita* in tomato. Ciência Rural.2017; 47:1-7.
- 366 27. Sharma RD, Gomes AC. Effects of *Bacillus* spp. Toxins on oviposition and juvenile
367 hatching of *Heterodera glycines*. Nematologia Brasileira. 1996; 20: 53-62.
- 368 28. Sansinenea E, Ortiz A. Secondary metabolites of soil *Bacillus* spp. Biotechnology
369 Letters. 2011; 33: 1523–1538.
- 370 29. Velusamy P, Gnanamnickam SS. The Effect of bacterial secondary metabolites on
371 bacterial and fungal pathogens of rice. Em: KARLOVSKY, P, editor. Secondary Metabolites
372 in Soil Ecology .14ª ed. Berlin: Springer; 2008.
- 373 30. Lian LH, Tian BY, Xiong R, Zhu MZ, Xul J, Zhang KQ. Proteases from *Bacillus*: a new
374 insight into the mechanism of action for rhizobacterial suppression of nematode populations.
375 Letters in Applied Microbiology. 2007; 45: 262–269.
- 376 31. Hammami I, Rhouma A, Jaodu B, Rebai A, Nesme X. Optimization and biochemical
377 characterization of a bacteriocin from a newly isolated *Bacillus subtilis* strain 14 B for
378 bioncontrol of *Agrobacterium* spp. strains. Letters in Applied Microbiology. 2009; 48: 253-
379 260.

- 380 32. Ngugi HK, Dedej S, Delaplane KS, Savelle AT, Scherm H. Effect of flower-applied
381 Serenade biofungicide (*Bacillus subtilis*) on pollination-related variables in rabbiteye
382 blueberry. *Biological Control*. 2005; 33: 32-38.
- 383 33. Yao BH, Karimov S, Boturov U, Sanginboy S, Sharipov A. Effect of FZB 24R *Bacillus*
384 *subtilis* as a biofertilizer on cotton yields in field tests. *Archives of Phytopathology and Plant*
385 *Protection*. 2006; 39: 323-328.
- 386 34. Araújo FF, Marchesii GVP. Uso de *Bacillus subtilis* no controle da meloidoginose e na
387 promoção do crescimento do tomateiro. *Ciência Rural*. 2009; 39: 1558-1561. Portuguese.
- 388 35. Araújo JJ S, Muniz MFS, Filho GM, Rocha FS, Castro JMC. *Bacillus subtilis* no
389 tratamento de mudas de bananeira infectadas por fitonematoides. *Revista Ceres*. 2018; 65:
390 99-103. 2018. Portuguese.
- 391 36. Araujo FF, Bettiol W. Supressividade dos nematoides *Meloidogyne javanica* e
392 *Heterodera glycines* em soja por adição de lodo de esgoto ao solo. *Ciência Rural*. 2005; 35:
393 806-812. Portuguese.
- 394 37. Alves LFA, Alves SB, Lopes RB, Augus NT. Estabilidade de uma formulação de *Bacillus*
395 *sphaericus* armazenada sob diferentes temperaturas. *Scientia Agricola*. 2001; 58: 21-26.
396 Portuguese.
- 397 38. D'agostino F, Morandi MAB. Análise da viabilidade comercial de produtos à base de
398 *Bacillus subtilis* e *Bacillus pumilus* para controle de fitopatógenos no Brasil. EM: Bettiol W,
399 Morandi MAB, editores. *Biocontrole de doenças de plantas: Uso e perspectivas*. 1ª ed.
400 Jaguariúna: Embrapa Meio Ambiente. 2009. Portuguese