

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32

34

5
6
7
8
9
10
11
12
13
14

15
16
17
18
19
20
21
22
23
24

25

26

27

28
29
30
31
32

33 Nigeria where it is grown as sole crop or as an intercrop with other crops such as root crops, grain
34 legumes, cereals (sorghum, millet, rice) and even vegetables (Gwinner *et al.*, 1996). Despite the
35 importance and wide cultivation of this cereal, its growth and productions are usually impaired by
36 diseases caused by fungal, bacterial, viral or nematode which results in considerable yield loss and
37 decrease grain quality (Akande and Lamidi, 2006).

38 Fungi were ranked as the second most important cause of maize diseases and the major genera
39 commonly encountered on maize in tropical regions are *Fusarium*, *Aspergillus* and *Penicillium*
40 (Ominski *et al.*, 1994; Orsi *et al.*, 2000). *Fusarium* is considered as the most devastating fungal
41 menace of maize while *Fusarium verticillioides* is the prevalent specie causing root rot, stalk rot and
42 ear rots in maize (Masuka *et al.* 2003; Alankoya *et al.*, 2008). Meanwhile, the severe economic
43 losses in corn and other cereal crops worldwide, in addition to the potential occurrence of
44 fumonisins and other mycotoxins in consequence of *F. verticillioides* infection are a matter of
45 concern in current mycotoxicology (Ono *et al.* 2010).

46 In the recent times, the use of fungicide which is a chemical control measures commonly employed
47 in the management of diseases caused by fungi have been discouraged due to its environmental and
48 food contamination (Akanmu *et al.* 2013). Hence, there is need to embrace a safer and novel
49 biological approach to plant disease management. In order to improve plants' resistance to diseases
50 effort has been directed at the search for new anti-microbial materials from natural sources, which
51 are mostly low-molecular weight secondary metabolites essential for plant disease resistance (Singh
52 *et al.*, 2010; Maddox *et al.* 2010). While accumulation of certain plant secondary metabolites such as
53 phytoalexins is induced upon pathogen attack, the increase accumulation of phenolic phytoalexins in
54 plants can promote host defense against pathogens (Boudet, 2006; Maddox *et al.* 2010). More so,
55 phenolic compounds with less complex structures, such as catechol and coumarin, have exhibited
56 bactericidal and fungicidal activities (Cowan, 1999). Hence, the ability of plants to release phenol as
57 a defense mechanism against infection by pathogens is of importance in determining its resistance
58 (Samapundo, 2007). This study therefore investigates the phenol production by five maize varieties
59 in response to infection caused by *Fusarium verticillioides*

MATERIALS AND METHODS

Experimental Site: The experiment was conducted in the screen house of the Department of Botany University of Ibadan, Ibadan, Nigeria.

Sources of materials used: Seeds of maize varieties; SWAM1-SR, BR-9928-DMR-SR, ART-98-SW1, ART-98-SW6, BR-9943-DR-SR were obtained from maize germplasm of the Institute of Agricultural and Research Training (IAR&T), Apata, Ibadan. Pure culture of the characterised *Fusarium verticillioides* isolate was obtained from the Plant Pathology Laboratory, International Institute of Tropical Agriculture (IITA), Ibadan, Nigeria.

Multiplication of the inoculum: *Fusarium verticillioides* isolate was sub-cultured, maintained on solidified potato dextrose agar (PDA) and incubated at room temperature (28 ± 2) °C for 7 days to produce multiple cultures of the organism. Each of the cultured plates was flooded with 2ml sterile distilled water. A sterile scalpel was used to harvest the spores and mycelia growths by scraping them into a sterilized flask. The solution was adjusted with sterile distilled water, stirred and filtered out the spore suspension with the use of muslin cloth. The inoculum load of 1.4×10^7 spores/ml was counted using haemocytometer.

Soil preparation and planting: Prior to the conduct of the experiment, the screenhouse was adequately sanitized. Agricultural soil of between 0 to 15 cm depth was collected from the experimental site of Botany Department, University of Ibadan, Ibadan. The soil was sterilized using electric soil sterilizer at 120°C for 1 hour, and was filled in polythene bag at 5kg per bag after cooling. Maize seeds were surface sterilized in a beaker containing sodium hypochlorite (3.5% w/v) for 3 minutes after which the seeds were dried in between layers of whatmann filter paper for another 5 minutes. The seeds were planted at three seed per hole across the varieties.

Experimental Design: The experiment was laid out in a completely randomized design (CRD) in ten replications. Treatments across the five maize varieties consisted of two inoculation methods; soil and seedling inoculations. These were carried out at varying volume of the standardized (1.4×10^7 spores/ml) inoculum at 0 ml, 10 ml and 20 ml of the spore suspensions. The treatment

with soil inoculation was conducted at the period of planting while seedling inoculation was carried out on the respective plots at 2 weeks after planting. The control experiments were treated with respective volume of sterile distilled water.

Determination of disease incidence and severity: The percentage of disease incidence and severity were determined using the formulas:

$$\text{Disease incidence (\%)} = \frac{\text{number of infected maize plants}}{\text{number of maize plants}} \times 100$$

$$\text{Disease severity (\%)} = \frac{\text{area of plant tissue affected}}{\text{total area}} \times 100$$

Extraction of plant samples: Two replicates from each treatment across the varieties were carefully uprooted at 4th, 5th and 6th week after planting (WAP). Soil adhering to the root of maize plant was removed by agitation in water, the plants were then air dried at room temperature. Two gramme (2g) of leaf sample per treatments was treated with 10ml of 80% methanol in a tightly clogged flask. The treated leaf was allowed to stand for 3 minutes after which the liquid fraction was filtered using muslin cloth. The step was repeated three times with 80% ethanol and the supernatants were collected into volumetric flasks. Final volume of the extracts was made to 50 ml with 80% ethanol and all the extracts from each sample was combined and centrifuged at 2000 revolutions per minutes for 20 minutes in a centrifuge.

Determination of phenol content: This was done using the Folin-Ciocalteu method as described by Singh *et al.* (2012) in which gallic acid served as a standard and total phenol is expressed as mg/g gallic acid equivalents (GAE). The amount of total soluble phenol present in the sample was calculated according to Singh *et al.* (2012).

Data collection and statistical analysis: Data were collected on plant height (cm), leaf number (cm), disease incidence (%) and disease severity (%) weekly for 9 weeks. All the data on growth, disease and phenol contents collected were subjected to ANOVA using SAS 9.1 statistical analysis software and Means were separated by Duncan Multiple Range System at 95% confidence interval.

RESULTS

The fitted model for the interactive effect of maize varieties, inoculation methods, inoculums' quantity and duration of experiment produced a significant ($p<0.01$) result on the growth and disease parameters measured. All the factors evaluated were significant with plant height, number of leaves and disease severity while only inoculum volume and duration of the experiment (WAP) was significant with disease incidence caused by *F. verticillioides* (Table 1).

Maize variety; BR-9928-DMR-SR followed by ART-98-SW6 recorded the most significant ($p<0.05$) growth on plant height and number of leaves across the varieties evaluated. This was followed by ART-98- SW1 and BR-9943-DR-SR which also recorded similar level of significance while SWAM 1-SR showed the least growth performances (Table 2).

All the maize varieties recorded more than 70% stalk rot infections with BR-9928-DMR-SR having highest rate of 73.11%. The stalk rot severity ranged from 19.01% in variety ART-98-SW1 to 25.21% in ART-98-SW6 (Figure 1).

Table 3 showed the effect of different inoculation methods and inoculums volume on the growth and disease occurrence in maize plants. While no significant differences was recorded between the inoculation methods with respect to plant height and number of leaves, soil inoculation method produced a significant ($p<0.05$) disease incidence and severity compare to seedling inoculation which showed no significant difference from the control. Whereas, significant ($p<0.05$) reduction was recorded in the growth rate with increasing inoculums volume. No significant result was obtained in the disease incidence with respect to inoculums levels while at 20 ml, there was significant increase in stalk rot severity (Table 3).

There was a significant increase in plant height and leaf number with the increasing weeks after planting (Table 4)

Figure 2 showed a linear relationship between the disease incidence caused by *F. verticillioides* in maize and the period of experiment. A consistent increase in the disease progression was observed

140 until the 5th week when the infection reached its climax and maintained this position till the 9th week
141 of experiment (Figure 2).

142 The r^2 value of 0.8694 reinstated the reliability of result obtained in the severity of stalk rot with
143 respect to period of experiment. Having received *F. verticillioides* inoculation in the 2nd WAP,
144 disease severity increased consistently from 3rd week (9.95%) to 9th week (36.29%) after planting
145 (Figure 3).

146

UNDER PEER REVIEW

147 **Table 1:** ANOVA table of growth and disease occurrence in maize plants after inoculations with *Fusarium verticillioides*

Sources of variation	df	Plant height (cm)		Leaf number		Disease incidence		Disease severity	
		F value	P value	F value	P value	F value	P value	F value	P value
Model	21	162.46	0.0001**	157.48	0.0001**	210.05	0.0001**	72.74	0.0001**
Maize varieties	4	12.48	0.0001**	14.55	0.0001**	0.30	0.8765	8.82	0.0001**
Inoculums volume	2	43.49	0.0001**	39.33	0.0001**	11.00	0.0011**	34.64	0.0161*
Inoculation method	2	16.16	0.0001**	9.44	0.0001**	0.29	0.5882	5.81	0.0001**
Weeks After Planting	8	405.5	0.0001**	395.34	0.0001**	392.13	0.0001**	172.33	0.0001**
Replicates	9	1.56	0.1693	1.05	0.3864			0.42	0.8372
Error	1327								
Corrected Total	1348								


148

149 *Significant ** Highly Significant.

150

151

152



153

154

155 Figure 1: Disease incidence (DI) and severity (DS) across the maize varieties caused by *F.*
156 *verticillioides*.

157

158

159

160

161

162

163

164

165

166

167

168

169

170

Table 2: Effect of *F. verticillioides* on plant heights and leaf number across the maize varieties

Varieties	Plant height (cm)	Number of leaves
SWAM 1-SR	43.27 ^c	4.53 ^c
BR-9928-DMR-SR	51.31 ^a	5.47 ^a
ART-98-SW6	54.56 ^a	5.68 ^a
ART-98- SW1	49.52 ^b	5.07 ^b
BR-9943-DR-SR	48.94 ^b	4.97 ^b
R ²	0.72	0.71

Mean with different letters are significantly different ($p \leq 0.05$)

187 Table 3: Effect of inoculation method and inoculum volume on growth and disease occurrence in
188 maize plants

Parameters	Variables	Plant height (cm)	Number of leaves	Disease incidence (%)	Disease severity (%)
Inoculums' volume (ml)	Control	54.11a	5.63a	70.18a	21.51b
	10	48.88b	4.90b	71.78a	22.18ab
	20	41.58c	4.63c	72.65a	23.99a
Inoculation method	Control	52.99a	5.23a	69.56b	20.69b
	Seedling inoculation	46.72b	5.06a	70.18b	21.51b
	Soil inoculation	50.58b	5.17a	74.87a	25.59a
	R ²	0.72	0.71	0.94	0.53

189

190 Means with different letters are significantly different ($p \leq 0.05$)

191

192 Table 4: Effect of time (WAP) on the growth of treated maize plants


	Weeks After Planting	Plant height (cm)	Leaf number
	1	2.08i	0.01i
	2	11.78h	1.59h
	3	24.61g	3.07g
	4	39.63f	4.35f
	5	54.69e	5.61e
	6	66.92d	6.50d
	7	75.94c	7.43c
	8	82.57b	8.38b
	9	87.65a	9.36a
	R ²	0.72	0.71

193

194 Means with different letters in a column are significantly different

195

196



197


198 Figure 2: Effect of time (WAP) on disease incidence caused by *F. verticillioides* in maize plants.

199

200

201

202



203

204 **Figure 3:** Effect of time (WAP) on disease severity of *F. verticillioides* in maize plants

The model for the reaction of quantity of phenol produced was significant ($p<0.01$) through the period of experiment. The effect of maize varieties and inoculums volume also recorded significant result at the 4th, 5th and 6th (WAP). The inoculums' method used showed significance ($p<0.01$) at 4th and 5th WAP while it was significant at $p<0.05$ at 6th WAP (Table 5).

Maize variety ART-98-SW6 showed the most significant ($p<0.05$) phenol content across the period of this study. This result was followed by BR-9928-DMR-SR while ART-98-SW1 and BR-9943-DMR-SR showed no significant difference (Table 6).

The seedling inoculation method showed significantly higher phenol contents than soil inoculation, although the two methods produced results which were significantly higher than the control. Moreover, inoculums volume of 10ml recorded a significantly higher phenol contents than at 20ml (Table 7).

The inoculation method were positively and significantly ($p<0.01$) correlated with inoculums volume ($r=0.64$). Also, inoculation method was significantly associated with phenol production at week 4 ($r=0.60$), week 5 ($r=0.68$) and week 6 ($r=0.71$) after planting. Similarly, inoculums' volume was significantly correlated with phenol content obtained at week 4 ($r=0.81$), week 5 ($r=0.90$) and week 6 ($r=0.89$) of the experiment (Table 8).

Table 5: ANOVA table for Phenol (mg/g) produced by the maize plants after days of infection

Source	df	4th WAP		5th WAP		6th WAP	
		F value	P>F	F value	P>F	F value	P>F
Model	7	90.24	0.0001**	227.49	0.0001**	310.19	0.0001**
Maize varieties	4	49.92	0.0001**	67.34	0.0001**	74.98	0.0001**
Inoculation methods	1	9.59	0.0065**	9.79	0.0061**	4.21	0.0560*
Inoculums volume	1	139.13	0.0001**	306.88	0.0001**	267.68	0.0001**
Error	17						
Corrected Total	24						

**=Highly significant *= Significant

Table 6: Phenol (mg/g) content of the maize plants after days of infection

Varieties	4th WAP	5th WAP	6th WAP
SWAM1-BR-SR	7.40 ^d	11.12 ^d	14.04 ^d
BR-9928-DMR-SR	13.80 ^b	16.92 ^b	20.96 ^b
ART-98-SW6	16.16 ^a	19.04 ^a	21.28 ^a
ART-98-SW1	11.20 ^c	15.04 ^c	19.22 ^c
BR-9943-DMR-SR	10.52 ^c	14.54 ^c	18.74 ^c
R ²	0.97	0.99	0.99

Means with different letters are significantly different ($p \leq 0.05$)

Table 7: Phenol (mg/g) content of the maize plants days after inoculation with different volumes of *Fusarium verticillioides*

Parameter	Variable	4th WAP	5th WAP	6th WAP
Inoculation method	Control	4.72 ^c	5.94 ^c	7.88 ^c
	Seedling	14.32 ^a	16.33 ^a	17.58 ^a
	Soil	12.86 ^b	14.03 ^b	15.48 ^b
	R ²	0.97	0.99	0.99
Inoculums volume	Control	4.97 ^c	5.94 ^c	7.88 ^c
	10ml	13.37 ^a	15.32 ^a	18.61 ^a
	20ml	10.81 ^b	13.04 ^b	15.63 ^b
	R ²	0.97	0.99	0.99

Means with different letters are significantly different ($p \leq 0.05$).

233 **Table 8:** Extent of association between the maize varieties, pathogen inoculation, and time with
 234 phenol content

Correlation	Maize varieties	Inoculation method	Inoculums' volume	Phenol contents		
				Week 4	Week 5	Week 6
Maize varieties						
Inoculation method	0.00					
Inoculums' volume	0.00	0.64**				
Week 4	-0.13	0.60**	0.81**			
Week 5	-0.11	0.68**	0.90**	0.98**		
Week 6	-0.72	0.71**	0.89**	0.94**	0.99**	

236 * Significant, **Highly Significant

238 DISCUSSION

239 High incidence of stalk rot infection recorded across the *Fusarium verticillioides* inoculated maize
 240 varieties supported the claim that *F. verticillioides* is the major causal agent of highly variable
 241 disease symptoms ranging from asymptomatic plants to severe rotting and wilting in maize (Oren et
 242 al., 2003; Sobowale et al., 2005; Aliakbari, 2007). Virulence of *F. verticillioides* that ranged from
 243 25.21% in ART-98-SW6 to 19.01% in ART-98-SW1 demonstrated the variations in the severity of
 244 systemic infection on the host, while environmental factors also play a major in the activities of this
 245 cosmopolitan pathogen (Murillo-Williams and Munkvold, 2008; Thompson and Raizada, 2018).
 246 More so, variation recorded in the growth rate and resistance of maize varieties to *F. verticillioides*
 247 infections is in consonance some earlier findings in which host resistance was reported as a major
 248 determinant of variations in the activities of the pathogen (Sharma, 2003; Olowe et al. 2017).

249 Similar rate of maize growths observed in both seedling and soil inoculation methods is contrary to
 250 the results obtained by Khan et al. (2012) in which a significant increase in the growth parameters of
 251 plants that received *Alternaria inocula* through different methods was reported. Also, higher
 252 incidence and severity of stalk rot recorded in the soil inoculation method conforms to the earlier

claims that attributed varying severities caused by artificial inoculation of *F. verticillioides* with different inoculation methods (Drepper and Renfro, 1990; Sobowale, 2011). Furthermore, the increase in inoculums' quantity which resulted in corresponding decrease in growth rate of maize plants revealed varied efficiency as regards inoculum delivery with different inoculums' level (Sobowale et al., 2007).

The prevalence of stalk rot incidence observed to reach its peak at 5th week after planting affirmed that fungus colonizes maize stalks systemically without necessarily causing visible disease symptoms and that *F. verticillioides* strains can be vertically transmitted through seed-to-plant transmission and systemic stalk infection (Munkvold, 1997; Murillo-Williams and Munkvold, 2008). Meanwhile, the severity rate of 36.29% at 9th week after inoculation justifies the epidemiological claim that *Fusarium* stalk rot reduces output in maize by 10% typically and by 30–50% in severely affected areas (Li et al. 2010).

The result of maize variety ART-98-SW6, followed by BR-9928-DMR-SR, ART-98-SW1, BR-9943-DMR-SR and SWAM1-BR-SR as the decreasing order of phenol production after inoculated with *F. verticillioides* was in agreement with some earlier reports that both constitutive or induced synthesis contribute to abundance and composition of phenolic compounds in cereal grains and this is highly variable depending on the species, variety and environmental conditions (Adom and Liu 2002; Lattanzio et al. 2006). The observation of Reddy and Sireesha (2013) that nutritional status and concentration of biochemical constituents in plants prior to infection determines the severity of disease possibly explains the reasons maize varieties ART-98-SW6 and BR-9928-DMR-SR with higher stalk rot severities also produced higher phenol contents than other varieties tested. This further corroborated the reports that plants respond to pathogen invasion through the activation of complex defense strategies such as the accumulation of flavonoids, phytoalexins and phenolic compounds (Delledonne et al., 2001; Hefny et al., 2012). In this work, total phenol produced by the maize varieties acts as biochemical markers to analyse disease incidence and severity which conforms to the work done by Singh et al. (2012) on biochemical response and host-pathogen relationship of stalk rot fungi in early stages of maize (*Zea mays*).

Higher phenol contents produced in seedling inoculation method compare to soil inoculation could be associated with plants' reaction to direct inoculation which possibly resulted into a more effective delivery of *F. verticillioides* inoculums. Thus, the possibility of the pathogen inducing phenol production could be substantiated by the report that biochemical resistance, tolerance or susceptibility in plants against any disease depends mainly on preexisting, preformed or induced substances by the pathogen in the host (Reddy and Sireesha, 2013). In this study, the higher phenol production that occurred at lower inoculum volume was consistent with the findings of Perveen et al. (2010) who reported a decrease in total phenol of the leaves of *M. arvensis* with increase in initial inoculum of *S. sclerotiorum*, a situation attributed to the altered rates of synthetic activity because of infection by pathogen (Howlett, 2006).

The inoculation methods and inoculums volume used in this study effectively delivered infective stalk rot inoculum dosage, the rate of which were found strongly correlated with phenol production in maize plants at 4th, 5th and 6th week after inoculation. Thus, agreed with the claim that majority of phenolic compounds are a part of the preformed general defense system against potential pathogens (Stuper-Szablewska et al. 2017).

In general, phenol production in the maize varieties corresponds with the rate of severity of *F. verticillioides* infections. Therefore, close association exists between stalk rot disease and maize plants' resistance to infection through phenol production. Hence, researches that will increase phenol production in crops especially maize, may thus be suggested.

REFERENCES

- Adom KK, Liu RH. (2002). Antioxidant activity of grains. *J Agric Food Chem*. 50:6182–6187
- Akande S.R. and Lamidi G.O. (2006). Performance of quality protein maize varieties and disease reaction in the derived-savanna agro-ecology of South-West Nigeria. *African Journal of Biotechnology*. 5(19), 1744-1748.
- Akanmu A. O., Abiala M. A., Akanmu A.M., Adedeji A. D, Mudiaga P. M and Odebode A. C. (2013) Plant Extracts Abated Pathogenic *Fusarium* Species of Millet Seedlings. *Archives of Phytopathology And Plant Protection*, 46 (10) 1189-1205,

Alankoya A.E. Monda E.O. and Ajanga S. (2008). Variation in *in vitro* fumonisin B1 production by different *Fusarium verticillioides* isolates in Kenya. *American-Eurasian Journal for Agriculture and Environmental Science*. 4: 368-371.

Aliakbari F., Mirabolfathy M., Emami M., Mazhar S.F. and Karami-Osboo R., (2007). Natural Occurrence of *Fusarium* species in Maize Kernels at Gholestan Province in Northern Iran. *Asian Journal of Plant Sciences*, 6: 1276-1281.

Boudet A (2006) Evolution and current status of research in phenolic compounds. *Phytochemistry* 68:2722–2735

Cowan M (1999) Plant products as antimicrobial agents. *Clin Microbiol Rev* 12:564–582

Delledonne, M., J. Zeier, A. Marocco and C. Lamb (2001). Signal interactions between nitric oxide and reactive oxygen intermediates in the plant hypersensitive disease-resistance response. *Proc. Natl. Acad. Sci. USA.*, 98: 13454-13459.

Drepper W.J. and Renfro B.L. (1990). Comparison of Methods for Inoculation of Ears and Stalks of Maize with *Fusarium moniliforme*. *Plant Dis.* 74:952-956.

Gwinner J., Harnisch R., Much O. 1996. Manuel sur la manutention et la conservation des grains apres-recolte. GTZ, Eschborn, Germany. P 38.

Hefny M., Atta S., Bayoumi T., Ammar S.H. and El- Bramawy M. (2012). Breeding Maize for Resistance to Ear Rot Caused by *Fusarium moniliforme*. *Pakistan Journal of Biological Sciences*, 15: 78-84.

Howlett B.J. (2006) Secondary metabolite toxins and nutrition of plant pathogenic fungi *Cur. Opinion Plant Biol.*, 9, pp. 371-375

Khan M.M., Khan M.R. and Mohiddin F.A. (2012). The Relative Performance of Different Inoculation Methods with *Alternaria brassicae* and *A. brassicicola* on Indian Mustard. *Plant Pathology Journal*, 11: 93-98.

Lattanzio V, Lattanzio V.M., Cardinali A. (2006). Role of phenolics in the resistance mechanisms of plants against fungal pathogens and insects. *Phytochem: Adv Res.* 661:23–67.

Li W.J., He P and Jin J.Y. (2010) Effect of potassium on ultrastructure of maize stalk pith and young root and their relation to stalk rot resistance. *Agricultural Sciences in China*, 9:1467–1474.

Maddox C.E., Laur L.M. and Tian L. (2010) Antibacterial activity of phenolic compounds against the phytopathogen *Xylella fastidiosa*. *Curr Microbiol.* 60(1):53-58.

Masuka A.J., Cole D.L., and Mguni C. 2003. List of plant diseases in Zimbabwe. Plant Protection Research Institute.

Munkvold, G. P., McGee, D. C., and Carlton, W. M. (1997). Importance of different pathways for maize kernel infection by *Fusarium moniliforme*. *Phytopathology* 87:209-217

Murillo-Williams A. and Munkvold G.P. (2008) Systemic Infection by *Fusarium verticillioides* in Maize Plants Grown Under Three Temperature Regimes. *Plant Disease* 92 No. 12, pages 1695 - 1700

Okoro-Robinson M.O., Olawuyi O.J., Bello W.O. and Babalola B.J. 2014. Comparative evolution of organic manure on growth and yield of maize. *Agricultural and Biological Research*. 30(1): 60-73.

355 Olakojo S.A. and Iken J.E. 2001. Yield performance and stability of some improved maize (*Zea*
356 *mays* L.) varieties. *Moor Journal of Agricultural Research*. 2: 21-24.

357 Olawuyi O.J., Bello O.B., Ntube C.V. and Akanmu A.O. (2015) Progress from selection of some
358 maize cultivars' response to drought in the derived savanna of Nigeria. *Agrivita*, 37 (1) 8-17

359 Olawuyi O.J., Odebode A.C., Alfar A., Olakojo S.A. and Adesoye A.I. 2010. Performance of maize
360 genotypes and arbuscularmycorrhizal fungi in Samara District of Southwest Region of Doha-Qatar.
361 *Nigerian Journal of Mycology*, 3(1): 86-100.

362 Olowe, O.M., Odebode, A.C., Olawuyi, O.J. and Sobowale, A.A. (2017) Molecular Variability of
363 *Fusarium verticillioides*(Sacc.) in Maize from Three Agro-Ecological Zones of Southwest Nigeria.
364 *American Journal of Molecular Biology*, 7, 30-40

365 OnoE.Y.S., Fungaro M.H.P., Sofia S.H., Miguel T.D., Sugiura Y., Hirooka E.Y. (2010). *Fusarium*
366 *verticillioides* strains isolated from corn feed: characterization by fumonisin production and RAPD
367 fingerprinting. *Braz. arch. biol. technol.* vol.53 no.4 Curitiba

368

369 Oren L., Ezrati S., Cohen D., and Sharon A. (2003) Early Events in the *Fusarium verticillioides*-
370 Maize Interaction Characterized by Using a Green Fluorescent Protein-Expressing Transgenic
371 Isolate. *Appl Environ Microbiol.* 2003 Mar; 69(3): 1695–1701.

372

373 Orsi R.B., Correa B., Possi C.R., Schammas E.A., Nogueira J.R., Dias S., Malozzi M.A.B. 2000.
374 Mycoflora and occurrence of fumonisin of freshly harvested and stored hybrid maize. *Journal of*
375 *storage and Product Research*. 36-87.

376

377 Perveen K., Haseeb A. and Shukla P.K. (2010) Effect of *Sclerotinia sclerotiorum* on the disease
378 development, growth, oil yield and biochemical changes in plants of *Mentha arvensis*. *Saudi Journal*
379 *of Biological Sciences*, 17, Issue 4, Pages 291-294

380

381 Reddy M.N. and Sireesha C.H. 2013. Role of oxidative enzymes and biochemical constituents in
382 imparting resistance to groundnut (*Arachis hypogea* L.) against stem rot of diseases caused by
383 *Sclerotium rolfsii*. *Bioresearch Bulletin*. 36-41pp.

384 Samapundo S., De Meulenaer B., Osei-Nimoh D., Lamboni Y., Debevere J., Devlieghere F. (2007).
385 Can phenolic compounds be used for the protection of corn from fungal invasion and mycotoxin
386 contamination during storage? *Food Microbiology* 24 (2007) 465–473.

387

388 Sharma, T.R. (2003) Molecular Diagnosis and Application of DNA Markers in the Management of
389 Fungal and Bacterial Plant Diseases. *Indian Journal of Biotechnology*, 2, 99-109

390

391 Singh N., Ambika R., Meena S. and Girish M. 2012. Biochemical response and host-pathogen
392 relation of stalk rot fungi in early stages of maize (*Zea mays* L.). *African Journal of Biotechnology*.
393 11 (82): 14837-14843.

394 Singh P., Shukla R., Prakash B., Kumar A., Singh S. and Mishra P.K. 2010. Chemical profile,
395 antifungal, antiaflatoxic and antioxidant activity of *Citrus maxima* Burm. and *Citrus sinensis* (L.)
396 Osbeck essential oils and their cyclic monoterpene, DL-limonene. *Food and chemical toxicology*. 48
397 (6), 1734-1740.

398 Sobowale A.A. (2011). Determination of infective, non-lethal dosage of *Fusarium verticillioides* in
399 maize (*Zea mays*) stem and effective inoculation method in the screenhouse. *Journal of Agriculture*
400 *and Biological Sciences* Vol. 2(5) pp.118-122

401

- 402 Sobowale A.A., Cardwell K.F., Odebode A.C., Bandyopadhyay R., Jonathan S.G. (2007).
403 Persistence of *Trichoderma* species within maize stem against *Fusarium verticillioides*. *Arch.*
404 *Phytopathol. Plant Prot.* 40, 3: 215-231.
- 405
406 Sobowale A.A., Cardwell K.F., Odebode A.C., Bandyopadhyay R., Jonathan S.G. (2005). Growth
407 inhibition of *Fusarium verticillioides* (Sacc.) Nirenberg by isolates of *Trichoderma Pseudokoningii*
408 strains from maize plant parts and its rhizosphere. *Journal of Plant Protection Research*, 45 (4) Pages
409 249-265.
- 410 Stuper-Szablewska K, Kurasiak-Popowska D, Nawracała J, Perkowski J. (2017). Response of non-
411 enzymatic antioxidative mechanisms to stress caused by infection with *Fusarium* fungi and chemical
412 protection in different wheat genotypes. *Chem Ecol.* 33:949–962.
- 413 Thompson M. and Raizada M. (2018). Fungal Pathogens of Maize Gaining Free Passage Along the
414 Silk Road. *Pathogens*, 7, Number 4, 81
- 415