

FLUORESCENCE AND GROWTH OF EGGPLANT UNDER IRRIGATION LEVELS AND SILICON DOSES

ABSTRACT

Aims: The study was conducted to evaluate the fluorescence and growth of eggplant under influence of water deficit and silicon doses.

Study Design: The design was a randomized complete block design, in a 5 x 2 factorial arrangement, with four replications and one plant per plot, totaling 40 experimental units.

Length and place of study: The research was implemented between September and November 2016, in a greenhouse of the Center of Science and Technology Agrifood, at the Federal University of Campina Grande (UFCG / CCTA), Campus of Pombal-PB, Brazil.

Methods: Doses of 0, 75, 150, 225 and 300 mg L⁻¹ of silicon and the irrigation slides of 50 and 100% of real evapotranspiration - ETr were used, which were obtained by the weighing the pots daily in order to keep the weight close to the field capacity.

Results: The use of 50% level of the ETr provides reductions of 5.58; 7.92 and 6.84% of fluorescence in the initial, maximum and quantum yield of the photosystem. The fresh and dry mass of the stem presented the maximum productivity (140.8 and 48.1 g) in the plants irrigated with 100% Etr and submitted to the doses of 106 and 110 mg L⁻¹ of Si.

Conclusion: The eggplants plants fluorescence and biomass is reduced by the decrease of the irrigation level; The doses of silicon did not provide development in the fluorescence and dry mass of the leaf; The use of 100% level of the ETr, combined with the application of 108 mg L⁻¹ of silicon showed the best results in the stem biomass accumulation.

Keywords: *Solanum melongena*, *silicic fertilization*, *water deficit*, *abiotic stress*.

1. INTRODUCTION

The eggplant (*Solanum melongena* L.) is a crop of high socioeconomic importance, producing worldwide approximately 52.3 million tons in an area of 1.85 million ha. China (32.9 million tons), India (12.5 million tons) and Egypt (1.3 million tons) are the largest producers [1]. In Brazil, recent data estimate that the production of this vegetable is around 90 thousand tons [2], being mainly cultivated by small and medium producers.

The cultivation of this oilseed crop has been gaining notoriety, mainly due to its nutritional and phytotherapeutic properties, being an important source of phenolic compounds, carotenoids and alkaloids [3]. Cultivation at temperatures between 20 and 35 °C were observed in great conditions, especially by being a tropical climate crop and presenting itself as a good alter-native to the semi-arid.

One of the main limitations imposed on agriculture in the semi-arid region is related to low water availability due to low rainfall and high annual evapotranspiration [4]. In the eggplant crop, water deficits can negatively affect characteristics such as mineral composition, leaf

29 water potential, photosynthesis and fruit yield [5,6]. Thus, strategies that minimize the
30 deleterious effects and/or increase the efficiency of water use is being studied [7,8].

31 Silicon (Si) is the second most abundant element of the earth. It is absorbed by plants as
32 monostetic acid (H_4SiO_4), being reported in the literature as the beneficial element for some
33 crops and essential for other species such as rice and sugar cane. Some of the benefits
34 provided by Si include increased tolerance to biotic and abiotic stresses, thus Si deposition
35 on the cell wall may decrease cuticular transpiration, thereby reducing plant water loss [9].
36 Benefits of Si application on the depletion of water stress was reported in potato [10], pepper
37 [11], arugula [12] and tomato [13].

38 Therefore, the objective of this study was to evaluate the fluorescence and growth of the
39 eggplant under influence of water deficit and silicon doses.

41 2. MATERIAL AND METHODS

42
43 The research was carried out between September and November 2016, in a greenhouse of
44 the Center of Science and Technology Agrifood, at the Federal University of Campina
45 Grande (UFCG / CCTA), Campus of Pombal-PB, Brazil, at geographical coordinates 6 ° 46
46 '16' of Latitude S and 37 ° 49' 15 " longitude W, at an altitude of 144 m.

47 The experimental design was a randomized block design in a 5 x 2 factorial arrangement for
48 five silicon doses (0, 75, 150, 225 and 300 mg L⁻¹) and two irrigation levels (50 and 100%
49 of the actual evapotranspiration - ETr), with four replicates and one plant per plot, totaling 40
50 experimental units.

51 In order to determine the real evapotranspiration, we used the weighing lysimeter method in
52 the treatments that received 100% of the ETr. The weight of the pots was determined at field
53 capacity (FCw) from saturation by capillarity followed by drainage until constant weight,
54 considering the FCw and weighed daily the present weight (Pw) of each pot, with these data
55 used the equation 1, in which the ETr was determined with the division of the subtraction of
56 these numbers by the area of the pot.

$$57 \quad ETr = \frac{FCw - Pw}{A} \quad \text{mm} \quad \text{Eq. 1}$$

58 As an experimental unit, a 12.8 L pot was filled with the sample of a Fluvic Neosol collected
59 in the 0-40 cm depth [14]. It was sieved and analyzed to obtain its physical and chemical
60 characteristics, following the methodologies described by Embrapa [15], in the Laboratory of
61 Soils and Plant Nutrition of CCTA / UFCG, as indicated in Table 1.

62 **Table 1. Chemical and physical attributes of the soil used in the experiment**

Chemical characteristics								
pH	EC _{se}	P	K ⁺	Na ⁺	Ca ²⁺	Mg ²⁺	Al ³⁺	H + Al ³⁺
CaCl ₂ 1:2,5	dS m ⁻¹	mg dm ⁻³	cmol _c dm ⁻³					
6.50	0.91	7.00	0.52	0.36	4.55	2.35	0.00	0.00
Physical characteristics								
Sand	Silt	Clay	AD	DP	Total porosity		Textural	
g kg ⁻¹						%	class	
715	213	72	1.48	2.86	48		Sandy loam	

63 *pH – hydrogen potential, Ca^{2+} and Mg^{2+} extracted with 1 M L^{-1} KCl at pH 7.0; Na^+ and K^+ extracted*
64 *using 1 M L^{-1} NH_4OAc at pH 7.0; $\text{Al}^{3+} + \text{H}^+$ extracted using 0.5 M L^{-1} CaOAc pH 7.0; ECse – electrical*
65 *conductivity of the saturation extract; AD - apparent density; DP –particle density.*

66 The eggplant (*Solanum melongena* L.) seedlings "Embu" cultivar were cultivated in 128-cell
67 expanded polystyrene trays, using as substrate the Tropstrato® commercial mix with two
68 seeds per cell with subsequent thinning, leaving only one seedlings per cell. Transplanting
69 was carried out at 40 days after sowing (DAS), when the plants had two true leaves and a
70 height of approximately 15 cm.

71 Fertilization with macronutrients (except N) and micronutrients were performed accord-
72 ing to the Malavolta [16] recommendation for potting. The following doses were applied, in mg dm^{-3} :
73 P = 100; K = 160; Ca = 230; Mg = 20; S = 155; B = 0.5; Cu = 1.5; Fe = 10; Mn = 4; Mo =
74 0.15 and Zn = 5.0 and the subsequent sources: simple superphosphate, KCl, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$
75 H_3BO_3 , $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, Fe-EDTA, $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$, ammonium molybdate and $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$
76 respectively. In potassium fertilization, the amounts of K supplied by potassium silicate were
77 discounted to balance the nutrient doses between the treatments.

78 Silicon leaf fertilization was supplied by six sprays of the product Quimifol Silicio® (10% Si +
79 8.3% K, density = 1.31 kg L^{-1}). Applications started seven days after transplanting (DAT) and
80 the other applications were provided biweekly. The amount was applied respecting the
81 vegetative development of the crop and increasing gradually in each application.

82 The solutions were prepared in one liter containers, each one representing a dose of silicon.
83 As the product used to provide adequate amounts of the studied element (Quimifol) had
84 potassium in its composition, it was necessary to use the potassium nitrate (KNO_3) to
85 balance the nutrient concentrations at lower doses of Si. Urea was used to provide the
86 concentrations of nitrogen applied along with KNO_3 in silicon treatment. In order to avoid
87 possible problems, its application was separated in two days: on the first day, the doses for
88 KNO_3 and the second on Quimifol + Urea.

89 At 73 DAS, using a portable modulated fluorometer, the emission of chlorophyll a
90 fluorescence was quantified and it was possible to determine the initial fluorescence (F_0),
91 the maxi-mum fluorescence (F_m), the variable fluorescence (F_v) and the maximum quantum
92 yield of pho-tosystem II (F_v / F_m) in each plant. At 114 DAS, through a destructive
93 evaluation of the experi-ment, the biomass of the eggplants were determined. These were
94 collected, fractionated and weighed for the determination of leaf fresh mass (LFM), stem
95 fresh mass (SFM) and shoot fresh mass (SHFM). Then the material was packed in paper
96 bags and placed in an air circulation oven at 65°C for 72 hours for the determination of leaf
97 dry mass (LDM), stem dry mass (SDM) and shoot dry mass (SHDM).

98 The data were submitted to analysis of variance by the F test ($p < 0.05$). The mean values of
99 the irrigation levels were compared by F test ($p < 0.05$), which is conclusive for two factors
100 from the same source of variation. Mean values for the silicon doses were analyzed by
101 polynomial regression at 5% probability. Statistical software Sisvar version 5.6 was used for
102 data analysis [17].

103 3. RESULTS AND DISCUSSION

104

105 According to the summary of the analysis of variance, there was a significant effect for
106 interaction between irrigation levels and silicon doses for fresh leaf and stem masses, and
107 for stem dry mass (Table 1). The irrigation levels provided significance for the initial and
108 maximum fluorescence, quantum yield of photosystem II and the fresh and dry masses of

the aerial part. The silicon doses promoted an isolated effect on the initial, variable and maximum fluorescence, quantum yield of photosystem II and on the dry and fresh masses of the shoot and dry leaves.


Table 2. Analysis of variance, by the mean square values for the initial fluorescence (F0), maximum fluorescence (Fm), variable fluorescence (Fv), quantum yield of

Source of variation	DF	Mean squares									
		F0	Fm	Fv	Fv/Fm	SFM	SDM	LFM	LDM	SHFM	SHDM
Blocks	3	64 ^{ns}	28140 ^{**}	26026 ^{**}	0.0006 ^{ns}	783 [*]	117 [*]	80 ^{ns}	3.6 ^{ns}	798 ^{ns}	228 ^{**}
Irrigation levels (IL)	1	725 [*]	9517 ^{ns}	15171 [*]	0.0135 ^{**}	12179 ^{**}	3417 ^{**}	40 ^{ns}	1.1 ^{ns}	3023 ^{**}	2799 ^{**}
Doses silicon (DS)	4	252 [*]	10674 [*]	10758 [*]	0.0016 [*]	1108 ^{**}	161 ^{**}	158 ^{ns}	9.7 [*]	841 [*]	168 [*]
Interaction (IL* DS)	4	34 ^{ns}	2213 ^{ns}	1530 ^{ns}	0.0005 ^{ns}	641 [*]	91.8 [*]	194 [*]	3.6 ^{ns}	141 ^{ns}	57.0 ^{ns}
Residue	40	100	4065	3336	0.0005	239	25.8	70	3.6	342	43.8
Averages		156.8	668.5	511.2	0.760	105.5	33.7	44.18	9.9	141.2	42.6
CV (%)		6.46	9.54	11.30	3.00	14.65	15.09	18.93	19.31	13.29	15.51

photosystem II (Fv / Fm), stem fresh mass (SFM) and stem dry mass (SDM), leaf fresh mass (LFM) and leaf dry mass (LDM), shoot fresh mass (SHFM) and shoot dry mass (SHDM) of the eggplant submitted to silicate fertilization and irrigation levels.

** $P < 0.01$; * $P < 0.05$; ^{ns} $P > 0.05$

The 100% irrigation level provided higher efficiency of the photosynthetic apparatus, represented by the initial and variable fluorescence, presenting the largest increases of 161.05 and 530.7 quantum⁻¹ electrons, respectively (Figure 1A and 1B). The results obtained in the 50% Etr level shows a reduction of 5.58 and 7.92% regarding the level of 100%. Then, the greater availability of water to the plant provided, the greater absorption and translocation capacity of nutrients in the vegetal tissues. Thus, the higher availability of water results in lower transpiration losses, resulting in a greater nutritional contribution to the plant and improving the cooling of plant tissues through energy dissipation [18].



126 **Figure 1. Initial fluorescence - F0 (A), variable fluorescence - Fv (B) and quantum yield**
 127 **of photo-system II - Fv/Fm (C) submitted to different irrigation levels.**

128 The quantum yield of photosynthetic II showed similarity to F0 and Fv, where the
 129 highest efficiency was obtained in the 100% Etr (0.78) level, presenting a superiority of
 130 6.84% (Figure 1C). This result shows that the plants submitted to the 50% level of Etr
 131 promoted stress to the photosynthetic apparatus, since values below 0.75 quantum⁻¹
 132 electrons are considered stress conditions. The photosynthetic apparatus is intact when
 133 values vary between 0.75 and 0.85 quantum⁻¹ electrons [19,20].

134 Results that validate those obtained by Magalhaes et al. [21], values of 0.78 for
 135 quan-tum efficiency of FSII in the level of 125% Etr found in common bean (*Phaseolus*
 136 *vulgaris* L.). Neves et al. [22] observed in sunflower (*Helianthus annuus*) values within the
 137 tolerable limits (0.75-0.85) of plants grown under ideal conditions of water regime.

138 The effect of the silicon doses on the initial fluorescence (F0) presented results that best fit
 139 the increasing linear effect of 164.81 quantum⁻¹ electrons at the dose of 300 mg L⁻¹ of Si
 140 (Figure 2A). This increase in F0 rates can be considered destructive to the photosynthetic
 141 apparatus, since the uptake efficiency is reduced as F0 is raised, providing the FSII
 142 inactivation or the inhibition of excitation transfer between the antenna complex and the
 143 center of reaction [23].




Figure 2. Initial fluorescence - F0 (A), variable fluorescence - Fv (B), maximum fluorescence (C) and quantum yield of photosystem II - Fv/Fm (D) submitted to different irrigation levels

**** $P < 0.01$; * $P < 0.05$**

It was observed for the variable and maximum fluorescence that the results fit the quadratic model, with the highest values in the plants submitted to the control treatment with 572.76 and 715.6 quantum⁻¹ electrons, resulting in decreases of 9.6 and 4.9% in relation to the highest dose tested, respectively (Figure 2B and 2C). The effect promoted by the Si on the fluorescence indices in this study differs from those obtained by Ferraz et al. [24] in cotton (*Gossypium hirsutum* L.), where they verified that the Si promoted increases in the maximum and variable fluorescence in the cultivars BRS Rubi and BRS Topázio. Maghsoud et al. [25] found that the application of 6 mM Si reduced the maximum fluorescence in wheat plants (*Triticum aestivum* L.).

The quantum yield of photosystem II behaved in a decreasing linear manner with the increasing doses of silicon at the maximum increment (0.795 quantum⁻¹ electrons) in the control treatment, reaching a reduction of 7.5% at the maximum dose (300 mg L⁻¹) of Si tested (Figure 2D). Thus, the application of Si promoted disturbances in the activity of FSII, reducing the photochemical efficiency and reducing the use and conversion of light energy. This effect is related to the increase of F0, promoting damage to the photosynthetic apparatus due to the decrease of FSII efficiency through the inactivation of reaction centers [26].

Al-Aghabary et al. [27] observed that the application of 2.5 mM Si increased the quantum yield of FSII in tomato plants (*Lycopersicon esculenta* L.) under conditions of saline stress. Maghsoud et al. [25] verified that the application of 6 mM Si in wheat plants promoted an increase in the quantum efficiency of FSII.

It was verified for the fresh and dry masses of the stem the effect of the interaction between the irrigation levels and Si doses, with the maximum increments (140.8 and 48.1 g) in the plants irrigated with 100% ETr and submitted to the doses of 106 and 110 mg L⁻¹ of Si, respectively (Figures 3A and 3B). The plants irrigated with 50% ETr presented average values of 88.1 and 24.4 g plant⁻¹ for the fresh and dry mass of the stem, reaching reductions of 59.8 and 97.1%. These results indicate that the application of Si promotes the development in plants under water deficit in function of promoting improvements in nutritional balance, providing a greater accumulation of Si in the cell wall and favoring the accumulation of biomass [28,12].




Figure 3. Fresh (A) and dry matter of the stem (B), leaf fresh mass (C) and leaf dry mass (D) of eggplant submitted to irrigation levels and silicate fertilization.

** $P < 0.01$; * $P < 0.05$

The accumulation of fresh leaf mass was superior in the 100% level ETr with 51.44 g plant⁻¹ at dose 0 of Si (Figure 3C). While the application of 135 mg L⁻¹ of Si promoted the increase (49.6 g plant⁻¹) in leaf mass content in plants irrigated with 50% ETr. This effect suggests that decreasing water availability reduces leaf emission and mass production. Thus, the Si applied to plants with low water availability promotes improvements due to the deposition of Si in the roots, leaves and stem, reducing water loss through transpiration [12].

The dry matter of the leaves had an effect only for the Si doses. The silicon application linearly reduced the accumulation of dry matter, promoting losses of 21.2% when comparing the values of the lowest and highest dose applied (Figure 3D). The absorption and translocation did not occur efficiently due to the low root and xylem activity of the transporter, since the increase of the applied dose does not guarantee that it is absorbed by the plant [28].

The fresh shoot mass behaved similar to the dry mass of the leaves, reducing the accumulation as a function of Si doses with losses of 15.6% when comparing the values of the lowest and highest dose applied (Figure 4A). This response may be due to the cellular wall stiffness as a result of Si accumulation in the tissues, resulting in reduced leaf water potential [29] causing a low translocation capacity of photo-assimilates in the plant.




Figure 4. Shoot mass (A) in function of doses of silicon and irrigation levels (B).

**** $P < 0.01$; * $P < 0.05$**

The effect of the 100% ETr irrigation level promoted the highest increases in fresh and dry shoot mass, increasing 13.1 and 48.7% compared to the values obtained in plants irrigated with 50% ETr (Figure 4B). The water supply under reduced conditions may promote limitations to the stomatal activity and to the photosynthetic apparatus, resulting in the decline of the partitioning and accumulation of biomass by the plant [30].

The benefit of silicon fertilization on the accumulation of biomass in plants is still controversial, as in *Corymbia citriodora*, where the fertilization with Si did not have an effect on the fresh and dry masses of the roots and shoot [31]. The application of Si via irrigation water in melon (*Cucumis melo* L.) did not have an effect on the accumulation of dry shoot matter and in total plant dry mass [28]. The leaf fertilization of 150 mg L⁻¹ of Si in okra (*Abelmoschus esculentus* L.) plants under salt stress promoted increases of 40% and 36% in fresh and dry shoot mass and 32% and 25% in fresh and dry root mass, respectively [32].

4. CONCLUSION

The fluorescence and biomass of the eggplant plants is reduced by the decrease of the irrigation level;

The silicon doses did not provide increases in the fluorescence and dry mass of the leaf;

The 100% irrigation level of the ETr associated with the application of 108 mg L⁻¹ of silicon caused the best results in the stem mass accumulation.

6. REFERENCES

1. FAO. FAOSTAT: database. Accessed: 29 April 2019. Available: <http://www.fao.org/faostat/en/#data/QC/visualize>.
2. Marouelli WA, Braga MB, Silva HR, Ribeiro CSC. Irrigação na cultura da berinjela. Embrapa Hortaliças: Circular Técnica. 2014(135). Portuguese.
3. Gürbüz N, Uluişik S, Frary A, Frary A, Doğanlar S. Health benefits and bioactive compounds of eggplant. Food chemistry. 2018; 268(1):602-610. <https://doi.org/10.1016/j.foodchem.2018.06.093>

- 228 4. Tavares VC, Arruda ÍRP, Silva DG. Desertificação, mudanças climáticas e secas no
229 semiárido brasileiro: uma revisão bibliográfica. Geosul. 2019;34(70): 385-405.
230 Portuguese. <https://doi.org/10.5007/2177-5230.2019v34n70p385>
- 231 5. Díaz-Pérez JC, Eaton TE. Eggplant (*Solanum melongena* L.) plant growth and fruit
232 yield as affected by drip irrigation rate. HortScience. 2015;50(11):1709-1714.
233 <https://doi.org/10.21273/HORTSCI.50.11.1709>
- 234 6. Mariano-Nasser FAC, Borges CV, Ramos JA, Nasser MD, Lundgren GA, Furlaneto
235 KA, Vieites RL. Bioactive compounds and enzymatic activity in minimally processed
236 eggplant packed under active modified atmosphere. Semina: Ciências Agrárias.
237 2019; 40(1):139-148. <http://dx.doi.org/10.5433/1679-0359.2019v40n1p139>
- 238 7. Mohawesh O. Utilizing deficit irrigation to enhance growth performance and water-
239 use efficiency of eggplant in arid environments. Journal of Agricultural Science and
240 Technology. 2016;18(1): 265–276.
- 241 8. Çolak YB, Yazar A, Sesveren S, Colak I. Evaluation of yield and leaf water potential
242 (LWP) for eggplant under varying irrigation regimes using surface and subsurface
243 drip systems. Scientia horticulturae. 2017;219(1):10-21.
244 <https://doi.org/10.1016/j.scienta.2017.02.051>
- 245 9. Pontigo S, Ribera A, Gianfreda L, Mora ML, Nikolic M, Cartes P. Silicon in vascular
246 plants: uptake, transport and its influence on mineral stress under acidic conditions.
247 Planta. 2015;242(1): 23-37. <https://doi.org/10.1007/s00425-015-2333-1>
- 248 10. Pilon C, Soratto RP, Broetto F, Fernandes AM. Foliar or soil applications of silicon
249 alleviate water-deficit stress of potato plants. Agronomy Journal. 2014;106(6): 2325-
250 2334. <https://doi.org/10.2134/agronj14.0176>
- 251 11. Pereira TS, Lobato AKS, Silva MHL, Lobato EMSG, Costa DV, Uchoa EB, et al.
252 Differential responses produced by silicon (Si) on photosynthetic pigments in two
253 pepper cultivars exposed to water deficiency. Australian Journal of Crop
254 Science. 2015;9(12): 1265-1270.
- 255 12. Jesus EGD, Fatima RT, Guerrero AC, Araújo JLD, Brito ME. Growth and gas
256 exchanges of arugula plants under silicon fertilization and water restriction. Revista
257 Brasileira de Engenharia Agrícola e Ambiental. 2018; 22(2): 119-124.
258 <http://dx.doi.org/10.1590/1807-1929/agriambi.v22n2p119-124>
- 259 13. Silva ON, Lobato AKS, Ávila FW, Costa RCL, Neto CO, Santos Filho BG, et al.
260 Silicon-induced increase in chlorophyll is modulated by the leaf water potential in
261 two water-deficient tomato cultivars. Plant, Soil and Environment. 2012;58(11):481-
262 486.
- 263 14. EMBRAPA. Sistema brasileiro de classificação de solos. 5 ed. Brasília: Embrapa.
264 2018; 353.
- 265 15. EMBRAPA. Manual de métodos de análise de solo. Manual de análises químicas de
266 solos, plantas e fertilizantes. 3 ed. Brasília: Embrapa. 2017; 627.
- 267 16. Malavolta E, Vitti GC, Oliveira SA. Avaliação do estado nutricional das plantas:
268 princípios e aplicações. Piracicaba: Potafos. 1997; 319.

- 269 17. Ferreira DF. Sisvar: A Guide for its Bootstrap procedures in multiple comparisons.
270 Ciência e Agrotecnologia. 2014; 38(2):109-112. Available:
271 <http://dx.doi.org/10.1590/S1413-70542014000200001>
- 272 18. Taiz L, Zeiger E, Møller IM, Murphy A. Fisiologia e desenvolvimento vegetal. 6.ed.
273 Porto Alegre: Artmed. 2017; 888.
- 274 19. Suassuna JF, Melo AS, Sousa MSS, Costa FS, Fernandes PD, Pereira VM, et al.
275 Desenvolvimento e eficiência fotoquímica em mudas de híbrido de maracujazeiro
276 sob lâminas de água. Bioscience Journal. 2010; 26(4):566-571.
- 277 20. Silva FG, Dutra WF, Dutra AF, Oliveira IM, Filgueiras LMB, Melo AS. Trocas
278 gasosas e fluorescência da clorofila em plantas de berinjela sob lâminas de
279 irrigação. Revista Brasileira de Engenharia Agrícola e Ambiental. 2015;19(10):946-
280 952. <http://dx.doi.org/10.1590/1807-1929/agriambi.v19n10p946-952>
- 281 21. Magalhaes ID, Lyra GB, Souza JL, Teodora I, Cavalcante CA, Ferreira RA, et al.
282 Physiology and grain yield of common beans under evapotranspired water
283 reposition levels. Irrigat Drainage Systems Engineering. 2017;6(1):183-191.
284 <http://dx.doi.org/10.4172/2168-9768.1000183>
- 285 22. Neves JMG, Aquino LA, Berger PG, Neves JCL, Rocha GC, Barbosa EA. Silicon
286 and boron mitigate the effects of water deficit on sunflower. Revista Brasileira de
287 Engenharia Agrícola e Ambiental. 2019;23(3):175-182.
288 <http://dx.doi.org/10.1590/1807-1929/agriambi.v23n3p175-182>
- 289 23. Souza CAS, Pinto DG, Aguilar MAG, Coelho RL, Gasparini Júnior AJ, Cao JR, et al.
290 Influência do silício sobre o crescimento, a fluorescência da clorofila a e na
291 incidência de insetos-pragas em genótipos de cacau. Agrotrópica. 2012;24(1):31-
292 40. Portuguese.
- 293 24. Ferraz RLS, Beltrão NEM, Melo AS, Magalhães ID, Fernandes PD, Rocha MS.
294 Trocas gasosas e eficiência Fotoquímica de cultivares de algodoeiro herbáceo sob
295 aplicação de silício foliar. Semina: Ciências Agrárias. 2014;35(2):735-748.
296 Portuguese. <http://dx.doi.org/10.5433/1679-0359.2014v35n2p735>
- 297 25. Maghsoudi K, Emam Y, Ashraf M. Influence of foliar application of silicon on
298 chlorophyll fluorescence, photosynthetic pigments, and growth in water-stressed
299 wheat cultivars differing in drought tolerance. Turkish Journal of Botany.
300 2015;39(1):1-10. <http://dx.doi.org/10.3906/bot-1407-11>
- 301 26. Oliveira WJ, Souza ER, Santos HRB, Silva EFF, Duarte HHF, Melo VM.
302 Fluorescência da clorofila como indicador de estresse salino em feijão caupi.
303 Revista Brasileira de Agricultura Irrigada. 2018;12(3):2592-2603. Portuguese.
304 <http://dx.doi.org/10.7127/RBAI.V12N300700>
- 305 27. Al-Aghabary K, Zhu Z, Shi Q. Influence of silicon supply on chlorophyll content,
306 chlorophyll fluorescence, and antioxidative enzyme activities in tomato plants under
307 salt stress. Journal of Plant Nutrition. 2005;27(12):2101-2115.
308 <https://doi.org/10.1081/PLN-200034641>
- 309 28. Gomes FAL, Araújo RHCR, Nóbrega JC, Fátima RT, Silva MS, Santos AS, et al.
310 Application of silicon to alleviate irrigation water salinity in melon growth. Journal of

311 Experimental Agriculture International. 2018;25(6):1-9.
312 <https://doi.org/10.9734/JEAI/2018/43767>

313 29. Zanetti LV, Milanez CRD, Gama VN, Aguilar MAG, Souza CAS, Campostrini E, et
314 al. Leaf application of silicon in young cacao plants subjected to water deficit.
315 Pesquisa Agropecuária Brasileira. 2016;51(3):215-223.
316 <http://dx.doi.org/10.1590/S0100-204X2016000300003>

317 30. Anjos RAR, Santos LCS, Oliveira DB, Amaro CL, Rios JM, Rocha GT. Initial growth
318 of *Jatropha curcas* plants subjected to drought stress and silicon (Si) fertilization.
319 Australian Journal of Crop Science. 2017;11(4):478-484.
320 <http://dx.doi.org/10.21475/ajcs.17.11.04.377>

321 31. Castro EB, Santos LTD, Fernandes LA, Trajano CV. Silicato de alumínio em
322 substrato para produção de mudas de *Corymbia citriodora*. Floresta e Ambiente.
323 2016;23(2):229-236. <http://dx.doi.org/10.1590/2179-8087.106814>

324 32. Abbas T, Balal RM, Shahid MA, Pervez MA, Ayyub CM, Aqueel MA, et al. Silicon-
325 induced alleviation of NaCl toxicity in okra (*Abelmoschus esculentus*) is associated
326 with enhanced photosynthesis, osmoprotectants and antioxidant metabolism. Acta
327 Physiologiae Plantarum. 2015;37(6). <https://doi.org/10.1007/s11738-014-1768-5>

328