

**BIOCONTROL POTENTIAL OF *Bacillus thuringiensis* ISOLATED FROM SOIL  
AGAINST MOSQUITO LARVA**

ABSTRACT

A major challenge for achieving successful mosquito control is overcoming insecticide resistance. This study was carried out to assess the larvicidal activity of *Bacillus thuringiensis* isolated from different soil samples within Sokoto metropolis using standard methods. Confirmatory identification of the organism was made based on biochemical characterization and microscopic observation. The larvicidal activity of *Bacillus thuringiensis* isolated was tested against the larva of mosquito using three dilutions of the Bacillus culture in a bioassay. The isolated organisms were confirmed as *Bacillus thuringiensis*. The result of the bioassay showed variation in the level of efficacy of the bacteria and depended on time of the exposure. Mortality rate greater than 20% was observed after 60 minutes and increased to 100% after time of exposure was increased for all dilutions of *B. thuringiensis* used. The results showed that *Bacillus thuringiensis* toxins can be bacteriocidal to mosquito larvae in a matter of minutes depending on the concentration ingested by the larvae. This in essence proved that *Bacillus thuringiensis* is an effective biolarvicide that can be used to reduce and possibly eradicate the nuisance of disease-causing mosquitoes and aid in the rollback of malaria.

Keywords: Mosquito control, *Bacillus thuringiensis*, bioassay, larvicidal activity

24  
25

## INTRODUCTION

26 Mosquitoes are associated with the transmission of pathogens to humans and other vertebrates.  
27 Some of these include the causative agents of malaria, filariasis and dengue, as well as other  
28 mosquito-borne zoonotic arbovirus, like Saint Louis Encephalitis Virus (SLEV), West Nile  
29 Virus (WNV) and Eastern Equine Encephalitis Virus (EEEV) [1]. Significant morbidity and  
30 mortality are recorded as results of these diseases are as a result of the inherent difficulty of  
31 controlling mosquitoes [2, 3]. Increase in the distribution of mosquitoes was associated with the  
32 emergence of viruses and diseases in new areas [4]. WHO [3] reported malaria to be the world  
33 most important vector-borne disease. Cases of these diseases have been reported in more than  
34 100 countries, with approximately more than 3 billion people living in endemic areas. More than  
35 200 million cases of malaria and eight hundred thousand (800,000) malaria-related deaths are  
36 recorded every year [5]. The increase in number of malaria cases is as a result of deteriorating  
37 health systems, increase resistance of anopheline to insecticides, time taken to develop an  
38 effective vaccine and as well resistance of plasmodium to antimalarial drugs [1].

39 Dengue, including dengue hemorrhagic fever and Dengue Shock Syndrome (DSS) transmitted  
40 by *Aedes* mosquitoes is rapidly becoming a worldwide disease, threatening a third of the world  
41 population, with an estimate of 50-100 million cases every year [6, 7]. So also, *Lymphatic*  
42 *filariasis* caused by *Wuchereria bancrofti*, which is transmitted by mosquitoes, affects more than  
43 120 million people around the world [8]. Lack of effective vaccine against these diseases has left  
44 the control of mosquito population as the most effective way to prevent vector-borne diseases [9,  
45 10].

46 Chemical insecticides have been used in the last century to successfully control mosquitoes of  
47 the genus *Aedes* and *Anophelles*. Current ecological and environmental protection standards halt

48 the use of these chemicals, because of their adverse effects on non-target species, including  
49 humans, environmental impact, contamination soil and water and development of mosquito  
50 resistance to insecticides [11]. New strategies were created to replace the use of chemical  
51 insecticides. They include Integrated Pest management (IPM) that has guidelines. Guidelines of  
52 which are based on environmental planning, public awareness and biological control that control  
53 the mosquitoes more efficiently while preserving the environment from contamination [1].  
54 Commercial preparations of *Bacillus thuringiensis* (*Bt*) as a biocontrol agent has been the  
55 greatest success in microbial pesticides, with more than 95% of the microbial pesticides sold  
56 being of this bacterial agent [12]. *Bacillus thuringiensis* (*Bt*) is a gram-positive, rod-shaped and  
57 spore forming bacteria that is mostly found in the soil and produces polymorphic crystal proteins  
58 [13]. The insecticidal activity of *Bt* is due to the proteic parasporal inclusions that are produced  
59 during sporulation [14]. The insecticidal proteinaceous crystals (ICPs) comprised one or more  
60 crystal (Cry) and Cytolytic (Cyt) proteins recognized as  $\delta$ -endotoxin. When ingested by the  
61 target insect, the ICPs dissolve in the midgut of the larva releasing prototoxins that eventually  
62 lead to the formation of pores that causes cell-cytolysis [15, 16].  
63 Despite the use of *Bacillus thuringiensis* as a biocontrol agent for over 30 years, no significant  
64 resistance was recorded. However, the search for natural *Bt* isolates with increase activity against  
65 mosquito and other insect is still encouraged. Recently, *Bt* with increased activity against *Aedes*  
66 *caspius* and *Culex pipiens* were isolated [17]. In this study, the larvicidal activity of *Bt* isolated  
67 from soil samples will be evaluated.

68

69

70

## 71 MATERIALS AND METHODS

### 72 Sample Collection

73 A total of 4 soil samples were collected from Tamaje, Maberu, Arkilla and Kantin Daji areas of  
74 Sokoto state, where there is no previous record of application of *Bacillus thuringiensis* based  
75 insecticides. The soil samples were collected aseptically from top of 5cm depth. The samples  
76 were placed immediately in plastic bags and labeled appropriately [18]. The soil samples were  
77 transported to the laboratory and stored at room temperature. In the same vein, mosquito larva  
78 was collected from stagnant waters around Sokoto metropolis.

### 79 Media Preparation

80 The media to be used to culture the bacteria such as Nutrient agar and Luria-Bertani will be  
81 prepared according to the manufacturer's instruction.

#### 82 Nutrient Agar

83 Twenty-eight grams (28g) of nutrient agar was weighed and dissolved in 1000ml of distilled  
84 water in a sterile conical flask. The mixture was heated using a hot plate to dissolve the medium.  
85 The conical flask was plugged with cotton wool stoppers and wrapped with aluminum foil. It  
86 was then sterilized using an autoclave at 121°C for 15 minutes. The medium was cooled at about  
87 45 - 50°C after sterilization and then poured into sterile petri dishes (about 20ml per plate) under  
88 aseptic conditions. The plates were allowed to solidify and incubated at 37°C for 24 hours and  
89 the sterility of the medium was checked [19].

#### 90 3.2.2 Luria-Bertani Agar

91 This media is made up of Tryptone, Yeast extract, NaCl, NaOH and Bacto agar for jelling. Ten  
92 grams (10g) of Tryptone, ten grams (10g) of NaCl, five grams (5g) of NaOH and ten grams  
93 (10g) of yeast extract were weighed and dissolved in 950ml of distilled water in a sterile conical

94 flask. The mixture was heated using a hot plate to dissolve. The final volume was added up to  
95 1000ml and fifteen grams (15g) of Bacto agar was added. The conical flask was plugged with  
96 cotton wool stoppers and wrapped with aluminum foil. It was then sterilized using an autoclave  
97 at 121<sup>0</sup>C for 15 minutes. The medium was cooled at about 45 – 50<sup>0</sup>C after sterilization and then  
98 poured into sterile petri dishes (about 20ml per plate) under aseptic conditions. The plates were  
99 allowed to solidify and incubated at 37<sup>0</sup>C for 24 hours and the sterility of the medium was  
100 checked [20].

#### 101 **Isolation of *Bacillus thuringiensis* from Soil**

102 Five grams (5g) of each soil sample was weighed and added to 100ml of distilled water. The  
103 samples were heated on a hot plate for 10 minutes to eliminate all bacteria incapable of  
104 producing endospores. Since it is known that *Bacillus thuringiensis* produces spores, it will be  
105 safe to assume that if it was present in the soil, it would be in our heated sample. The samples  
106 were then diluted 5 fold to eliminate all humic materials within the samples and to reduce the  
107 overall colony forming units within each sample [18].

#### 108 **Culturing of *Bacillus thuringiensis***

109 The diluted samples were cultured on nutrient agar plates for 24 hours at 37<sup>0</sup>C in to order to give  
110 the spores chance to germinate on media with adequate nutrients and optimal temperature [18].  
111 The media, however, offers favorable growth for a wide range of bacteria as well as *Bacillus*  
112 *thuringiensis*. The colonies were sub-cultured onto Luria-Bertani plates and incubated at 37<sup>0</sup>C  
113 for 24 hours, so as to obtain pure cultures of *B. thuringiensis*. Series of tests which include gram  
114 staining and biochemical tests were further employed to identify *Bacillus thuringiensis* after  
115 formation of colonies with smooth round shape and earthy odor.

116

## 117 **Gram Staining Techniques**

118 A smear of colonies isolated after the identification were made on a clean glass slides using a  
119 sterile wire loop. They were air dried and fixed. The smears were flooded with crystal violet for  
120 about 60 seconds and were washed with tap water. They were then tipped off with lugol's iodine  
121 for 30 seconds and then washed with tap water. They were decolorized with acetone and washed  
122 off with tap water. The fixed smears were counterstained with safranin and allowed for 60  
123 seconds and then washed off with tap water and allow to air dry. Oil immersion was added to the  
124 stained slides and viewed under a microscope using x100 objective for the morphological  
125 characteristics of the isolates [21].

## 126 **Characterization of the Isolated Bacteria**

127 The colonies that form on the T3 agar will again be confirmed by biochemical tests based on:  
128 Indole test, Catalase test, Triple Sugar Iron test (T.S.I), Methyl Red test (M.R.), Vogues-  
129 Proskauer (V.P.).

### 130 **Indole Test**

131 A test tube of sterile peptone water, enriched with 1% tryptophan will be inoculated with young  
132 culture of isolates and incubated at 37<sup>o</sup>C for 48hrs. About 4 drops of kovac's reagent will be  
133 added and shaken gently. A red color will occur immediately at upper part of the test tube  
134 indicating a positive test. A yellow color at the surface will denote a negative result [21].

### 135 **Catalase Test**

136 The container containing 3% hydrogen peroxide solution will be shaken to expel the dissolved  
137 oxygen. One drop of the solution will be placed on a clean glass slide. Presence of gas bubbles  
138 indicates a positive test while absence of gas bubbles indicates negative reaction [19].

139

140 **Triple Sugar Iron Agar (T.S.I test)**

141 A speck of the isolate will be inoculated by streaking and stabbing into the medium and will be  
142 incubated at 37°C for 24 hours. Fermentation of any of the sugar will be indicated by a change in  
143 color, from red to yellow and crack or raised in the medium indicates gas production [21].

144 **Methyl Red Test (M.R.)**

145 A speck of the isolate will be inoculated into the medium, which would be incubated at 37°C for  
146 48 hours. Few drops of methyl red would be added to the culture. M.R positive test will indicate  
147 red color while no changes denotes negative [21].

148 **Vogues-Proskauer Test (V.P)**

149 A speck of the isolate will be inoculated into glucose phosphate water medium and incubated at  
150 37°C for 2 days. Ethanoic solution of 5%  $\alpha$ -naphthol (1.2 ml) and 0.4 ml potassium hydroxide  
151 solution will be added to 2ml of culture and will be shaken vigorously. It will be placed in a  
152 sloping position (for maximum exposure of the culture to air) and will be examined after 30 to  
153 60 minutes. The evolution of red color indicates a positive test for Voges-Proskauer [21].

154 **Coagulase Test**

155 About 2 or 3 colonies was emulsified in 0.05ml of saline contained in a serological tube. 1ml of  
156 plasma was added and incubated at 35<sup>0</sup>-37<sup>0</sup>C and it was checked after 1hour, 2hours, 3hours and  
157 4hours of incubation for signs of clotting of the plasma. Increase in viscosity or complete clotting  
158 indicates a positive coagulase test, while absence of viscosity or clotting indicates a negative  
159 coagulase test [21].

160 **Motility Test**

161 A small quantity of each isolate was stabbed into triple sugar iron agar and incubated at 37°C for  
162 24 hours. Motility was observed by spread of the organism outwards from the stabbed area.

163 **Urease Test**

164 A speck of each isolate was inoculated into Christensen's urea agar and incubated at 37<sup>o</sup>c for 24  
165 hours. Liberation of red color indicates urease positive test while initial yellow color indicates  
166 negative test.

167 **Citrate Utilization Test**

168 A speck of each isolate was inoculated into Koser's citrate medium and was incubated at 37<sup>o</sup>C  
169 for 72 hours. A positive citrate is confirmed by the promotion of blue color while the initial  
170 green color denotes a negative result [21].

171 **Bioassay**

172 The *Bacillus thuringiensis* isolates selected were tested against larva of mosquito. The stock  
173 cultures of *Bacillus thuringiensis* from slant bottles were picked using a sterile wire loop and  
174 diluted five-fold 10<sup>-1</sup>- 10<sup>-5</sup> in sterile distilled water in five test tubes. Five (5) ml each of the  
175 cultures in the first, third and fifth test tubes was added to three (3) disposable cups containing  
176 45ml of sterile distilled water, providing each cup with different dilution factors. Twenty-five  
177 (25) larvae were transferred into each of the disposable cups. The cups were kept at 25<sup>o</sup>C – 30<sup>o</sup>C  
178 for 6 hours. At intervals of 30 minutes, each cup was observed for larval presence and larval  
179 mortality rate was calculated.

180 **RESULTS**

181 The microscopic and biochemical characteristics of the isolated organisms are shown in table  
182 4.1. The characteristics of which confirmed the isolated organism to be *Bacillus thuringiensis*.  
183 Table 4.2 shows the result for the bioassay of 10<sup>-1</sup> diluents of *Bacillus thuringiensis* against  
184 mosquito larvae. The mortality rate was found to increase as the incubation time increases. A  
185 mortality rate of 52% was recorded after 150 minutes and a 100% mortality rate was recorded



186 after 330 minutes. The result illustrated in table 4.3 shows the bioassay for the  $10^{-3}$  diluents of *B.*  
187 *thuringiensis* on mosquito larvae. Mortality rate of 52% was recorded after 180 minutes, after  
188 which a 100% mortality rate recorded after 360 minutes. Illustrated in table 4.4 is the bioassay of  
189 the  $10^{-5}$  diluents of *B. thuringiensis* against mosquito larvae. Mortality rate of 60% was recorded  
190 after 210 minutes, after which a mortality rate of 100% was recorded after 360 minutes.

191

192

193

UNDER PEER REVIEW

194 **Table 4.1: Biochemical and morphological characteristics of *Bacillus thuringiensis***

Isolates	Gram Reaction	Catalase	Coagulase	Glucose	Sucrose	Lactose	Gas	Motility	Citrate	MR	VP	Urease	Indole
<b>A</b>	+ve Rod	+	-	-	-	-	-	+	+	+	-	+	+
<b>B</b>	+ve Rod	+	+	-	-	-	-	+	+	+	-	+	+
<b>C</b>	+ve Rod	+	-	-	-	-	-	+	+	+	-	+	+
<b>D</b>	+ve Rod	+	+	-	-	-	-	+	+	+	-	+	+

195

196 **Key:** + = Positive

197 - = Negative

Time (min)	0	30	60	90	120	150	180	210	240	270	300	330	360
------------	---	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----

<b>No. of larva</b>	25	24	20	17	14	12	10	7	4	2	1	0	0
<b>Control</b>	25	25	25	25	25	25	25	25	25	25	25	25	25
<b>Mortality</b>	0	1	5	8	11	13	15	18	21	23	24	25	25
<b>Mortality rate (%)</b>	0	4	20	32	44	52	60	72	84	92	96	100	100

198

199

200

**Table 4.2: Bioassay of  $10^{-1}$  Diluents of *Bacillus thuringiensis* Culture Against Mosquito Larvae .**

201

202

**Table 4.3: Bioassay of  $10^{-3}$  Diluents of *Bacillus thuringiensis* Culture Against Mosquito Larvae.**

<b>Time (min)</b>	<b>0</b>	<b>30</b>	<b>60</b>	<b>90</b>	<b>120</b>	<b>150</b>	<b>180</b>	<b>210</b>	<b>240</b>	<b>270</b>	<b>300</b>	<b>330</b>	<b>360</b>
<b>No. of larva</b>	25	24	23	21	19	16	12	9	7	6	3	1	0
<b>Control</b>	25	25	25	25	25	25	25	25	25	25	25	25	25
<b>Mortality</b>	0	1	2	4	6	9	13	16	18	19	22	24	25
<b>Mortality rate (%)</b>	0	4	8	14	24	36	52	64	72	76	88	96	100

203

204

205

206

**Table 4.4: Bioassay of  $10^{-5}$  Diluents of *Bacillus thuringiensis* Culture Against Mosquito Larvae.**

<b>Time (min)</b>	<b>0</b>	<b>30</b>	<b>60</b>	<b>90</b>	<b>120</b>	<b>150</b>	<b>180</b>	<b>210</b>	<b>240</b>	<b>270</b>	<b>300</b>	<b>330</b>	<b>360</b>
<b>No. of larva</b>	25	25	24	23	23	21	18	10	6	3	1	0	0
<b>Control</b>	25	25	25	25	25	25	25	25	25	25	25	25	25
<b>Mortality</b>	0	0	1	2	2	4	7	15	19	22	24	25	25
<b>Mortality rate (%)</b>	0	0	4	8	8	16	28	60	76	88	96	100	100

207

208

209

210

211

212

UNDER PEER REVIEW

214

## 215 **DISCUSSION**

216 Mosquitoes are a great nuisance and they pose a serious threat to human health in the society.

217 Many chemical insecticides have been produced for the control of mosquitoes in the past years,

218 some of which have been very effective while others have done little or no good at all. Most of

219 the insecticides used are made of synthetic chemicals and were found to have negative effect on

220 the diversity of many insects, well being of humans and as well the environment. These

221 therefore, call the need for the search of biological control methods that cause less harm to

222 human health, diversity of the insects and the environment. The efficacy of *Bacillus*

223 *thuringiensis* as a larvicide for controlling mosquito larva yielded great results. The microscopic

224 and the biochemical characteristics of the organisms as shown in Table 4.1 confirmed the basic

225 characteristics of *Bacillus thuringiensis*, being Gram positive and having a rod shape. The

226 biochemical characteristics, showed the motile nature of the organism among others. These

227 characteristics are similar to what was reported by Ahmed *et al.* [22].

228 In the bioassay for the control of mosquito larvae, different diluents used showed varying degree

229 of effectiveness with 100% mortality rate recorded after 360 minutes. This might be attributed to

230 the ability of the organism to produce binary toxin (Bin) which is a primary insecticidal

231 component produced during sporulation and vegetative stage of *B. thuringiensis* in controlling

232 the growth of the mosquito larvae. This is in agreement with what was reported by Oei *et al.*

233 [23]. In all the diluents, very low mortality was recorded after 30 minutes of incubation, which

234 could be attributed to the time of exposure of the larvae as well as the number of the organisms

235 present in the container. But more than half of the larvae were death after 240 minutes, with a

236 high mortality rate of over 70% recorded in all the diluents of the *B. thuringiensis*. This could

237 also be attributed to the time of exposure of the larvae and as well the increase in number of the  
238 cells in the medium that could be attributed to the increase in the number of organisms ingested  
239 by the mosquito larvae, which causes damage in the midgut of the larvae [24]. Thereby  
240 releasing the crystallized binary toxins, that in turns are solubilized in the midgut, releasing two  
241 proteins [25], that are cleaved by endogenous proteins to form active toxins [26]. This is in  
242 agreement with what was reported by Aissaoui and Boudjelida [27].

243

## 244 **Conclusion**

245 *Bacillus thuringiensis* naturally found in the soil has proved to be a good larvicidal agent against  
246 mosquito larvae in the laboratory. The organism and its product can be further studied to search  
247 for novel compounds that can be used in control of mosquito-borne diseases such as malaria.

248

## 249 **References**

250

- 251 1. Wilke ABB Marelli MT. Paratransgenesis: A Promising New Strategy for Mosquito  
252 Vector Control. *Parasites and Vectors*. 2015; 8:342. DOI 10.1186/s13071-015-0959-2.
- 253 2. World Health Organization (WHO). Progress Report 2000-2009 and Strategic Plan 2010-  
254 2020 of the Global Program to Eliminate Lymphatic filariasis: Halfway Towards  
255 Eliminating Lymphatic Filariasis. 2010; ISBN 978-92-4-150072-2.
- 256 3. World Health Organization (WHO). WHO Global Malaria Programme. New Report  
257 Signals Showdown in The Fight Against Malaria, 2012. Geneva.
- 258 4. Fonseca DM Smith JL Wilkerson RC Fleischer RC. Pathways of Expansion and Multiple  
259 Introduction Illustrated by large Genetic Differentiation Among World Wide  
260 Populations of the Southern House Mosquito. *American Journal of Tropical Medicine  
261 and Hygiene*. 2006; 74: 284-289.
- 262 5. Bremen JG Egan A Keusch GT. The Intolerable Burden of Malaria: A new Look at the  
263 Numbers. *American Journal of Tropical Medicine and Hygiene*. 2001; 64: iv-vii.
- 264 6. Reiter P. Oviposition, Dispersal and Survival in *Aedes aegypti*: Implications for the  
265 Efficacy of Control Strategies. *Vector-Borne Zoonotic*. 2007; 7: 261-274.

271

272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317

7. Mairiang D Zhang H Sodija A Murali T Suriyaphol P Malasit P *et al.* Identification of New Protein Interactions Between Dengue Fever Virus and its Hosts, Human and Mosquitoes. *PlosOne*. 2013; 8, ee53535.
8. Pidiyar VJ Jangid K Patole MS Shouche YS. Studies on Cultured and Uncultured Microbiota of *Culex quinquefasciatus* Mosquito Midgut Based on 16s ribosomal RNA gene Analysis. *American Journal of Tropical Medicine and Hygiene*. 2004; 70: 597-603.
9. Pates H Curtis CF. Mosquito Behavior and Vector Control. *Annual Review of Entemology*. 2005; 50: 53-70.
10. Vreysen M Robinson AS Hendrichs J. Area-Wide Control of Insect Pests: From Research t Field Implementation. The Netherlands: Springer. 2007; P. 325-361.
11. Dorta DM Vasuki V Rajavel A. Evaluation of Organophosphorus and Synthetic Pyrethroid Insecticides Against Six Vector Mosquito Species. *Rev Saude Publica*. 1993; 27:391-397.
12. Federici BA Hyun-Woo P Dennis KB. Overview of the Basic Biology of *Bacillus thuringiensis* with Emphasis on Genetic Engineering of Bacterial Larvicides foe Mosquito Control. *The Open Toxicology Journal*. 2010; 3: 83-100.
13. Cruz D Ian NB Bautista JR Teves FG. Isolation and Identification of *Bacillus thuringiensis* from *Harpaphe haydeniana* and its Entemotoxic Evaluation Against *Aedes* and *Culex* Larvae. *International Research Journal of Biological Sciences*. 2015; 4(3): 21-26.
14. Federici BA. *Bacillus thuringiensis in Biological Control*. In: Handbook of Biological Control. F Fosher (Ed.) Academic Press (Ed.). 1999; 577-593. ISBN 10: 0-12-257305-6.
15. Otieno-Ayayo ZN Zaritsky A Wirth MC Manasherob R Khasdan V Cahan R Ben-Dov E. Variations in the Mosquito Larvicidal Activities of Toxins from *Bacillus thuringiensis* Spp. *Israelensis*. *Environmental Microbiology*. 2008; 10: 2191-2199.
16. Angolo-Falho RC Loguercio LL. *Bacillus thuringiensis* is an Environmental Pathogen and Host-Specificity has Developed as an Adaptation to Human Generated Ecological Niches. *Insects*. 2014; 5: 62-91.
17. El-Kersh TA Al-akeel RA Al-Sheikh YA. Isolation and Distribution of Mosquitoes Cry genes in *Bacillus thuringiensis* Strains native to Saudi Arabia. *Tropical BioMed*. 2014; 31: 616-632.
18. Travers RS Martin PAW Reichelderfer CF. Selective Process for Efficient Isolation of Soil *Bacillus* sp. *Applied Environmental Microbiology*. 1987; 53: 1263-1266.



- 318 19. Cheesbrough M. District Laboratory Practices in Tropical Countries (Part 2). Cambridge  
319 University Press. 2002; 137-139.  
320
- 321 20. Sambrook J Russell DW. Molecular Cloning a Laboratory Manual. Cold Spring Harbor  
322 Laboratory Press. Cold Spring Harbor, NY. 2001.  
323
- 324 21. Oyeleke SB Manga SB. Essentials of Laboratory Practicals in Microbiology 1<sup>st</sup> Edition.  
325 To Best Publisher Minnaa, Niger State, Nigeria. 2008; 36-58.  
326
- 327 22. Ahmed HA Ali SG Abdul-Raouf UM. Isolation, Characterization and Molecular  
328 Identification of *Bacillus thuringiensis* Alex-13 isolated from Egypt Against *Spodoptera*  
329 *littoralis*. *International Journal of Microbiology and Allied Sciences*. 2015; 2(2): 34-44.  
330
- 331 23. Oei C Hindley J Berry C. Binding of *Bacillus sphaericus* Binary Toxin and its Deletion  
332 Derivatives to *Culex Quinquefasciatus* Gut: Elucidation of Functional Binding Domains.  
333 *Journal of General Microbiology*. 1992; 138: 1515-1526.  
334
- 335 24. Wei D Cai Q Yuan Z. Mosquitocidal Toxin From *Bacillus sphaericus* Induces Stronger  
336 Delayed Effects than Binary Toxin on *Culex quenequefasciatus* (Diptera; Culcidae).  
337 *Journal of Medical Entemology*. 2006; 43: 726-730.  
338
- 339 25. Berry C Jackson Tap J Oei C Hindley J. Nucleotide Sequence of Two Toxin Genes from  
340 *Bacillus sphaericus* IAB59: Sequence Comparisons Between Five Highly Toxicogenic  
341 Strains. *Nucleic Acid Research*. 1989; 17(18): 7516.  
342
- 343 26. Darboux I Nielsen-Leroux C Charles JF Pauron D. The Receptor of *Bacillus sphaericus*  
344 Binary Toxin in *Culex pipens* (Diptera culcidae) midgut: Molecular Cloning and  
345 Expression. *Insect Biochemical and Molecular Biology*. 2001; 31(10): 981-990.  
346
- 347 27. Aissaoui L Boudjelida H. Larvicidal Activity and Influence of *Bacillus thuringiensis*  
348 (Vectobac G), on Longevity and Fecundity of Mosquito species. *European Journal of*  
349 *Experimental Biology*. 2014; 4(1): 104-109.  
350