

**OCCURRENCE OF MULTIDRUG RESISTANCE *ESCHERICHIA COLI* AND OTHER BACTERIA SPECIES ASSOCIATED WITH URINARY TRACT INFECTION IN TWO GEOPOLITICAL ZONES OF ONDO STATE, NIGERIA**

**Abstract**

Urinary tract infection (UTI) is a common bacterial infection known to affect different parts of the urinary tract of both male and female. *Escherichia coli* has been found to be responsible for causing 80% to 90% of the infection. An investigation was carried out to determine the prevalence of bacteria, especially *E.coli* implicated in UTI, and to ascertain their antibiotics susceptibility pattern. Early morning mid-stream urine samples were collected from 250 patients aged 18 to 60 years, between March and July of 2016 from 5 major Hospitals in the study location. The isolates were identified using standard microbiological methods and susceptibility tests were carried out using ten antibiotics. Results showed that 65(30.7%) of the isolates were *E. coli*. *Pseudomonas aeruginosa* 45(21.2%), *Klebsiella pneumoniae* 42(19.8%), *Staphylococcus aureus* 32(15.1%) and *Proteus mirabilis* 28(13.2%). The percentages of resistance of *E. coli* isolates to antimicrobial agents were chloramphenicol (64.9%), sparfloxacin (59.5%), ciprofloxacin (73.0%), septrin (73.0%), amoxicillin (91.9%), augmentin (83.8%), gentamycin (48.7%), perfloxacin (40.5%), ofloxacin (40.5%) and streptomycin (54.1%). The need for constant antimicrobial susceptibility surveillance by health managements system that will help clinicians to provide safe and effective therapy is advocated.

**Key words:** Uropathogen, *Escherichia coli*, Urinary tract infection

**INTRODUCTION**

Urinary tract infection is one of the most frequent bacterial infectious diseases that affect both inpatient and outpatients around the world [8], and is one of the most common diseases that affect people of all ages from neonate to adulthood and globally [12, 30]. UTI is more common among females than male with about 60% of women having at least one episode in their lifetime [21]. This is due to shorter urethra in females which is more readily transverse by microorganisms [53]. The highest incidence of UTI occurs mostly in child bearing age due to the fact of high increase in sexual activity and aging. Gupta [23] reported that about 150 million people are diagnosed with urinary tract infection each year.

*Escherichia coli* have been found to be responsible for 90% of the first urinary tract infections in young women [21]. They are known as a component of the intestinal microflora of human or warm blooded animals [9], which can exist as commensal, it also exist in a symbiotic state

providing resistance against pathogenic organisms or it can be pathogenic and cause diseases of the intestinal and extra intestinal sites

Over 150 million people experienced an episode of UTI yearly and this is costing the world economy over 6 billion US dollars in treatment and workloss [23]. Studies that have being carried out in Enugu, Yola, Zaria and Ile-Ife showed that the same etiological agent was isolated in Nigeria. The prevalence rate of urinary tract infection in Yola was 67.2% and 77% in Enugu [20]. Olowu [41] reported an incidence of 28.1% in a population of 2780 outpatients at Lagos University Teaching Hospital while Aiyegoro [2] reported 11.96% in Ile-Ife. The common bacteria associated with urinary tract infections aside *E. coli* are *Klebsiella*, *Pseudomonas*, *Proteus* and also the Gram positive organism like *Stappylcocci* sp, *Enterococci* sp. *Candida albicans* has been isolated in patients with underlying physiological debilitation [35].

Commensal strains of *E. coli* can also cause diseases in immunocompromised hosts, the pathogenic and commensal *E. coli* are classified into four main phylogenetic groups: A, B1, B2 and D [13]. The phylogenetic group A and B1 consists mainly of the commensal strains that are found in large intestine of humans and animals, but they do not naturally carry any known virulence factor [28]. Whereas, the phylogenetic group B2 and to a lesser extent the group D consist of pathogenic strains that normally carry virulence-associated genes, which are mostly associated with extra-intestinal diseases. The pathogenic *E. coli* strains causes enteric or diarrhoeogenic disease, urinary tract infection and blood infection. The aim of this study is to isolate and identify microorganisms that are associated with urinary tract infection and to assess the antibiotic susceptibility pattern of the isolates.

## **MATERIALS AND METHODS**

### **Sample collection**

A total of 250 urine samples were collected from volunteers (between the ages of 18 and 60 years) that visited the General Hospital, Ikare-Akoko; Federal Medical Center Owo; Adekunle Ajasin University Health Center Akungba-Akoko and Union Diagnostic Center, Akure, Ondo State, Nigeria between Jan and May in 2016.

### **Culture and identification of isolates**

Each sample was inoculated on MacConkey agar, Eosin Methelene Blue (EMB) agar and Cystine Lactose Electrolyte Deficient (CLED) agar plates and incubated at 37<sup>0</sup>C for 24 hours. The identification and characterization of isolated bacteria were done by Gram staining, followed by microscopic examination and biochemical tests such as indole production, citrate utilization and phenol red lactose broth tests.

### **Susceptibility test**

Antimicrobial sensitivity testing of all the isolates was performed using disk diffusion method according to [14]. Clinical samples were inoculated on Mueller Hinton plates and spread evenly to form a lawn culture, after which, the antibiotic discs were placed on the surface and incubated at 37°C for 24 hours. The antibiotics used include: cotrimoxazole (30ug), chloramphenicol (30ug), sparfloxacin (10ug), ciprofloxacin (10ug), amoxicillin (10ug), augmentin (30ug), gentamycin (10ug), perfloxacin (10ug), ofloxacin (10ug) and streptomycin (30ug).

## RESULTS

A total of 212 (84.8%) of urine samples yielded significant growth, out of 155 (62%) females that participated, 150(96.8%) showed positive culture, while 62(65.3%) showed positive culture out of 95(24.8%) males that participated (Table 1). Five different uropathogens were isolated *E.coli* was predominant 65(26%) followed by, *Pseudomonas aeruginosa* 45(18.0%), *Klebsiella pneumonia* 42(16.8%), *Staphylococcus aureus* 32(12.8%), and *Proteus mirabilis* 28(11.2%). Thirty eight (15.2%) samples had no bacteria growth (Table 2). The frequency of bacteria isolated from male and female volunteers was shown in Table 3.

Distribution according to age group revealed that the highest rate of infection occurred in age group 18-45 and the least infection was recorded in volunteers above 46 years (Table 1). The antimicrobial drugs used on the uropathogenic *E.coli* isolated showed that ofloxacin (40.5%) and perfloxacin (40.5%) were the most efficient drugs of choice *in vitro*. This was because they showed high efficacy on the isolates than any other antimicrobials used in this study, this was followed by gentamycin (48.7%). The bacteria showed high degree of resistance to amoxicillin (91.9%), augmentin (83.8%), ciprofloxacin (73.0%), cotrimoxazole (73.0%), chloramphenicol (64.9%) sparfloxacin (59.5%) and streptomycin (54.1%) (Table 4) *Pseudomonas aeruginosa* isolated in this study were resistant to chloramphenicol, sparfloxacin, cotrimoxazole, amoxicillin, augmentin, perfloxacin and streptomycin, but were sensitive to gentamycin, ofloxacin and ciprofloxacin (Table 4). The resistance pattern of *K. pneumonia*, *S. aureus* and *P. mirabilis* were also shown in Table 4, the zone of inhibition was 15 to 40mm.

**Table 1:** Age and Sex distribution of volunteers with significant bacteriuria

Age (years)	Males	Females	Total	Total Percentage
18-45	39	95	134	63.2%
46-60	23	55	78	36.8%
Total	62	150	212	100%
	29.2%	70.8%	84.8%	

N= 250

**Table 2:** Prevalence of microorganisms isolated from urine of patients

Organisms isolated	Number of isolates	Percentage (%)
<i>Escherichia coli</i>	65	26.0
<i>Pseudomonas aeruginosa</i>	45	18.0
<i>Klebsiella pneumonia</i>	42	16.8
<i>Staphylococcus aureus</i>	32	12.8
<i>Proteus mirabilis</i>	28	11.2
No bacteria growth	38	15.2
<b>Total</b>	<b>250</b>	<b>100</b>

**Table 3:** Frequency of isolated bacteria from positive cultures of volunteers

Bacteria isolates	Female	Male	Total (%)
<i>Escherichia coli</i>	49(75.4%)	16(24.6%)	65(30.7)
<i>Pseudomonas aeruginosa</i>	32(71.1%)	13(28.9%)	45(21.2)
<i>Klebsiella pneumonia</i>	28(66.7%)	14(33.3%)	42(19.8)
<i>Staphylococcus aureus</i>	26(81.3%)	6(18.8%)	32(15.1)
<i>Proteus mirabilis</i>	12(42.9%)	16(57.1%)	28(13.2)
<b>Total</b>	<b>147(69.3%)</b>	<b>65(30.7%)</b>	<b>212(100)</b>

119

120 **Table 4:** Antimicrobial sensitivity pattern showing resistance to bacterial agents

Isolates	CH	SP	CPX	SXT	AM	AU	CN	PEF	OFX	S
<i>Escherichia coli</i>	64.9	59.5	73.0	73.0	91.9	83.8	48.7	40.5	40.5	54.1
<i>Pseudomonas auriginosa</i>	65	80	30	55	70	65	25	50	30	65
<i>Klebsiella pneumoniae</i>	50	57.1	50	78.6	78.6	71.4	64.3	35.7	14.3	85.7
<i>Staphylococcus aureus</i>	58.3	66.7	50	58.3	33.3	41.7	41.7	50.0	25.0	66.7
<i>Proteus mirabilis</i>	80	50	30	70	80	80	20	60	30	70

121 CH=Chloramphenicol, SP=Sparfloxacin, CPX=Ciprofloxacin, SXT=Septrin , AM=Amoxacillin,

122 AU=Augmentin, CN=Gentamycin , PEF= Perfloxacin , OFX=Ofloxacin, S=Streptomycin

123 **DISCUSSION**

124 Out of 250 patients (155 females and 95 males between the ages of 18 – 60years), that  
 125 participated in this study, 212(84.8%) urine samples had significant bacteriuria. This was higher  
 126 than the study on profile of urinary tract infection and quinolones resistance among *E.coli* and  
 127 *Klebsiella* spp isolated by Namratha [37], who reported the prevalence of 62%. Also prevalence  
 128 of 67.2% was reported in Yola [20], 60% reported in Lafia [30], 22% in Ibadan [40], 77.9% in  
 129 Enugu [33] and 35.5% in Jos [19] all in Nigeria. The high prevalence observed might be due to  
 130 factors like peer group influence, promiscuity, pregnancy, low socio-economic status that are  
 131 common among Nigerian young men and women [30].

132 This study revealed that *E. coli* was one of the predominant organisms incriminated in urinary  
 133 tract infection. Out of 212 positive isolates *E. coli* had 65(30.7%) followed by *P. aeruginosa*  
 134 45(21.2%), *K. pneumoniae* 42(19.8%), *S. aureus* 32(15.1%) and *P. mirabilis* 28(13.2%). The  
 135 pattern and frequency of occurrence of the bacterial isolates found in this study is similar to those  
 136 reported by other workers. Alausa and Onile [5] reported in their study that *Escherichia coli* was  
 137 the most commonly isolated pathogens in significant bacteriuria. Also, Bishara [10] in Israel  
 138 agreed with this statement, the authors reported that *E. coli* was responsible for 52% of cases of

urinary tract infection, *Klebsiella* spp.(14%), *Proteus* species (95%) and *Enterococcus* spp. was 4%. A variant result was observed in a study in Ile-Ife Southwestern Nigeria by Okonko [41] where *S. aureus* 11(28.9%) was second to *E. coli* 13(42.1%), *Klebsiella* spp 7(18.4%) and *Pseudomonas* spp 2(5.3%). A variant was also observed in a tertiary hospital in Abuja, Central Nigeria, where Iregbu and Nwajiobi-Princewill [25] proved that *E.coli* and *Klebsiella* spp were the two predominant uropathogens 323(37%) and 202(25%) respectively, *P. aeruginosa* 75(8.4%), *Proteus* spp 67(7.5%) and *S. aureus* 60(6.8%). The same trend of *E. coli* leading other uropathogens was reported by [36, 56].

The frequency of urinary tract infections in female was 150(70.8%) and the occurrence in male was 62(29.3%). This finding was in agreement with studies carried out by [24, 41]. This also correlates with other studies by [55, 29] where urinary tract infection was higher in females compared to males. Anochie [7] reported a predominance of female patients in a study carried out to determine the influence of instruction about the method of urine collection and storage on the prevalence of urinary tract infection. Similar findings were reported by [42]. The author reported that urinary tract infection was more frequent in women than men [20, 1, 45]. Okafor [39] reported 20.7% cases of urinary tract infection in males which was similar to 29.3% cases reported in this study.

The higher incidence of urinary tract infection in females has been attributed to the shorter female urethra and its close proximity to the anus that make it easier for enteric flora to colonize this area [15, 49]. Other contributory factors may include the use of contraceptives, childbirth and menopause [44, 17].

The highest rate of infection (UTI) was recorded in volunteers between the age group 18-45 and the least infection (UTI) was recorded in volunteers above 46 years, this is in agreement with the study conducted by [6]. The high incidence of urinary tract infection between the ages of 18-45 years is due to high sexual activities and multiple sex partners among the age group.

The antibiotic resistant *E coli* isolated from urinary tract infections are on high increase and this is a serious problem. Many of the isolated strains are multi-resistance [4, 32]. In this study 54.1% of the *E coli* isolates were noted to be multiple resistant, they are resistant to more than six antibiotics. The antibiotic sensitivity test of this study shows that ofloxacin and perfloxacin were the most efficient antibiotics in vitro testing followed by gentamycin and sparfloxacin, meanwhile most of the organisms are highly resistance to amoxicillin followed by augmentin ciprofloxacin, septrin and chloramphenicol. Sensitivity test in this study revealed that ofloxacin and perfloxacin have the highest efficacy in vitro (59.5%) and (59.5%) respectively, this is in agreement with [26] where ofloxacin had (70.5%) as the most effective among antimicrobials. In this study 91.9% resistance to amoxicillin was found, and this is in agreement with the report of [34] The authors reported 95% resistance to amoxicillin, also Datta [16] reported 88.57% and 100% resistance to amoxicillin was reported by [55].

The percentages of the pathogens resistance varied between 91.9% and 40.5%. In this study, the isolates have demonstrated extremely high percentage of multidrug resistance phenotype. The

rate of resistance to Septrin found is higher than those reported by others [54, 57]. Septrin or trimethoprim alone has been used widely in treating urinary tract infections caused by *E. coli* for decades this study shows low efficacy of septrin also Ciprofloxacin resistance in this study is found to be higher than that reported in previous studies [48].

Current study shows that resistance to Ciprofloxacin is on high increase and if urgent measures are not taken to put the situation under control the need for search of new drugs to combat bacterial infection will arise in Nigeria, meanwhile few ciprofloxacin resistance strains were found in other countries [17].

The high level of resistance seen in Amoxicillin, Augmentin, Septrin, Ciprofloxacin, Chloramphenicol and Sparfloxacin is most likely due to selective pressure that results from uncontrolled and inappropriate use of these agents in our hospitals and in this country as a whole, this is promoted by poverty, lack of awareness about the usage of drugs, lack of antibiotic policy and also the readily availability of antibiotics sold over the counter in Nigeria. The high rate of antimicrobial resistance has pose a problem in the medical sector, 54.1% of our *Escherichia coli* population were noted to be multiple resistance.

In this study *Pseudomonas aeruginosa* isolated were resistance to chloramphenicol, sparfloxacin, septrin, amoxicillin, augmentin, perfloxacin and streptomycin, but were sensitive to gentamycin, ofloxacin and ciprofloxacin. Olowu and Oyetunji [43] also reported multi-resistance in *Pseudomonas aeruginosa* in their study of nosocomial urinary tract infection. *Klebsiella pneumonia* showed high level of resistance to most of the antibiotics used a resistance of 78.6% was seen against amoxicillin, 85.7% were seen against streptomycin, 71.4% were resistance to augmentin and 64.3% to gentamycin while 14.3% were resistance to ofloxacin this result was similar to that reported by [2].

*Staphylococcus aureus* had a resistance pattern of 58.3% to Chloramphenicol, 66.7% to sparfloxacin, 50% to ciprofloxacin, 58.3% to cotrimoxazole, 50.0% to perfloxacin, 66.7% to streptomycin this is contrary to [3] report which presented a report in which 67.3% of *Staphylococcus* spp were resistance to cloxacillin, 64.9% to amoxicillin, 51.8% to augmentin, 70.2% to tetracycline, 48.8% to erythromycin, 36.9% to cotrimoxazole, 11.9% to chloramphenicol and 1.8% to gentamycin in a study in Ibadan Nigeria

## CONCLUSION

The choice of drugs in the treatment of urinary tract infection (UTI) is quite narrow due to the wide -scale resistance that the UTI pathogens exhibited. *E. coli* remains the most frequent bacteria isolate. There is the need for constant antimicrobial sensitivity surveillance which will help clinicians to provide safe and effective therapy. The role of personal hygiene to curtail its spread cannot be overemphasized.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## 215 REFERENCES

- 216 1. Ahmad, S. (2012). Pattern of urinary tract infection in Kashmir and antimicrobial  
217 susceptibility. *Bangladesh Med Res Counc Bull* 38:79-83.
- 218 2. Aiyegoro, O.A., Igbinosa O.O., Ogunmwonyi, I.N., Odjadjare, E.E., Igbinosa, O.E and  
219 Okoh, A.I. (2007). Incidence of urinary tract infections (UTI) among children and  
220 adolescents in Ile-Ife, Nigeria. *African journal of Microbiology Research* pp 013-019.
- 221 3. Ako-Nai, A.K., Adeyemi, F.M., Aboderin, O.A. and Kassim, O.O. (2005). Antibiotic  
222 resistance profile of staphylococci from clinical sources recovered from infants. *Afr. J.*  
223 *Biotechnol.* 4(8): 816-822.
- 224 4. Al-Assil, B., Mahfoud, M. and Hamzeh, A. R. (2013). Resistance trends and risk factors of  
225 extended-spectrum b-lactamases in *Escherichia coli* infections in Aleppo, Syria. *Am J Infect*  
226 *Control* 41, 597–600.
- 227 5. Alausa, K.O and Onile, B.A. (1984). The epidemiological pattern of bacterial septicaemia at  
228 the University College Hospital, Ibadan. *Nig Med. J.* 14: 55-62.
- 229 6. Ani, O.C. and Mgbechi, E.K. (2008). Prevalence of urinary tract infections (UTI) in sexually  
230 active women of Abakaliki, Ebonyi State, Nigeria. *Animal Research International* 5(2):  
231 876-887.
- 232 7. Anochie, I.C, Nkanginieme, K.E.O. and Eke F.U. (2001). The influence of instruction about  
233 the method of urine collections and storage on the prevalence of urinary tract infection. *Nig.*  
234 *J. Paediatr.* 28(2): 39-42.
- 235 8. Basu, S., Mukherjee, S.K., Hazra, A. and Mukherjee, M. (2013). “Molecular characterization of  
236 uropathogenic *Escherichia coli*: nalidixic acid and ciprofloxacin resistance, virulent factors  
237 and phylogenetic background. *Journal of Clinical and Diagnostic Research.* 7(12): 2727-  
238 2731.
- 239 9. Bettelheim, K.A. (1997). *Escherichia coli* in the normal flora of humans and animals. In  
240 *Escherichia coli: Mechanisms of Virulence. Edited by M. Sussman. Cambridge, UK:*  
241 *Cambridge, University Press. pp. 85-109*
- 242 10. Bishara, J, Leibovic, L, Huniner, D, Drucker, M, Samra, Z, Konisberger, H. and Fitlik, S.  
243 (1997). Five years prospective study of bacteraemic urinary tract infection in a single  
244 institution. *Evr. J. Clin. Microbiol. Infect. Dis.* 16 (8): 563-567.
- 245 11. Blumberg, E.A, and Abrutyn, E. (1997). Methods of the reduction of urinary tract infection.  
246 *Curr. Opin. Urol.*; 7: 47-51.
- 247 12. Chang, S.L and Shortliffe, L.D. (2006). Pediatric urinary tract infections. *Pediatr Clin North*  
248 *Am* 53:379-400.
- 249 13. Clermont, O, Bonacorsi, S, and Bingen, E. (2000) Rapid and simple determination of the  
250 *Escherichia coli* phylogenetic group. *Appl Environ Microbiol* 66: 4555-4558.
- 251 14. Clinical and Laboratory Standards Institute. (2014). Performance Standards for  
252 Antimicrobial Susceptibility Testing; Twenty-Third Informal Supplement. *CLSL*  
253 *Document Wayne, PA* M100-S23.



- 254 15. Dai, B., Liu, Y., Jia, J. and Mei, C. (2010). Long-term antibiotics for the prevention of  
255 recurrent urinary tract infection in children: a systematic review and meta-analysis.  
256 *Archives of Disease in Childhood* .95(7):499–508.
- 257 16. Datta, P., Gupta, V. and Sidhu, S.(2014). Extended spectrum beta lactamase positive  
258 uropathogenic *E. coli* – epidemiological factors and resistance. *Br J Med Practic*. 7(2):a718.
- 259 17. Diekema, D.J., Boots, B.J., Miller, T.E., Vaughn, R.F., Woolson, J.W., Yankey, E.J., Ernst,  
260 S.D., Flach, M.M., Ward, C.L., Franciscus, M.A., Pfaller, T.M and Doebbeling, B.N.(2004).  
261 Antimicrobial resistance trends and outbreak frequency in United States hospitals. *Clin.*  
262 *Infect. Dis* 38: 78-85.
- 263 18. Dielubanza, E.J. and Schaeffer, A.J. (2011). Urinary tract infections in women. *The Medical*  
264 *clinics of North America*;95(1): 27–41.
- 265 19. Ebie, M., Kandki-Olukemi, Y.T., Ayanbadejo, J. and Tanyigna, K.B. (2001). Urinary tract  
266 infections in a Nigerian Military Hospital. *Niger. J.Microbiol*. 15(1):31-37.
- 267 20. El-Mahmood, M.A. (2008). Antimicrobial susceptibility pattern of pathogenic bacteria  
268 causing urinary tract infection in a specialist hospital in Yola, Adamawa state. *Journal of*  
269 *Clinical Medicine and Research* 1(1):001-008.
- 270 21. Foxman, B., Barlow, R., D'Arcy, H., Gillespie, B. and Sobel, J.D. (2000) Urinary tract  
271 infection: self-reported incidence and associated costs. *Ann Epidemiol* 10: 509-515.
- 272 22. Geo, F.B., Janet, S.B. and Stephen, A.M. (2001). Urinary tract infections. Enteric gram  
273 negative rods (Enterobacteriaceae). *Jawetz, Melnik and Adelberg's Medical Microbiology*.  
274 22nd Edition. Pp 221.
- 275 23. Gupta, K. (2001). Increasing antimicrobial resistance and the management of uncomplicated  
276 community acquired urinary tract infections. *Int. J. Antimicrob. Agents*. 135: 41-50.
- 277 24. Ibadin, O.M., Peada, O. and Abel, U.G. (2006). Urinary tract infection in Adolescent/ Young  
278 Adult Nigerians with Acquired Human Immuno Deficiency Disease in Benin City. *A peer*  
279 *review journal of Biomedical Sciences* 5(2):55-60.
- 280 25. Iregbu, K.C. and Nwajiobi-Princewill, P.I. (2013). Urinary Tract Infections in a Tertiary  
281 Hospital in Abuja, Nigeria. *African Journal Clinical Experimental Microbiology*,  
282 14(3):169-173.
- 283 26. John, D.O. and Rilwanu, A.M. (2013). Urinary Tract Infections Associated with *Escherichia*  
284 *Coli*: A 2005 to 2009 Clinical Assessment of Trends in Fluoroquinolones Activities in  
285 Maiduguri-City, Nigeria. *Journal of Applied Pharmaceutical*. 3(8): 84-91
- 286 27. Johnson, J.R., Moseley, S.L., Roberts, P.L, and Stamm W.E. (1988). Aerobactin and other  
287 virulence factor genes among strains of *Escherichia coli* causing urosepsis: association with  
288 patient characteristics. *Infect Immun* 56: 405-412.
- 289 28. Johnson, J.R, Delavari, P, Kuskowski, M. and Stell, A.L. (2001). Phylogenetic distribution of  
290 extraintestinal virulence-associated traits in *Escherichia coli*. *J Infect Dis*. 183: 78-88.
- 291 29. Kibret, M. and Abera, B. (2014). Prevalence and antibiogram of bacterial isolates from  
292 urinary tract infections at Dessie Health Research Laboratory, Ethiopia. *Asian pac J Trop*  
293 *Biomed* 4(2):164-168.

30. Kolawole, A.S., Kolawole, O.M., Kandaki-Olukemi, Y., Babatunde, S.K., Durowade, K.A. and Kolawole, C.F.(2009). Prevalence of urinary tract infections (UTI) among patients attending Dalhatu Araf Specialist Hospital, Lafia, Nasarawa State,Nigeria. *Int. J. Med. Sci.*1(1):163-167.
31. Kucheria, R., Dasgupta, P., Sacks, S.H., Khan, M.S. and Sheerin, N.S.(2005). Urinary tract infections: new insights into a common problem. *Postgrad Med J*.81:83-6.
32. Ma, K. L. and Wang, C. X. (2013). Analysis of the spectrum and antibiotic resistance of uropathogens in vitro: results based on a retrospective study from a tertiary hospital. *Am J Infect Control* 41, 601–606.
33. Mbata, T.I. (2007). Prevalence and antibiogram of UTIs among prison inmates in Nigeria.*Int. J. Microbiol.* 3(2): 6-12.
34. Mittal, S., Sharma, M, and Chaudhary, U. (2014). Study of virulence factors of uropathogenic *Escherichia coli* and its antibiotic susceptibility pattern. *Indian J Pathol Microbiol.* 57:61-64.
35. Moore, K.N., Day, R.A. and Albers, M. (2002). Pathogenesis of urinary tract infections: a review. *Journal Clinical Nurse.* 11(5): 568-74.
36. Mubanga, P., Steinberg, W.J. and Van Rooyen, F.C. (2015). Antimicrobial susceptibility profile of uropathogens in Maluti Adventist Hospital patients. *Afr J Prm Health Care Fam Med. Art* 7(1): 800.
37. Namratha, W.N. (2015). Profile of Urinary Tract Infection and quinolones resistance among *Escherichia coli* and *Klebsiella* species isolates. *Int. J. Curr. Microbiol. App. Sci* 4(7):749-756
38. Nataro, J.P and Kaper, J.B. (1998). Diarrheagenic *Escherichia coli*. *Clin. Microbiol. Rev.* 11:142.
39. Okafor, H.V, Okoro, B.A and Ibe, B.C. (1993). Prevalence of asymptomatic bacteriuria among nursery school children. *Nig. J. Paediatr.* 20: 84-88.
40. Okesola, A.O. and Oni, A.A. (2009). Antimicrobial resistance among common bacterial pathogens in south western Nigeria. *American-Eurasian J.Agric. Environ. Sci.* 5(3): 327-330.
41. Okonko, I.O., Donbraye-Emmanuel, O.B., Ijandipe, L.A., Ogun, A.A., Adedeji, A.O. and Udeze, A.O.(2009). Antibiotics Sensitivity and Resistance Patterns of Uropathogens to Nitrofurantoin and Nalidixic Acid in Pregnant Women with Urinary Tract Infections in Ibadan, Nigeria. *Middle-East Journal of Scientific Research* 4(2):105-109
42. Olowu, W.A.(1996). Symptomatic bacteriuria in children.*Nig. Med Pract.*27: 76-80.
43. Olowu, W.A and Oyetunji, T.G. (2003). Nosocomial significant bacteriuria prevalence and pattern of bacterial pathogens among children hospitalized for non-infective urinary tract disorders. *West Afr. J. Med.* 22 (1): 72-75.
44. Phipps, S, Lim, Y.N, McClinton, S, Barry, C, Rane A, and N'Dow, J. (2006). Short term urinary catheter policies following urogenital surgery in adults. In Phipps, Simon."Cochrane Database of Systematic Reviews". *Cochrane Database Syst Rev*; (2):43-74.

- 334 45. Randrianirina, F., Soares, J.L., Carod, J.F., Ratsima, E., Thonnier, V., Combe, P., Grosjean,  
335 P. and Talarmin, A.J. (2007). Antimicrobial resistance among uropathogens that cause  
336 community acquired urinary tract infections in Antananarivo, Madagascar.  
337 *Antimicrob Chemother* 59(2):309-312.
- 338 46. Rijavee, M, Müller-Premru, M, Zakotnik, B. and Žgur-Bertok, D. (2008). Virulence factors  
339 and biofilm production among *Escherichia coli* strains causing bacteraemia of urinary tract  
340 origin. *J Med Microbiol* 57: 1329-1334.
- 341 47. Rozen, S. and Skaletsky, H. (1999). Bioinformatics methods and protocols on Primers 3 on the  
342 WWW for general users and for Biologist programmers. *Methods Mol Biol* 2000(132):365-  
343 386
- 344 48. Sahm, D.F., Thornsberry, C., Mayfield, D.C., Jones, M.E. and Karlowsky,  
345 J.A. (2001). Multidrug-resistant urinary tract isolates of *Escherichia coli*: prevalence and  
346 patient demographics in the United States in 2000. *Antimicrob. Agents Chemother* 45:1402-  
347 1406.
- 348 49. Salvatore, S., Cattoni, E., Siesto, G., Serati, M., Sorice, P. and Torella, M. (2011). Urinary  
349 tract infections in women. *European journal of Obstetrics, Gynecology, and Reproductive*  
350 *Biology* 156(2):131-6.
- 351 50. Scholes, D, Hooton, T.M, Roberts, P.L, Stapleton, A.E, Gupta, K. and Stamm, W.E. (2000).  
352 Risk factors for recurrent urinary tract infection in young women. *J Infect Dis* 182:1177-82.
- 353 51. Stecher, B, and Hardt, W.D. (2008). The role of microbiota in infectious disease. *Trends*  
354 *Microbiol* 16(3):107-114
- 355 52. Swetha, V.V., Rao, U.S., Prakask, P.H. and Subbarayudu, S. (2014). Aerobic Bacteriological  
356 Profile of Urinary Tract Infections in a Tertiary Care Hospital. *Int J Curr Microbiol App Sci*  
357 3(3):120-125.
- 358 53. Uehling, D.T. (1995). Vaginal mucosa immunization in recurrent urinary tract infections.  
359 *Infect. Urology vol* 57-61.
- 360 54. Valdivieso, F., Trucco, O., Prado, V., Diaz, M.C., and Ojeda, A. (1999). Antimicrobial  
361 resistance of agents causing urinary tract infections in 11 Chilean hospital. *PRONARES*  
362 *Project. Rev. Med. Chil.*, 127(9): 1033-1040.
- 363 55. Xavier, A.M. and Amutha, S. (2016). Antibiotic resistance, RAPD-PCR Typing of Multidrug  
364 Resistant strains of *Escherichia coli* from Urinary Tract Infections (UTI). *Journal of clinical*  
365 *and diagnostic research dol.* 10(3):5-9.
- 366 56. Yabaya, A., Lucy, M. and Maimuna, H. (2012). Bacteria associated with urinary tract  
367 infections and their Susceptibility pattern to routine antibiotics. *Journal of medical and*  
368 *applied biosciences* 4(62):1360
- 369 57. Zhanel, G.G., Karlowsky, J.A., Harding, G.K.M., Carrie, A, Mazzulli, T. and Low, D.E.  
370 (2000). A Canadian national surveillance study of urinary tract isolates from outpatients:  
371 comparison of the activities of trimethoprim-sulfamethoxazole, ampicillin, mecillinam,  
372 nitrofurantoin and ciprofloxacin. *Antimicrob. Agents Chemother.* 44: 1089-1092.

373