

## Original Research Article

### DOSE-DEPENDENT CHEMOPREVENTIVE EFFECT OF METHANOL EXTRACT OF *CARICA PAPAYA* SEED ON POTASSIUM BROMATE- INDUCED NEPHROTOXICITY IN RATS

**Comment [Office1]:** Comment: add latin name?  
The words of manuscript is between 2000-2500

#### ABSTRACT

The dose-dependent preventive effect of methanol extract of *Carica papaya* seed was investigated on potassium bromate-induced nephrotoxicity in rats. The animals were concurrently administered a single oral dose of 100mg/kg body weight potassium bromate and varying dosages of 200mg/kg, 400mg/kg and 600mg/kg body weight methanol extract of *Carica papaya* seed obtained by maceration, and the effect examined 48hours after administration. The result showed significant ( $P<0.05$ ) decreases in serum levels of urea, creatinine, uric acid and all electrolytes studied in a dose-dependent fashion from 200mg/kg to 400mg/kg and 600mg/kg in that order when compared to control. Furthermore, the activities of biomarker enzymes of the renal brush border membrane namely  $\gamma$ -glutamyltransferase (GGT), alkaline phosphatase (ALP), maltase (Mal) and leucine aminopeptidase (LAP) in homogenates prepared from renal cortex and medulla were significantly ( $P<0.05$ ) decreased with  $KBrO_3$  administration but this was prevented by methanol extract of *Carica papaya* seed in a dose dependent fashion. Furthermore, there were significant changes in parameters of oxidative stress. While the activities of the antioxidant enzymes studied namely; catalase, superoxide dismutase and glutathione peroxidase as well as reduced glutathione were significantly decreased, the level of malondialdehyde significantly ( $P<0.05$ ) increases in the renal homogenates following  $KBrO_3$  administration but these were prevented by concurrent administration of  $KBrO_3$  and methanol extract of *Carica papaya* seed in a dose dependent manner. It was suggested that *Carica papaya* seed extract can ameliorate potassium bromate- induced renal toxicity and oxidative stress by preventing the changes in all the kidney function parameters studied and improving the antioxidant defense system respectively.

Keywords: Chemopreventive, *Carica papaya* seed, Potassium bromate, Nephrotoxicity

**Comment [Office2]:** Comment: please, structure in abstract is arranged with aim, material, method, result and conclusion respectively

#### 1. INTRODUCTION

Potassium bromate is a white crystalline substance that is used as food additive in bread, as improver and a dough conditioner in flour. The cosmetics industry also uses bromate as component of permanent hair waving solutions. Potable water disinfection by ozonation, which has been preferred as a better method of sterilization vis-à-vis chlorination because it does not precipitate hazardous substances such as trihalomethanes, also generates bromated [1]. Ingestion of  $KBrO_3$  can cause toxicity of several organs with the kidney being the primary target.  $KBrO_3$  causes mutation in renal tissues and ingestion of higher doses of the substance over a long period induces carcinomas in rats, hamsters and mice [2]. The International Agency for Research on Cancer, IARC has classified bromate as a probable human carcinogen and a complete carcinogen in animals [3]. Researchers have suggested upsurge in production of reactive oxygen species (ROS) and free radicals as fundamental in mediating  $KBrO_3$ -induced toxicity [4]. Free radicals can cause tissue damage when they attack macromolecules such as proteins, nucleic acids and membrane lipids and leads to imbalance in homeostasis thus causing tissue injury [5]. Lethal oral doses of bromate in humans range from 154 and 385 mg/kg body weight while

**Comment [Office3]:** Comment: please mention short name this agent with ( $KBrO_3$ ) for first after that with with short name

45 serious poisoning occurs at doses of 46–92 mg/kg body weight [6]. Oral doses of 185–385 mg/kg body  
46 weight can cause irreversible toxic effects like renal failure and deafness in humans while lower doses  
47 are associated with vomiting, diarrhea, nausea and abdominal pain [6].

48 *Carica papaya* is a fast growing tree-like herbaceous plant in the family *caricaceae* with four genera. The  
49 genus *Carica linn* is the most common of the *Carica papaya* and is the most widely cultivated and best  
50 known species [7]. Economically, *Carica papaya* is the most important species within the *caricaceae*  
51 being widely cultivated for consumption as fresh drinks, jams and candies or as dried and crystallized  
52 fruit. The green fruit leaves and flowers are also cooked as vegetable [8]. Several workers have  
53 reported the therapeutic activities of *Carica papaya* seed to include anti-helminthic action [9], antibacterial  
54 [10], antifungal [11] and female anti fertility [10] among others. Seeds of *carica papaya* have also been  
55 used for centuries in folk medicine for the treatment of poison-related renal disorders among some ethnic  
56 groups in Nigeria and therefore could be considered a potential candidate for chemoprevention of the  
57 kidney against  $KBrO_3$ -induced nephrotoxicity hence the need for the present study [6].

## 58 2. MATERIALS AND METHODS

### 59 2.1 Chemical and Assay Kits

60 Epinephrine, reduced glutathione, tris (hydroxymethyl) aminomethane (Tris), [2-[4-(2-hydroxyethyl)-1-  
61 piperazinyl]ethanesulfonic acid], HEPES, Trichloroacetic acid (TCA), hydrogen peroxide,  $H_2O_2$ ,  
62 Thiobarbituric Acid (TBA). Potassium bromate was supplied by Labtech Chemicals Lagos, Nigeria. The  
63 assay kits for urea, creatinine, uric acid, sodium, potassium, chloride and bicarbonate were all obtained  
64 from Randox laboratories Ltd UK. All other chemicals used were of analytical grade

### 66 2.2 Plant Sample and Extraction

67  
68 65 matured unripe *Carica papaya* was bought from Na'ibawa fruit market Kano, Nigeria and identified at  
69 the plant Herbarium, Department of Plant Biology, Bayero University Kano, Nigeria with an accession  
70 number BUKHAN 0012. Each of the samples was cut to remove the seeds which was washed with tap  
71 water, shade-dried and ground into fine powder with an electric blender. Maceration as described by [12]  
72 was used. 500 g of the powdered dried *Carica papaya* seed was suspended in 1200 ml of methanol for  
73 24 hours and shaken at regular intervals. The extract was then sieved first with cheese cloth and then  
74 with Watmann filter paper No 1. The filtrate was concentrated to dryness in a water bath preset at  $50^\circ C$   
75 and was kept in the refrigerator at  $4^\circ C$  until required.

### 77 2.3 Experimental Animal

78  
79 Thirty (30) apparently healthy young male Albino Wister rats, each weighing between 120-150g were  
80 used for the study. The study was carried out at the animal house unit of the department of Biological  
81 Sciences, Bayero University Kano, Nigeria. All animal procedures were performed according to the guide  
82 for the care and use of laboratory animals of the National Institute of Health as well as the Animal Welfare  
83 Act. Prior to the experiment, the animals were allowed to acclimatize to the animal house for one week  
84 and were maintained on standard pellet rat diet with free access to water.

### 86 2.4 Experimental Design

87  
88 By the end of the seven days acclimatization period, the animals were randomly divided into six groups  
89 as shown below. Each group contains five rats. Solution of potassium bromate was administered orally as  
90 a single dose of 100mg/kg body weight to rats in the tests and  $KBrO_3$  control groups. Methanol extract of  
91 *Carica papaya* seed was reconstituted in distilled water and administered concurrently to animals in the  
92 tests groups in dosages of 200mg, 400mg and 600mg/kg body weight of rats while animals in the normal  
93 control group was administered equivalent volume of distilled water  
94 Group one, normal control: given distilled water  
95 Group two,  $KBrO_3$  control: given  $KBrO_3$ , 100mg/kg bw  
96 Group three, *papaya* control: given 600mg/kg bw CPS

**Comment [Office4]:** Comment please report the statistical report of the study. Please report what is the purpose of the study<

**Comment [Office5]:** Comment: no needed

97 Group four, treatment: given 200mg/kg bw CPS + 100mg/ kg bw KBrO<sub>3</sub>  
98 Group five, treatment given 400mg/kg bw CPS + 100mg/ kg bw KBrO<sub>3</sub>  
99 Group six, treatment: given 600mg/kg bw CPS + 100mg/ kg bw KBrO<sub>3</sub>  
100

#### 101 **2.4.1 Collection of blood sample**

102 All the animals were sacrificed by decapitation, 48 hours after the respective treatment and blood  
103 samples were collected in lithium heparin tubes and centrifuged at 4000 rpm for 5 minutes to collect the  
104 serum which is stored at 4C

#### 105 **2.4.2 Preparation of renal homogenates**

106 The renal homogenates were prepared using the method of [4]. After the animal sacrifice, the kidneys  
107 were excised, bisected and kept in ice-cold 154mM NaCl and 5 mM Tris-HEPES buffer, pH 7.5. The  
108 cortex and medulla were carefully separated using a sharp scalpel and homogenized separately in a  
109 glass Teflon homogenizer in 2 mM Tris-HCl, 50mM mannitol buffer, pH 7.0, to get a 10% (w/v)  
110 homogenate. These homogenates were diluted to 5% with Tris-mannitol buffer followed by high speed  
111 homogenization (20,000 rpm) in an Ultra Turrex Kunkel homogenizer. The renal homogenates were  
112 divided into aliquots and frozen immediately pending analysis.  
113  
114  
115

**Comment [Office6]:** Comment: please mention what's methods?

#### 116 **2.4.3 Determination of biochemical Parameters**

##### 117 **Urea, creatinine and uric acid**

118 Urea was determined in serum by the diacetyl monoxime method using kit from Randox Laboratories Ltd,  
119 UK. Creatinine level was determined in deproteinized serum based on its reaction with saturated picric  
120 acid to give a yellow-red complex using kits from Randox Laboratories Ltd, UK while uric acid level was  
121 determined by the measurement of quinoneimine dye complex using kit from Linear Chemicals  
122 Barcelona, Spain.  
123  
124  
125

##### 126 **Electrolytes**

127 Na<sup>+</sup>, K<sup>+</sup>, Cl<sup>-</sup> and HCO<sub>3</sub><sup>-</sup> were all estimated in serum by spectrophotometric measurement using kits from  
128 Teco Diagnostics Anaheim, USA  
129  
130

##### 131 **Renal brush border membrane enzymes**

132 GGT (EC 2.3.2.2) was determined in the homogenates of renal cortex and medulla by colorimetric  
133 method using kit from Spectrum Diagnostic, Germany. The reaction is based on the measurement of  
134 chromogen p-nitroanilide at wavelength of 418nm. ALP (EC 3.1.3.1) was determined by colorimetric  
135 method by measuring of an intense yellow colour complex, p-nitrophenol using kit from Dialab Production  
136 Neudorf, Austria while maltase (3.2.1.20) and LAP (3.4.11.1) were determined using kits from  
137 Elabscience Biotechnology Inc, USA and Bioway Nanjiang, China respectively  
138  
139

##### 140 **Parameters of oxidative stress**

141 The parameters that show the induction of oxidative stress determined include catalase (CAT) (EC  
142 1.11.1.6), superoxide dismutase (SOD) (EC 1.15.1.1), glutathione peroxidase (GPx) (EC 1.11.1.9),  
143 reduced glutathione (GSH) and malondialdehyde (MDA). All the parameters were determined in  
144 homogenates prepared from renal cortex and medulla separately. CAT activity in renal tissues were  
145 determined by the quantitation of chromic acetate formed at pH 7.0 according to the method of Singha  
146 [13] while SOD activities were determined by the method of Misra and Fridovich [14] by inhibition of auto  
147 oxidation of epinephrine at pH 10.2. GPx activities were determined by the splitting of H<sub>2</sub>O<sub>2</sub> with oxidation  
148 of GSH at pH 7.4 using the method of Rotruck *et. al.* [15] while the levels of GSH were quantified in  
149 deproteinised samples by measurement of 5', 5'-dithiobis-(2-nitrobenzoic acid) (DTNB) using the method  
150

151 of Beutler *et. al.* [16]. Malondialdehyde was determined by the measurement of thiobarbituric acid  
152 reactive substances (TBARS) as described by Varshney and Kale [17].

## 154 2.5 Statistical Analysis

155  
156 Results are expressed as mean  $\pm$  SDM and n =5 for all readings. One-way analysis of variance (ANOVA)  
157 was used to analyzed data and a difference of (P<0.05) was considered significant.

## 159 2.6 Histological Examination

160  
161 The method of Krause [18] was employed in the examination of the kidney tissues. The  
162 photomicrographs were observed using Leitz, DIALUX research microscope at x100 magnification.

# 164 3. RESULTS AND DISCUSSION

**Comment [Office7]:** Comment: please separate between result and discussion?  
Discussion contain the result the study compared other study

## 166 3.1 Kidney Function Parameters

167  
168 Administration of KBrO<sub>3</sub> significantly (P<0.05) increases the serum levels of urea, creatinine and uric acid  
169 however concurrent administration of KBrO<sub>3</sub> and *Carica papaya* seed extract resulted in significant  
170 (P<0.05) decreases in these parameters towards normal control values in a dose-dependent fashion from  
171 200mg/kg to 400mg/kg and 600mg/kg in that order. The parameter most affected by KBrO<sub>3</sub> was creatinine  
172 and the least affected was uric acid. Administration of *Carica papaya* seed extract alone did not give any  
173 significant change in all the kidney function parameters studied.

175 Table 1: Dose-dependent effect of concurrent administration of potassium bromate and methanol extract  
176 of *Carica papaya* seed on kidney function parameters of rats

	Normal control	KBrO <sub>3</sub> control	<i>Papaya</i> control	200mg/kg	400mg/kg	600mg/kg
Urea (mmol/l)	8.44 $\pm$ 0.56	14.82 $\pm$ 0.53*	8.52 $\pm$ 0.33	8.31 $\pm$ 0.31	8.47 $\pm$ 0.09	8.24 $\pm$ 0.51*
Creatinine(mmol/l)	3.80 $\pm$ 0.57	7.07 $\pm$ 0.25*	3.59 $\pm$ 0.33	3.52 $\pm$ 0.42	3.21 $\pm$ 0.24	3.20 $\pm$ 0.25*
Uric acid (mmol/l)	5.49 $\pm$ 0.21	5.94 $\pm$ 0.19*	5.56 $\pm$ 0.18	5.81 $\pm$ 0.11	5.69 $\pm$ 0.73	5.62 $\pm$ 0.21

177 n = mean  $\pm$ SD of five different samples; \* significantly different (P<0.05) from normal control

178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190

## 181 3.2 Serum Electrolytes

183 Potassium bromate administration resulted in significant (P<0.05) increases in serum levels of all the  
184 electrolytes studied with HCO<sub>3</sub><sup>-</sup> being the most adversely affected followed by K<sup>+</sup> and Cl<sup>-</sup>. Na<sup>+</sup> was the  
185 least affected electrolyte by KBrO<sub>3</sub> administration. However concurrent administration of KBrO<sub>3</sub> and  
186 methanol extract of *Carica papaya* seed led to significant decreases in the serum levels of all the  
187 electrolytes studied in a dose-dependent fashion with 600mg/kg being the most active dosage.  
188 Administration of *Carica papaya* seed extract alone did not significantly affect any of the electrolytes  
189 studied.

191  
192

193 Table 2: Dose-dependent effect of concurrent administration of potassium bromate and methanol extract  
194 of *Carica papaya* seed on serum electrolytes of rats

	Normal control	KBrO <sub>3</sub> control	<i>Papaya</i> control	200mg/kg	400mg/kg	600mg/kg
Na <sup>+</sup> (mmol/l)	139.86±2.01	144.76±2.09*	138.48±2.35	142.02±2.34	141.77±1.59	140.75±1.67
K <sup>+</sup> (mmol/l)	8.97±0.28	24.89±0.43*	9.19±0.52	11.49±0.99	9.33±0.77*	9.05±0.28*
Cl <sup>-</sup> (mg/dl)	103.83±3.50	143.60±3.11*	103.46±3.77	106.61±4.27	104.27±4.04	100.25±4.02*
HCO <sub>3</sub> <sup>-</sup> (mg/dl)	5.45±0.56	23.69±1.68*	5.15±0.52	5.67±0.89	5.32±0.73	5.27±0.46

195 n = mean ±SD of five different samples; \* significantly different (P<0.05) from normal control

196 .

### 197 3.3 BBM Marker Enzymes

198

199 Administration of KBrO<sub>3</sub> significantly (P<0.05) decreases the activities of all the BBM marker enzymes,  
200 namely γ-glutamyltransferase, alkaline phosphatase, maltase and leucine aminopeptidase. The effect  
201 was observed in both in cortex and medulla, with the cortex being more extensively affected than medulla  
202 for all the enzymes. The BBM enzyme most affected by KBrO<sub>3</sub> was ALP followed by maltase and LAP.  
203 GGT was least affected. However concurrent administration of KBrO<sub>3</sub> and methanol extract of *Carica*  
204 *papaya* seed resulted in significant (P<0.05) increases in activities of all the BBM marker enzymes toward  
205 normal control values in a dose-dependent fashion. There was no any change in the *papaya* control  
206 group

207

208 Table 3: Dose-dependent effect of concurrent administration of potassium bromate and methanol extract  
209 of *Carica papaya* seed on activities of marker enzymes in brush border membrane in homogenates of  
210 renal cortex and medulla of rats

	Normal control	KBrO <sub>3</sub> control	<i>Papaya</i> control	200mg/kg	400mg/kg	600mg/kg
GGT (U/L)						
Cortex	5.71±0.45	3.36±0.37*	5.52±0.52	4.50±0.52	5.54±0.42*	5.74±0.15*
Medulla	3.97±0.41	1.27±0.54*	3.78±0.15	3.73±0.11*	3.55±0.15*	3.62±0.48*
ALP (U/L)						
Cortex	6.48±0.77	2.32±0.62*	6.28±0.65	4.52±0.48	6.17±0.55	6.58±0.33*
Medulla	4.83±0.37	1.33±0.44*	4.69±0.80	3.04±0.11	4.03±0.12*	5.51±0.45*
Maltase (U/L)						
Cortex	25.87±0.87	11.91±0.95*	26.07±0.57	25.05±0.04	26.52±0.11	26.95±0.20*
Medulla	18.54±0.53	8.14±0.41*	18.64±0.92	16.08±0.10	18.16±0.69*	18.53±0.54*
LAP (U/L)						
Cortex	5.49±0.50	3.42±0.48*	5.30±0.34	4.36±0.35	4.86±0.18	5.11±0.34*
Medulla	3.92±0.30	2.46±0.50*	3.86±0.44	4.30±0.21	4.31±0.20	4.41±0.40

211 n = mean±SD of five different samples, ALP= Alkaline phosphatase, GGT= γ-glutamyltransferase, LAP =  
212 leucine aminopeptidase; \*significantly different (P<0.05) from normal control

213

### 214 3.4 Parameters of Oxidative Stress

215

216 Administration of KBrO<sub>3</sub> resulted in significant (P<0.05) increase in MDA concentration in the  
217 homogenates of both renal cortex and medulla of rats as compared to control values however  
218 concurrent administration of KBrO<sub>3</sub> and methanol extract of *Carica papaya* seed resulted in significant  
219 (P<0.05) decrease in the level of MDA toward normal control values in both the cortex and medulla in  
220 a dose-dependent fashion.

221

222 The result of the antioxidant enzymes studied (CAT, SOD, GPx) and GSH also showed significant  
223 decreases in all the enzyme activities and the level of GSH in both cortex and medulla with the cortex  
224 being extensively affected than the medulla. However concurrent administration of KBrO<sub>3</sub> and  
225 methanol extract of *Carica papaya* seed resulted in significant (P<0.05) increases in activities of all  
226 the AO enzymes studied and GSH in a dose dependent manner. Administration of *Carica papaya*  
227 seed extract alone did not significantly affect any of the parameters of OS studied

228

229 Table 4: Dose-dependent effect of concurrent administration of potassium bromate and methanol extract  
 230 of *Carica papaya* seed on parameters of oxidative stress in homogenates of renal cortex and medulla of  
 231 rats

	Normal control	KBrO <sub>3</sub> control	<i>Papaya</i> control	200mg/kg	400mg/kg	600mg/kg
CAT (units/mg prtn)						
Cortex	71.76±2.48	44.92±1.24*	72.76±1.46	56.56±1.51	63.06±1.15*	75.35±1.43*
Medulla	42.67±1.83	22.86±1.21*	43.56±1.14	36.01±0.73*	42.50±1.66*	43.38±0.30*
SOD (units/mg prtn/min)						
Cortex	21.16±1.70	13.58±0.56*	20.64±1.28	19.37±0.51	20.35±1.38	20.55±1.43*
Medulla	12.84±0.42	7.77±0.69*	12.74±0.61	11.81±0.43	11.42±1.69	11.44±1.57*
GPx(units/mg prtn)						
Cortex	49.49±1.11	24.89±0.41*	49.74±1.24	50.43±2.01	42.52±0.81	50.86±0.71*
Medulla	18.27±0.92	12.45±1.34*	19.63±0.94	19.14±1.27	18.32±1.27*	19.49±0.25*
GSH(μmol/min T)						
Cortex	3.15±0.57	0.54±0.09*	3.45±0.52	1.49±0.34	1.42±0.08	1.48±0.05*
Medulla	2.30±0.32	0.42±0.03*	2.55±0.63	0.56±0.23	1.37±0.11	1.52±0.21*
MDA(units/mg prtn)						
Cortex	15.41±1.01	32.50±0.71*	15.47±1.46	16.28±0.59	15.76±0.12*	15.29±0.63*
Medulla	8.18±0.63	23.39±1.11*	8.47±1.07	8.32±0.07	8.07±0.22	8.14±0.49*

232 n = mean±SD of five different samples, CAT = Catalase, SOD= Superoxide dismutase, GPx = glutathione  
 233 peroxidase; \*significantly different (P<0.05) from normal control

234  
 235  
 236  
 237  
 238  
 239  
 240  
 241  
 242  
 243  
 244




Plate 1: micrograph of rat kidney administered with distilled water (Mag. x100)  
 Showing normal architecture with intact Bowman's capsule, tubular epithelium and basement membrane

BM = basement membrane, BMC = Bowmans capsule, PCT = Proximal convoluted tubule

245  
 246  
 247  
 248

249  
250  
251  
252




Plate 2: micrograph of rat kidney administered with *Papaya* (Mag. x100) Showing normal architecture similar to normal control with intact Bowman's capsule, tubular epithelium and basement membrane




Plate 3: micrograph of rat kidney administered with  $KBrO_3$  (Mag. x100) Showing changes in glomerulus such as irregular dilatation of tubules, and distortions on the distal and proximal tubules and damaged basement membrane

Plate 4: micrograph of rat kidney administered with  $KBrO_3 + 200mg$  (Mag. x100) showing nearly normal architecture with intact Bowman's capsule, and tubular epithelium but basement membrane has not returned to normal

266  
267  
268  
269  
270

$KBrO_3$   
control




Plate 5: micrograph of rat kidney administered with  $KBrO_3 + 400mg$  (Mag. x100) showing nearly normal architecture with intact Bowman's capsule, and tubular epithelium but basement membrane has not returned to normal

271  
272  
273  
274  
275




Plate 6: micrograph of rat kidney administered with  $KBrO_3 + 600mg$  (Mag. x100) showing nearly normal architecture with intact Bowman's capsule, and tubular epithelium but basement membrane has not returned to normal

276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302

The kidney is exposed regularly to high level of reactive oxygen species (ROS) and therefore requires a functional antioxidant defense system to protect its structure and function, and maintain its metabolic properties [19]. This is important because redox interruptions are reported to have devastating effect on body system because it can generate ROS which can attack and modify macromolecules such as proteins, lipids and DNA [20]. Researchers have linked several environmental pollutants, therapeutic drugs and certain food additives such as  $KBrO_3$  with increase generation of ROS and have outlined some key events that are involved in renal damage and antioxidant defense mechanism [21, 22, 23, 24]. Thus, it is important to search for antioxidants that have ability to prevent the damaging effect of various toxicants' induced oxidative stress and the accompanying metabolic disorders. *Carica papaya* seed known for its uses in folk medicine for the treatment of poison-related renal disorders by several ethnic groups across Nigeria was used in the present study and its nephroprotective potential was investigated. A single oral dose of  $KBrO_3$  was used to induce nephrotoxicity in rats and seeds from matured, unripe *Carica papaya* fruit were shade-dried, ground into fine powder, extracted with methanol and tested for potency against the  $KBrO_3$ -induced nephrotoxicity in rats. The organic extract of the dried seed was found to prevent the devastating effect of  $KBrO_3$  on the kidney of rats studied

**Comment [Office8]:** Comment: 21-24

The observed significant ( $P < 0.05$ ) increases in serum urea, creatinine and uric acid levels in  $KBrO_3$  administered rats could be due to the kidney's inability to carry out its functions of filtration and transport of metabolites as a result of the morphological changes on the kidney tissues following administration of



303 KBrO<sub>3</sub>. Histological findings showed alterations in the glomerulus such as irregular dilatation of the  
304 tubules and necrosis leading to the distortion of the glomerular basement in KBrO<sub>3</sub> administered rats  
305 (plate 3). These could alter the normal physiology of the kidney and lead to changes such as alteration in  
306 intraglomerular hemodynamics [25, 26], decrease in renal blood flow and glomerular filtration rate (GFR)  
307 and therefore leading to reduced uric acid and creatinine clearance [26] hence the accumulation of the  
308 kidney function parameters observed however, concurrent administration of KBrO<sub>3</sub> and methanol extract  
309 of *Carica papaya* seed prevented these changes and restored the values towards normal control.

310  
311 The increased levels of serum Na<sup>+</sup>, K<sup>+</sup>, HCO<sub>3</sub><sup>-</sup> and Cl<sup>-</sup> corroborated the observed increases in the serum  
312 levels of urea, creatinine and uric acid. The kidney is responsible for the regulation of various electrolytes  
313 and maintenance of homeostasis [27], for example Na<sup>+</sup> and K<sup>+</sup> are major components of extracellular and  
314 intracellular fluids respectively and these physiological state are regulated by the kidney, therefore the  
315 elevated levels of these electrolytes could indicate renal dysfunction particularly at glomerular and tubular  
316 levels. Also, HCO<sub>3</sub><sup>-</sup> and Cl<sup>-</sup> are reabsorbed in the proximal tubule and therefore the elevated level of these  
317 electrolytes could imply renal damage at the tubular level. However the serum levels of all the electrolytes  
318 studied were restored towards normal with concurrent administration of KBrO<sub>3</sub> and methanol extract of  
319 *Carica papaya* seed

320  
321 The significant decreases (P<0.05) in the activities of all the brush border membrane (BBM) biomarker  
322 enzymes (GGT, ALP, Maltase, LAP) in the homogenates of both cortex and medulla of the kidney of rats  
323 following the administration of KBrO<sub>3</sub> could be due to KBrO<sub>3</sub> damaging effect on the structure and function  
324 of the BBM. Two likely reasons could be responsible for these suggestions:

- 325 (1) There could be direct loss of BBM or enzyme molecules into the lumen of the tubule following the  
326 toxic attacks by KBrO<sub>3</sub>, as it was reported for some nephrotoxicants by other researchers [28, 29]  
327 and  
328 (2) There could be enzymes inactivation due to conformational changes in the molecular structure of  
329 the BBM enzymes. Earlier researchers have reported oxidative modification of amino acid side  
330 chains of enzyme protein by reactive oxygen species that was generated by KBrO<sub>3</sub> toxicity and  
331 consequent inactivation of the BBM enzymes [30, 31, 32]

332 BBM forms the major lining of the epithelial cells of the proximal tubule of the kidney and it was  
333 reported to be the first barrier for various solutes during absorption in the kidney. Other workers have  
334 reported the BBM as major target of renal injury due to ischemia and nephrotoxic agents [4, 28, 33]. This  
335 of course could explain either of the two suggestions given above on why there was a significant (P<0.05)  
336 decline in the activities of all the BBM marker enzymes observed in this research. However, concurrent  
337 administration of KBrO<sub>3</sub> and methanol extract of *Carica papaya* seed was able to attenuate the decreases  
338 in activities of all the BBM marker enzymes studied in a dose dependent fashion and restored the values  
339 towards normal control.

340  
341 The significant decreases (P<0.05) in the activities of antioxidant enzymes studied (CAT, SOD, GPx) in  
342 homogenates of both cortex and medulla of the kidney of rats following administration of KBrO<sub>3</sub> could be  
343 due to the KBrO<sub>3</sub>-induced reactive oxygen species production that could have caused oxidative stress.  
344 ROS and oxidative stress have been reported to be important mediators of KBrO<sub>3</sub> nephropathy [34].  
345 Similarly, higher levels of ROS increases oxidative modification of cellular components in protein, lipids  
346 and nucleic acids, and causes damage to organelles such as the plasma membranes, mitochondria in the  
347 kidney and other tissues [35, 36].

348  
349 The significant decrease (P<0.05) in GSH level in renal cortex and medulla of rats following  
350 administration of KBrO<sub>3</sub> could be due to oxidative stress occasioned by ROS. GSH, a valuable  
351 physiological tri-peptide is reported to be a vital extracellular and intracellular protective antioxidant  
352 against oxidative stress. It reduces H<sub>2</sub>O<sub>2</sub> and hydroperoxides by its redox and detoxification reactions and  
353 protects protein thiols groups from oxidation [19]. Literature reported that decrease in the level of GSH will  
354 increase OS which will subsequently lead to cell damage [37].

355  
356 Free radical and ROS is key to the initiating of peroxidation of polyunsaturated fatty acid (PUFA). This  
357 reaction cascade which could lead to formation of lipid peroxide can be prevented by enzymatic and non  
358 enzymatic controlled antioxidant [19]. In the present study, the significant increase (P<0.05) in MDA

359 concentration of kidney tissues in KBrO<sub>3</sub> administered rats could be the consequence of an increment in  
360 the formation of oxygen free radicals generated by KBrO<sub>3</sub> and in fact indicates a compromised antioxidant  
361 defense system. MDA is one of the potent metabolites of peroxidized PUFA [19, 38] and is considered a  
362 biomarker of oxidative stress [39, 40]. Presence of MDA can induce cellular and functional damage [41].  
363

364 However, concurrent administration of KBrO<sub>3</sub> and methanol extract of *Carica papaya* seed prevented all  
365 the changes in the parameters of oxidative stress studied in both cortex and medulla in a dose dependent  
366 manner  
367

#### 368 4. CONCLUSION

369 Methanol extract of *Carica papaya* seed possess ameliorative effect against potassium bromate- induced  
370 nephrotoxicity and the most effective dose was 600mg/kg body weight. The effect of the extract was  
371 experienced in both the renal cortex and medulla. Since KBrO<sub>3</sub> is known to induces oxidative stress in the  
372 cell which in part is considered responsible for its toxicity, the preventive role of *Carica papaya* seed  
373 extract could be due to its ability to act as an antioxidant and a scavenger of reactive oxygen species.  
374 Thus *Carica papaya* seed extracts can be considered a potential preventive agent against renal damage  
375 caused by KBrO<sub>3</sub> and other structurally related compounds.  
376

#### 377 COMPETING INTERESTS

378 Authors have declared that no competing interests exist.  
379

#### 380 REFERENCES

- 381 1. Starek A, Beata S. Toxicological properties of potassium bromate. *Journal of pharmacological*  
382 *reports*. 2016; 1(3): 2-10.
- 383 2. Kolisetty N, Delker DA, Muralidhara S, Bull RJ, Cutruvo JA and Fisher JW. Changes in mRNA  
384 and protein expression in the renal cortex of male and female F344 rats treated with bromate.  
385 *Arch Toxicol*. 2013; 87:1911- 1925.
- 386 3. International Agency for Research on Cancer (IARC). Monographs on evaluation of the  
387 carcinogenic risk of chemicals to humans in some naturally occurring and synthetic food  
388 components, furoconmarins and ultra violet radiation. Lyon France. 1986; 40, Pp. 207 – 220.
- 389 4. Ahmad MK, Khan AA, Ali NS and Mahmood R. Chemo-protective effects of taurine on potassium  
390 bromate-induced DNA damage, DNA protein cross linking and oxidative stress in rat  
391 intestine. *Amino acids*. 2015; 45: 1109 – 1121.
- 392 5. Ahmad MK. and Mahmood R. Alterations in brush border membrane enzymes, carbohydrate  
393 metabolism and oxidative damage to rat intestine by rat intestine by potassium bromate.  
394 *Biochemie*. 2012; 94: 2776 – 2782.
- 395 6. Mark JE. Cataractogenic potential of bromate mediated oxidative stress in rats. *Animal Biol*.  
396 1988; 45: 567 – 660.
- 397 7. Anjum V, Ansari SH, Naquvi KJ, Arora P and Ahmad A. Development of quality standards of  
398 carica papaya. Linn leaves *Sch. Res. Lib*. 2013; 5 (2): 370 – 376.
- 399 8. Gurditta P. Baskeful benefits of papaya. *Int. Res. J. Pharm*. 2011; 2 (7): 6 – 12.
- 400 9. Arvind GD, Duravel S and Hurish G. Traditional medicinal uses of carica papaya. *J. Med Care*  
401 *Pap*. 2013; 1 (1): 2320 – 3862.
- 402 10. Yogiraj V, Kumar P, Chauhan C, Goyal A and Vyas B. Carica papaya Linn: An overview. *Int. J. of*  
403 *Herbal Medicine*. 2014; 2 (5): 01-08.
- 404 11. He X, Yi G, Wu L, Zhou HG. Chemical composition and antifungal activity of *Carica papaya* Linn  
405 seed essential oil against *Candida spp*. *Letters in applied microbiology*. 2017; 64 (5): 124-132.
- 406 12. Azwanida NN. A review on the extraction methods used in medicinal plants, principle, strength  
407 and limitation. *Med. Aromat. Plants*. 2015; 10: 417-422.
- 408 13. Singha AK. Colorimetric assay of catalase. *Analytical Biochemistry*. 1972; 47: 389-394
- 409 14. Misra HP, Fridovich I. The role of superoxide anion in the auto oxidation of epinephrine and a  
410 simple assay for superoxide dismutase. *J. Biol. Chem*. 1972; 247: (12) 3170-3175.  
411  
412  
413

Comment [Office9]: Comment: 1911 – 1925  
become 1911-25 for all references

- 414 15. Rotruck JT, Pope HE, Ganther AB, Swanson S. Biochemical role as component of glutathione  
415 peroxidase. *J science*. 1973; 179: 588-590.
- 416 16. Beutler E, Duron O, Kelly BM. Improved method for determination of glutathione. *J. Lab Clin Med*.  
417 1963; 61: 882-890
- 418 17. Varshne, R, Kale RK. Effects of calmodulin antagonists on radiation-induced lipid peroxidation in  
419 microsomes. *Int J. Rad Biol*. 1990; 58: 733-743.
- 420 18. Krause WJ. The art of examining and interpreting histologic preparations. A student  
421 handbook. Partheton publishing group London, UK. 2001 ; 9-10.
- 422 19. Noori S. An overview of oxidative stress and antioxidant defensive system. *Open Access Scientific*  
423 *Reports*. 2012; 1(8):1-9.
- 424 20. Ighodaro OM, Akinloye OA. First line defense antioxidants-superoxide dismutase (SOD),  
425 Catalase (CAT) and glutathione peroxidase (GPx): their fundamental role in the entire antioxidant  
426 defense grid. *Alexandria Journal of Medicine*, 2017; 12: 213-230.
- 427 21. Olagunju J, Adeneye A, Fagbohunka B, Bisuga N, Ketiku A and Benebo A. Nephroprotective  
428 activities of the aqueous seed extract of *Carica papaya* linn in carbon tetrachloride – induced  
429 renal injured Wistar rats: A dose and time-dependent study. *Biology and Medicine*. 2009 ; 1(1):  
430 11-19.
- 431 22. Nale L, More P, Ghumare B, Shendre S and Mote C. Protective effects of *Carica papaya* L. seed  
432 extract in gentamicin induced hepatotoxicity and nephrotoxicity in rats. *Int. J. Bio. Sci*. 2012; 3 (3):  
433 508 – 515.
- 434 23. Lakshmi MS, Reddy UK, Rani SK. A Review of Medicinal Plants for Nephroprotective  
435 Activity. *Asian Journal of Pharmaceutical and Clinical Research*. 2012; 5 (4): 8-14.
- 436 24. Naggayi M, Makubi N, and Iliya E. The protective effects of aqueous extract of *Carica papaya*  
437 seeds in paracetamol induced nephrotoxicity in male Wistar rats. *African Health Sciences*. 2015;  
438 15 (2): 599 – 605.
- 439 25. Montenegro J, Gonzalez O, Saracho R, Aquirre R and Marinez I. Changes in renal function in  
440 primary hypothyroidism. *American Journal of Kidney Disease*. 1996; 27 (2): 195 – 198.
- 441 26. Sarika A, Ranjina C., Devika T and Mallika V. Biochemical markers of liver and kidney function  
442 are influenced by thyroid function. A case controlled follow up study in Indian hypothyroid  
443 subjects. *Indian Journal of Clinical Biochemistry*. 2009; 24 (4): 370 – 374.
- 444 27. John AD, Larry AW. The autonomic nervous system and renal physiology. *International Journal of*  
445 *Nephrology and Renovascular disease*, 2013; 6: 149-160.
- 446 28. Fatima S, Yusifi AN, Mahmood R. Effect of cisplatin on renal brush border membrane enzymes  
447 and phosphate transport. *Hum Exp Toxicol* 2004; 23:547 – 554.
- 448 29. Scherberich JE, Wolf G and Schoeppe W. Shedding and repair of renal cell membranes following  
449 drug induced nephrotoxicity in humans. *Eur. J. Clin. Pharmacol*. 1993; 44:33-38.
- 450 30. Dudeja PK and Brasitus TA. Inactivation of rat small intestinal brush border membrane alkaline  
451 phosphatase by oxygen free radicals. *Gastroenterology* 1993; 105:357 – 366.
- 452 31. Gupta A, Sharma S, Nain CK, Sharma BK. and Ganguly NK. Reactive oxygen species-mediated  
453 injury in experimental ascending pyelonephritis. *Kidney Int*. 1996; 49:26-33.
- 454 32. Nath KA. and Norby SM. Reactive oxygen species and acute renal failure. *American Journal of*  
455 *Medical Sciences*. 2000; 109: 665 – 678.
- 456 33. Goldman M, Yaari A, Doshnitzki Z, Cohen – Luria R, Movan A. Nephrotoxicity of uranyl acetate:  
457 Effect on rat kidney brush border membrane vesicles. *Arch Toxicol*. 2006; 80: 387 – 393.
- 458 34. Zhang Y, Jiang L, Geng C, Li L, Shao J and Zhong L. Possible involvement of oxidative stress in  
459 potassium bromate induced gene toxicity in human HepG2 cells. *ChemBiol Interact*. 2011;  
460 189:186 – 191.
- 461 35. Giri U, Iqbal M and Athar M. Potassium bromate induces renal proliferative response and damage  
462 by elaborating oxidative stress. *Cancer let*. 1999; 135: 181 – 188.
- 463 36. Oloyode OB and Sunmonu TO. Potassium bromate content of selected bread samples in Ilorin,  
464 Central Nigeria and its effect on some enzymes of rat liver and kidney. *Food Chemistry*  
465 *Toxicology* 2009; 47: 2067 – 2070.
- 466 37. Khan N, Sharma S. and Sultana S. *Nigella sativa* (blac cumin) ameliorate potassium bromate  
467 induced early events of carcinogenesis: diminution of oxidative stress. *Human Exp. Toxicol*. 2003;  
468 23: 193 – 203.

- 469 38. Ohkawa H, Ohishi N and Yogi K. Assay for lipid peroxidation in animal tissues by thiobarbituric  
470 acid reaction. *Analytical Biochem.* 1979; 95: 351 – 358.
- 471 39. Kim YH, Mun KC, Lee SS, Seo SH, Kwak CS, Park SB and Kim HC. Oxidative damage in renal  
472 transplant patients. *Transplant Proc* 2000; 32:1777-8.
- 473 40. Dotan Y, Lichtenberg D and Pinchuk I. Lipid peroxidation cannot be used as a universal criterion  
474 of oxidative stress. *Prog Lipid Res.* 2000; 43:200-27
- 475 41. Cheeseman KH. Mechanisms and effects of lipid peroxidation. *Mol Aspect Med.* 1993; 14:191-7.

UNDER PEER REVIEW