

1                   **COMPARATIVE MORPHOMETRY OF THE GENUS THAIS FROM NEMBE,**  
2   **BAKANA AND CALABAR**

3  
4   **ABSTRACT**

5   AimMorphometry of *Thais* spp found in the Niger Delta Mangrove vegetation of Bakana, Calabar and Nembe were  
6   examined and compared. A total of 600 specimens (100 specimens per month) were collected during a period of six  
7   months (January to June 2018) from the three sampling communities. Three different species were identified  
8   namely: *Thais coronata*, *Thais haemastoma* and *Thais lacera*. Shell dimensions were measured to the nearest  
9   millimeter using Vernier calipers and weighed, to get the morphometric Characteristics: Whorl Diameter (WD),  
10   Shell length (SL), Shell width (SW), Body whorl length (BWL), Aperture length (AL), Aperture Width (AW) Shell  
11   Breadth (SB), and Animal Weight (AW). Number of whorls, number of primary spiral cord on the body and  
12   number of ridges or teeth inside of outer tip of the aperture were counted. The disparity between the morphometric  
13   traits across the different species identified were minimal as most of the species had similar values of  
14   morphometric traits. However, differences can be identified using their colour; *thais coronata* (dirty light grey), *T*  
15   *haemastoma* (light grey), and *T lacera* (plane grey). The Three (3) species had a modal length class of 3.5cm to  
16   4.5cm. *Thais coronata* and *Thais lacera* had a modal weight class of 9-11grams while *Thais haemastoma* had a  
17   modal weight class of 6-7grams. It was observed with the aid of length/weight relationship that the found in all  
18   study. It was observed with the aid of length/weight relationship that the *Thais* specimen found in all study  
19   locations exhibited a very weak linear relationship with very low  $R^2$  values across locations. The exponent b of  
20   *Thais coronata* and *Thais haemastoma* and *T. lacera* across the three study locations indicate a negative allometric  
21   growth pattern. The Month of April for samples collected from Nembe had the highest condition factor for the  
22   three (3) species. *T. coronata* (4.4), *T. lacera* (6.38) and *T haemastoma* (5.5).

23   **Keywords: comparative, morphometry, genus Thais, Nembe, Bakana, Calabar.**

24  
25  
26   **INTRODUCTION**

27   The genus *Thais* belong to the family *Muricidae*, and are gastropods that are found in  
28   the phylum Mollusca. They are one of the largest group of marine organisms and have  
29   been known for many years as a major source of protein consumed by human and  
30   other macro organisms. This class gastropods have been known to consist of snails  
31   that possess outer shells into which the animal can generally always withdraw.  
32   Gastropods were found and were also to known to successfully thrive and live in  
33   different habitats such as ocean, fresh water and land. They perform specific roles in  
34   keeping ecological balance intact and they, being a highly diversified group compared  
35   to the other group in the phylum Mollusca are commercially beneficially to humans.  
36   They are also used as ornaments and perform various ecological functions, especially  
37   maintaining the balance in the environment as well as to provide food and livelihood for  
38   humans This family contains a highly diverse group of species that are distributed in  
39   tropical, subtropical regions (Davis & Fitzgerald, 2004). In Nigeria, *thais* is found in  
40   Mangrove forests located in the Niger delta region. The Niger Delta mangrove forests  
41   forms a clear vegetation zone along the entire coastline and plays the traditional role of  
42   breeding and nursery ground of important fish and shell fish. The gastropod mollusks  
43   (*thais*, bivalves) are the permanent inhabitants of the mangrove community (Nazim *et*  
44   *al.*, 2015). The *Muricidae* are the third largest group in the class gastropoda and are a

45 taxonomically complex family consisting of around 1,502 species that are found  
46 worldwide (Bailly,2012). For classification and Nomenclatures of gastropod family, the  
47 family is separated into 13 sub- families that are further subdivided into more than 90  
48 genera. This classification is based largely on superficial shell and radular character  
49 due to poor phylogenetic knowledge associated with this family (Bieler, 1992).

50 Muricidae are members of the order neogastropod which contains more than 10,775  
51 estimated species and represent the largest order in the class Gastropoda and  
52 comprises close to 30,239 species (Radwin *et al.*, 1972; Bailly, 2012). Members of the  
53 Muricidae are distinguished from other neogastropods families by the presence of rows  
54 of protrusions or spines on their shells (Carpenter and Niem,1998). The shell sculpture  
55 is elongated possessing a long siphon canal, their operculum has either a marginal or  
56 lateral nucleus and their eggs are usually laid in protective corneous capsule that  
57 usually form when crawling juveniles hatch. Planktonic larva are carnivores that  
58 generally feeds on economically important mollusks as well as *barnacles* (Al-Yamani *et*  
59 *al.*, 2012). The soft body of their prey is reached by drilling hole with the aid of a  
60 softening secretion and scraping of a toothed structure known as radula. Their  
61 carnivorous tendencies make them to be considered as pests, as they may cause  
62 substantial destruction in exploited natural beds and areas of cultured commercial  
63 bivalves.

64 *Thais*, rock shell, dog whelk, dog winkles, ngolo *Thais*, rock shell, dog whelk, dog  
65 winkles, Ngolo. They are present on mangrove tree trunks, breathing roots, oyster  
66 beds, granite bunds, walls of intertidal monsoon drains, as well as on rocks and  
67 boulders on the shore and exhibit both restricted geographical and local distribution  
68 (Davis and Fitzgerald, 2004). They generally prey on barnacles, polychaetes, bivalves  
69 and other gastropods (e.g., Taylor, 1976, 1980). Some feed on the sap of a dead  
70 mangrove tree. Therefore, the aim of this study was to assess and compare the  
71 mophometry of the *Thais* species from Nembe, Bakana and Calabar.

72

73

## 74 **MATERIALS AND METHODS**

### 75 ***Study Area***

76 The study areas were in Bayelsa in Nembe, Rivers in Bakana, and cross river in their different  
77 fishing pond settlement namely mobogiri, golibogiri, and fisherman village in Nembe,  
78 Owuogono, ebekemoko in bakana. The vegetation's of the area is predominantly mangrove and  
79 swamps with no occurrence of *Nypa* palm and other coastal vegetation. The tidal amplitude is  
80 between 1.5 to 2m in normal tide and water level increases and decreases depending on the  
81 lunar cycle (Ogamba, 2003).


### 82 **Collection of Sample (*Thais* Sampling)**

83 The samples were collected by hand picking on the mangrove mud during low tide at the  
84 locations by the local fishermen and carried in sack bag until large enough before bringing it to  
85 the community where it's been brought by the traders and taken to the market for sales to the  
86 mzlarket women, in which the sample is brought and different species that are labeled  
87 separately differentiating the different species and taken to the laboratory where it is stored for  
88 analysis.

### 89 **Morphometric Measurements**

90 Shell dimensions will be measured to the nearest millimeter using Vernier calipers and weighed,  
91 to get the morphometric Characteristics: Whorl Diameter (WD), Shell length (SL), Shell width  
92 (SW), Body whorl length (BWL), Aperture length (AL), Aperture Width (AW) Shell Breadth  
93 (SB), and Animal Weight (AW).

94 Number of whorls, number of primary spiral cord in the body, number of ridges or teeth inside  
95 of outer tip of the aperture will be counted



96

97 Fig 3.1 Morphometric Parameters

### 98 **Laboratory Analysis**

99 Samples collected were washed properly to remove dirt, and were put in a sieve to drain and  
100 then stored in a polyethylene bag in the refrigerator for preservation.

101 (a) The number of whorls (NW) on each shell was counted and recorded.

- 102 (b) The shell length (SL) of each shell was measured with a Vernier caliper and  
 103 recorded in Centimeter (cm) to two places of decimal.
- 104 (c) The shell width (SW) of each shell was measured in centimeter (cm) in vernier  
 105 caliper and recorded.
- 106 (d) The aperture width (AW) also of all the specimen were measured and recorded in cm  
 107 nearest to two decimal places.
- 108 (e) The body whorl length (BWL) of each shell was measured in centimeter (cm) in  
 109 Vernier caliper and recorded.
- 110 (f) The shell thickness is measured of each shell is measured in centimeter to two  
 111 decimal places
- 112 (g) Number of whorl of each shell is counted and recorded (No of Whorl).
- 113 (h) Number of tubercles in the body whorl of each shell is counted.
- 114 (i) Number of ridges inside the upper lips is counted and recorded.
- 115 (j) The number of primary spiral cord of each shell is counted and recorded.
- 116 (k) Each shell with the contents (ws/m) was weighed in a Mettler Ae 163 balance and  
 117 recorded in grams nearest to two decimal places.
- 118 (l) Each shell was cracked to remove the fleshy body. The flesh was then put in a pre-  
 119 weighed watch glass and weighed. The weight of the flesh (weight of body mass –  
 120 wm) was obtained from weight of flesh + watch glass minus weight of watch glass.

121 The weights were recorded in grams to two places of decimals.

122 These readings (a – f) were all recorded in Appendix.

### 123 **Analysis of Data**

124 Shell dimensions will be measured to the nearest cm using vernier calipers and weighed, to get  
 125 the morphometric Characteristics shell length (SL), Shell width (SW), Body whorl  
 126 length(BWL), aperture length(AL), Aperture width(AW) shell weight(SW), and animal  
 127 weight(AW).

128 Number of whorls, number of primary spiral cord in the body, number of ridges or teeth inside  
 129 of outer tip of the aperture will be counted The animals would be relaxed in 7.5% magnesium  
 130 chloride solution mixed with an equal volume of seawater to examine soft body morphologies.  
 131 Juveniles and adults will be both examined, noting their colour (when dry or wet) and surface  
 132 morphology. The animals would be relaxed in 7.5% magnesium chloride solution mixed with  
 133 an equal volume of seawater to examine soft body morphologies.

### 134 **Length and Weight Relationship**

135 The length weight relationship was determined using cube law given by Lecren (1951).

$$136 \quad W = aL^b$$

137 Where W= Weight in grams (g)

138 L = Total Length in Centimeter (cm)

139 a = proportional constant or intercept

140 b = an Exponent

141 the equation was log transformed and were determined by linear regression analysis and scatter  
142 diagrams of length and weight were plotted

143 The logarithmic transformation of the formula is

144  $\text{Log } W = \text{Log } a + b \log L$

145 Where, W = weight of Thais in gram

146 L = observed total length in cm

147 a = regression intercept

148 b = the regression slope

149 The equation was log transformed to estimate the parameters “a” and “b”. If b is equal to 3, it is  
150 an isometric growth pattern, but if b is not equal to 3 (that is, b is > or < 3), it is an allometric  
151 growth pattern, which may be positive if b > 1 or negative if b < 1.

## 152 **Statistical Analysis.**

153 With the aid of JMP, SPSS and Microsoft Excel, statistical analysis was done on the data  
154 obtained from the study. Two sample student *t* test shall be used to compare the differences of  
155 the length and width of radula teeth, soft body shell ratios and other measured parameters. Chi-  
156 square tests shall be used to assess prevalence and intensity. L-W relationship shall be  
157 determined. Ratios of morph metric measurements against total Length were estimated.

158

159

160

## 161 **RESULTS AND DISCUSSION**

### 162 **Descriptive Analysis of Morphometric Traits**

163 During the sampling period there were different species of Thais found in the study locations of  
164 Nembe, Bakana and calabar. The species were picked randomly at the study sites. There was a  
165 combination of different species of thais namely *thais coronate*; which was the dominant  
166 species of at least five out of ten, followed by *thais Heamastoma* and then *thais lacera*.

167

#### 168 **Thais lacera**

169 Their shells have 2-5 body whorl with largest secondary spiral cord of (8-12) that are present  
170 between first two cells. its aperture is ovate and the inside of the outer lips is smooth while their  
171 siphonal canal is short and two groved sulcus present instead of outer lip colon. The shell  
172 surface colour is plane grayish or yellow tan.



173

174 **Plate 4.1: *Thais lacera***

175 *Thais Coronata*

176 Commonly known as the rock shell has a thick walled shell and mostly noticed to have  
 177 short wall with the shell closed by a long operculum, they are up to 5cm in length and their  
 178 colour is dirty grey to brown grey



179

180 **Plate 4.2 *Thais Coronata***

181 *Thais Heamostoma*

182 This conch shell is red mouthed up to 80cm long, is robust, oval and has a series of nodes that run  
 183 along the spiral shell and very short and the operculum is cod



184

185 **Plate 4.3 *Thais Heamostoma***

186 Table 4.1 shows the Comparative statement of Meristic Traits in three Species of Thais  
 187 identified during the study.

188 **Table 4.1 Comparative Statement of Meristic Traits in Three Species of Thais**

<b>Morphology</b>	<b><i>T. Coronata</i></b>	<b><i>T. haemastoma</i></b>	<b><i>T. lacera</i></b>
No of Whorl (Range)	2-6	3-6	2-5
Colour	Dirty grey	Light grey	Plane grey
No of Ridges (Range)	5-27	8-25	0
No of Spiral cord (Range)	7-51	17-43	22-51
No of Nodes on body wall (Range)	3-33	10-21	9-21
Shell Thickness	0.1	0.1	0.1
APL/AW (Ratio)	2.06	2.36	2.41
APL/BWL (Ratio)	0.79	0.41	0.83
BWL/WD (Ratio)	0.87	0.92	0.87
SL/BWL (Ratio)	1.16	1.16	1.13
SL/APL (Ratio)	1.45	2.87	1.32

189

190 **Comparative Meristics Trait**

191 Table 4.1 shows the result of the body ratio of the Aperture length to animal weight, Aperture  
 192 length to the body whorl length, body whorl length to the whorl diameter, shell length to the  
 193 body whorl length and shell length to the aperture length.

194 **Table 4.2: Descriptive Statistics of Morphometric traits**

<b>Statistics</b>	<b>Variable</b>	<b>T. coronate</b>	<b>T. haemastoma</b>	<b>T. lacera</b>
<b>Mean±SD</b>	SL	4.03±0.77	4.28±0.87	3.95±0.6
	SW	3.28±0.5	3.31±0.43	3.38±0.46
	BWL	3.45±0.53	3.68±0.47	3.48±0.62
	APL	2.77±0.41	2.98±0.44	2.89±0.45
	APW	1.34±0.34	1.39±0.26	1.33±0.25
	SWT	10.56±3.67	10.02±3.47	10.67±3.32
	AW	1.34±0.76	1.26±0.89	1.21±0.67
<b>Minimum`</b>	SL	2.3	3	2.4
	SW	2.1	2.3	2.4
	BWL	0.1	2.5	2
	APL	1.6	2	1.6

	APW	0.1	1	1
	SWT	3.85	5.39	5.45
	AW	0.2	0.4	0.4
Maximum	SL	9.3	9.3	4.8
	SW	4.5	4.3	4.5
	BWL	4.7	4.5	4.7
	APL	4.1	4.4	3.8
	APW	3	2.2	2
	SWT	25.88	23.34	20.79
	AW	6.5	6.5	4.5


Where SL (Shell Length), SW (Shell width), BWL (Body whorl length), APL (Aperture length), APW (Aperture width), SWT (Shell weight) and AW (Animal weight)

195 Descriptive analysis of Shell Morphometric traits

196 Table 4.2 Shows the mean standard deviation of the shell length, shell width, Body whorl  
 197 length, Aperture length, Aperture width, shell weight and animal weight of *T. coronata*, *T.*  
 198 *haemastoma* and *T. lacera* .

### 199 Length Size Class Frequency


200 *Thais coronata* found in all the study locations where measured to get Length size class (fig  
 201 4.1). Results show the most dominant size class or modal class to be 3.5cm-4cm (178) and 4cm-  
 202 4.5cm (123). Very few had size classes of 2cm-2.5cm (3) and 7cm-7.5cm (3). *Thais*  
 203 *Haemastoma* found in all the study locations where measured to get Length size class (fig 4.2).  
 204 Results show the most dominant size class to be 3.5cm-4cm (31) and 4cm-4.5cm (28). Very few  
 205 had size classes of 2cm-2.5cm (1) and 5cm-5.5cm (3cm). *Thais lacera* found in all the study  
 206 locations where measured to get Length size class (fig 4.3). Results show the most dominant  
 207 size class to be 4cm-4.5cm (31) and 3.5cm-4cm (23). Very few had size classes of 2cm-2.5cm  
 208 (4).



209

210 Fig 4.1 Length Size Class of *Thais Coronata* found in all the study Stations






211

212 Fig 4.2 Length Size Class of *Thais haemastoma* found in all the study Stations


213



214


215 Fig 4.3 Length Size Class of *Thais lacera* found in all the study Stations.

216 *Thais sp.* found in Calabar study location where measured to get Length size classes (fig 4.4).  
 217 Results show *Thais coronata* as dominant across most of the different size classes, followed by  
 218 *Thais haemastoma* and then *Thais lacera*. *Thais sp.* found in Nembe study location where  
 219 measured to get Length size classes (fig 4.5). Results show *Thais coronata* as dominant across  
 220 most of the different size classes, followed by *Thais haemastoma* and then *Thais lacera*. *Thais*  
 221 *sp.* found in Bakana study location where measured to get Length size classes (fig 4.6). Results  
 222 show *Thais coronata* as dominant across most of the different size classes, followed by *Thais*  
 223 *lacera*.



224


225 Fig 4.4 Length Size Class of species found in Calabar



226

227 Fig 4.5 Length Size Class of species found in Nembe

228



229

230 Fig 4.6 Length Size Class of species found in Bakana.


231 **Weight Class Frequency**

232 *Thais coronata* found in all the study locations where measured to get Weight class (fig 4.7).  
233 Results show the most dominant size class or modal class to be 10-11grams (68) followed by 9-  
234 10grams (55). Very few had weight classes of 3-4grams (1) and 21-22grams (1). *Thais*  
235 *Haemastoma* found in all the study locations where measured to get Weight class (fig 4.8).  
236 Results show the most dominant weight class to be 6-7grams (17) and 5-6grams (14). Very few  
237 had weight classes of 13-14grams (1) and 15-16grams (1). *Thais lacera* found in all the study  
238 locations where measured to get weight class (fig 4.9). Results show the most dominant size  
239 class to be 9-10grams (14) followed by 10-11grams (11). Very few had weight classes of 19-  
240 10grams (1) and 20-21grams (1). The results also show a random distribution of weight classes  
241 across species.

242


243

244



245

246 Fig 4.7 Weight Class of *Thais coronata* found in all the study Stations




247

248 Fig 4.8 Weight Class of *Thais haemastoma* found in all the study Stations

249

250

251



252

253 Fig 4.9 Weight Class of *Thais lacera* found in all the study Stations.

254 *Thais* sp. found in Calabar study location where measured to get Weight classes (fig 4.10).

255 Results show *Thais coronata* as dominant across most of the different size classes, followed by

256 *Thais haemastoma* and then *Thais lacera*. *Thais* sp. found in Nembe study location where


257 measured to get Length size classes (fig 4.11). Results show *Thais coronata* as dominant across

258 most of the different size classes, followed by *Thais haemastoma* and then *Thais lacera*. *Thais*

259 sp. found in Bakana study location where measured to get Length size classes (fig 4.12). Results

260 show *Thais coronata* as dominant across most of the different size classes, followed by *Thais*


261 *lacera*.



262

263 Fig 4.10 Weight Class of species found in Calabar


264



265

266 Fig 4.11 Weight Class of species found in Nembe

267



268

269 Fig 4.12 Weight Class of species found in Bakana

## 270 **Length and Weight Relationship**

271 The length and weight relationship of the different species across the different locations studied  
272 were analyzed. The result shows a very weak relationship (RSquare = 0.06) between the weight  
273 and the length of *Thais coronata* in Bakana. There is a negative allometric growth ( $a=0.67$ )  
274 between the weight and the length of *Thais coronata* in Bakana. The result shows a very weak  
275 relationship (RSquare = 0.037) between the weight and the length of *Thais coronata* in Calabar.  
276 It also shows a negative allometric growth ( $a=0.84$ ) between the weight and the length of *Thais*  
277 *coronata* in Calabar. The result shows a very weak relationship (RSquare = 0.057) between the  
278 weight and the length of *Thais coronata* in Nembe. It also shows a negative allometric growth  
279 ( $a=0.62$ ) between the weight and the length of *Thais coronata* in Nembe.

280 The result shows a very weak relationship (RSquare = 0.005) between the weight and the length  
281 of *Thais haemastoma* in Calabar. It also shows a negative allometric growth ( $a=0.87$ ) between  
282 the weight and the length of *Thais haemastoma* in Calabar. A very weak relationship (RSquare  
283 = 0.062) between the weight and the length of *Thais haemastoma* in Nembe. It also shows a  
284 negative allometric growth ( $a=0.64$ ) between the weight and the length of *Thais haemastoma* in  
285 Nembe.

286 There was a very weak relationship (RSquare = 0.023) between the weight and the length of  
287 *Thais Lacera* in Bakana. It also shows a negative allometric growth ( $a=0.68$ ) between the  
288 weight and the length of *Thais Lacera* in Bakana. A very weak relationship (RSquare = 0.02)  
289 between the weight and the length of *Thais Lacera* in Calabar was also observed. It also shows  
290 a positive allometric growth ( $a=1.05$ ) between the weight and the length of *Thais Lacera* in  
291 Calabar. The result shows a very weak relationship (RSquare = 0.09) between the weight and  
292 the length of *Thais Lacera* in Nembe. It also shows a negative allometric growth ( $a=0.57$ )  
293 between the weight and the length of *Thais Lacera* in Nembe.

294 A weak relationship (RSquare = 0.03) between the weight and the length of *Thais Coronata* in  
295 all locations was observed. It also shows a negative allometric growth ( $a=0.79$ ) between the  
296 weight and the length of all *Thais Coronata*. The result shows a weak relationship (RSquare =  
297 0.0298) between the weight and the length of *Thais haemastoma* in all locations. It also shows a  
298 negative allometric growth ( $a=0.75$ ) between the weight and the length of all *Thais*  
299 *haemastoma*. The result shows a very weak relationship (RSquare = 0.013) between the weight  
300 and the length of *Thais lacera* in all locations. It also shows a negative allometric growth  
301 ( $a=0.899$ ) between the weight and the length of all *Thais lacera*.

302

304 **Morphometric Traits**

305 *T. haemastoma* had an average shell length of 4.28cm and Shell width of 3.31cm, which varied  
306 minimally from *T. lacera* and *T. coronata* with shell lengths of 3.95cm and 4.03cm and shell  
307 widths of 3.38cm and 3.28cm respectively.

308 The disparity between the morphometric traits across the different species identified were  
309 minimal as most of the species had similar values of morphometric traits. Differences can be  
310 identified using their colour; *thais coronata* (dirty light grey), *T haemastoma* (light grey), and *T*  
311 *lacera* (plane grey) and the number of ridges *T. coronata* (5-27), *T. haemastoma* (8-28) and *T.*  
312 *lacera* lacking ridges

313 *T. lacera* has the ratio of aperture length (APL) to the body whorl length (BWL) of 0.83 and  
314 also aperture length to animal weight as 2.41 showing that the aperture length in *T. lacera* is  
315 quite large compare to the other species. *T. haemastoma* has the highest ratio of body whorl  
316 length to the body width of 0.92 and ratio of shell length (SL) to the aperture length (AL) is  
317 2.83.

318 This agrees with (Trussell & Etter, 2001) in their review of gastropods suggested that variations  
319 in morphometric traits become obvious as you proceed deeper from the brackish into the oceans  
320 as wave exposure has a direct relationship with length of the shell.

321 **Length and Weight Size Class**

322 The frequency distribution of shell length and Shell weight of the collected *Thais* snails from  
323 the results shows the estimated modal class in the frequency distributions estimated from the  
324 three sampling area and the combined data. Thus, the analysis of the modal Length size  
325 classes, modal weight size classes and interpretations are based on the combined population  
326 sampled across the months of study.

327 Most of them had a normal distribution. *Thais haemastoma* was absent from the Bakana  
328 study Station but had a modal length size class to be 3.5cm-4cm (31) and 4cm-4.5cm (28) in  
329 the two other locations. Very few had size classes of 2cm-2.5cm (1) and 5cm-5.5cm (3cm).  
330 *Thais lacera* found in all the study locations showed a dominant size class of 4cm-4.5cm (31)  
331 and 3.5cm-4cm (23) while, *Thais coronata* found in all the study locations had a dominant  
332 size class of 3.5cm-4cm (178) and 4cm-4.5cm (123).

333 In Calabar study station, most of the species had a Length size class ranging from 3.5cm to  
334 5cm with *Thais coronata* most dominant across most of the different size classes, followed  
335 by *Thais haemastoma* and then *Thais lacera*. In Nembe study station, most of the species had  
336 a modal size class ranging from 3.5cm to 5cm with *Thais coronata* most dominant across  
337 most of the different size classes, followed by *Thais haemastoma* and then *Thais lacera*. In  
338 Bakana study station, most of the species had a modal size class ranging from 3cm to 5cm  
339 with *Thais coronata* also the most dominant. In all the surveys of the population structure it  
340 was clear that small individuals (<10mm) were generally absent from most of the  
341 populations. And in the above size class it was observed that the population of the smaller  
342 size class 2.0cm -2.5cm of age one and below is very few compare to the size class of 3.5cm-  
343 4cm and this is as a result of the fact that the samples are market derived and the fishermen  
344 allows the smaller sizes to stay to up to a reasonable size before picking them. while the size  
345 class of 4.5 to 5.0 is not seen in the frequency table compared to FAO standard of *thais*  
346 *coronata* matured size as 0f 5cm and *T. haemastoma* standard mature size of 5cm to 6cm and  
347 this shows that the *thais* species is an endangered species due to the fact that they are not  
348 allowed to get to full maturity and there were no presents of eggs in any of the organism.

349 The length frequency also showed that *T coronata* as the dominant species found in the three  
350 locations with a size class of 3.5cm-4.0cm.



351 *Thais coronata* found in all the study locations had a modal weight class of 10-11grams (68)  
352 followed by 9-10grams (55). Very few had weight classes of 3-4grams (1) and 21-22grams  
353 (1). *Thais Haemastoma* found in all the study locations had a modal weight class of 6-7grams  
354 (17) and 5-6grams (14). Very few had weight classes of 13-14grams (1) and 15-16grams (1).

355 *Thais lacera* found in all the study locations also showed a most dominant size class of 9-  
356 10grams (14) followed by 10-11grams (11). Very few had weight classes of 19-10grams (1)  
357 and 20-21grams (1)

358

### 359 **Length-Weight Relationship**

360 The length–weight studies are made to determine mathematically the relationship  
361 between two variables and enable prediction of the other variable when one variable is  
362 known. As the animal grows it is said that the resultant increase in size, shape, and  
363 volume can be measured as length and weight relationship which has become a  
364 standard practice in fishery

365 It was observed with the aid of length/weight relationship that the found in all study  
366 locations exhibited a very weak linear relationship with very low  $r^2$  value across  
367 locations. According to Tesh, “If  $b$  values equals 3, it shows that the organism has a  
368 symmetric or isometric growth pattern while values of  $b$  which are more than or less  
369 than 3 shows that the fish growth pattern is allometric” (Tesh, 1971).

370 The exponent  $b$  of *Thais coronata* and *Thais haemastoma* and *T lacera* across the three  
371 study locations indicate a negative allometric growth pattern there by not showing any  
372 variance from  $b$  which is 3 which has been shown that the increase in weight of the  
373 animal is not proportionate to the cube of its length and that they maintain specific  
374 body shape throughout their life (Archya 1980)

375 This can be attributed to the nature of their habitat and their influence of their  
376 environment, condition of the growth and shell properties (Wilson & Owen 1969),  
377 Saad 1997, Gaur et al 2006. This also agrees with the study of Laximilathal, (2008)  
378 and Kesavan, (2012) who postulatded that in the temperate regions the growth line of  
379 the shell mollusk is said to be a pointer of age whereas at the tropical region due to the

380 lack of distinct season and limited variation of environmental parameters much  
381 difference in growth line is not visible.

382

## CONCLUSION

383 *Thais*, rock shell, dog whelk, dog winkles, ngolo *Thais*, rock shell, dog whelk, dog winkles,  
384 Ngolo. They are present on mangrove tree trunks, breathing roots, oyster beds, granite bunds,  
385 walls of intertidal monsoon drains, as well as on rocks and boulders on the shore and exhibit  
386 both restricted geographical and local distribution. Based on shell morphology alone, it is  
387 difficult to differentiate the species belonging to genus *Thais* because of large amount of  
388 plasticity, observed in the shell characters. The colour of the shells are poorly defined as  
389 species identifying character in *Thais* species. Due to lack of taxonomic clarity of the species  
390 in the Niger Delta region there is the need to know the different types of the species to help  
391 scientific studies currently ongoing in microbiology, biodiversity and parasitology.

392 The Study has shown that we have three different species of *Thais* found in the study  
393 locations of Nembe, Bakana and calabar. The species were picked randomly at the study sites  
394 and are namely *thais coronate*, *thais Heamastoma* and then *thais lacera*. Their sexual  
395 dimorphism and related characteristics, length weight relationship in the study showed that  
396 there is no relationship and that the species can be short and rounded but still have weight; it  
397 has a negative allometry that shows or indicate a decrease condition or elongation

398 The modal length class, that shows three modal age, of 0-1, 1year and two years and above  
399 with *T. coronate* the dominant species with a highest modal class.

400

## REFERENCES

- 401 1. Davis, R.A. & Fitzgenald, D.M. (2004). *Beaches and Coasts*. Blackwell, U.K., p. 419
- 402
- 403 2. Nazim, K., Moinuddin, A., Muhammad, U. Khan<sup>3</sup>, S. S. Shaukat,<sup>2</sup> A.K. & Agha,  
404 T.H.D. (2015). Population distribution of mollusks in mangrove forests, Pakistan.  
405 *FUUAST Journal of Biology*, 5(1), 37-41.
- 406 3. Bailly, N. (2012). Catalog of life 2012 annual checklist.
- 407
- 408 4. Bieler, R.. (1992). Gastropodee phylogeny and systematic. Rudiger Bieler *Annual*  
409 *Review of Ecology and Systematics*, 23, 311-338
- 410 5. Radwin, G.E., D'Attilio, A., & Mulliner, D.K. (1976). *Murex shells of the world: an*  
411 *illustrated guide to the Muricidae*, Stanford University Press, Stanford, California.

- 412 6. Al-Yamani, F. Y., Skryabin, V., Boltachova, N., Revkov, N., Makarov, M., Grinstov,  
413 V., & Kolesnikova, E. Illustrated Atlas on the Zoobenthos of Kuwait. Kuwait Institute  
414 for Scientific Research. (2012).  
415
- 416 7. Carpenter, K.E.; Niem, V.H. (eds) FAO species identification guide for fishery  
417 purposes. The living marine resources of the Western Central Pacific. Volume 1.  
418 Seaweeds, corals, bivalves and gastropods. Rome: FAO, (1998).  
419
- 420 8. (Davis, R. and Fitzgerald, D.. Evaluating the growth and age of the netted whelk  
421 *Nassarius reticulatus* (gastropoda: nassaridae) from statolith growth rings. *Marine*  
422 *Ecology Progress Series*, 2004; 342, 163-176.
- 423 9. Ogamba, E.N. Water quality status of Elechi Creek complex in relation to  
424 physicochemical parameters and plankton distributiojn; (2003).
- 425 10. Trussell, G. C., & Etter, R. J. Integrating genetic and environmental forces that shape  
426 the evolution of geographic variation in a marine snail, In *Microevolution Rate,*  
427 *Pattern, Process.* (pp. 321-337) Netherlands: Springer; 2001.
- 428 11. Tesh, G. Speciation and diversity on tropical rocky shores: a global phylogeny of  
429 snails of the genus *Echinolittorina*. *Evolution*. 1971; 58, 2227-2251.
- 430 12. Archya, E. Use of internal growth bands for measuring individual and population  
431 growth rates in *Mytilus edulis* from offshore production platforms. *Marine. Ecology*,  
432 1980; 66, 259-265.
- 433 13. Wilson, R.J.P. & Owen. Functional morphology, ecology and evolutionary  
434 conservatism in the Glycymerididae (Bivalvia). *Palaeo*,(1969; 18(2), 217–258.
- 435 14. Kesavan, K. (2012). Molluscan Diversity in Mangrove Ecosystem of Uran (Raigad),  
436 Navi Mumbai, Maharashtra, West coast of India. *Bulletin of. Environmental and.*  
437 *Pharmacological Life Science*. 1(6), 55-59.
- 438 15. Laxmilathal, D.G. (2008). *The littorinid mollusks of mangrove forests in the Indo-*  
439 *Pacific region. The genus Littoraria. British Museum, London.* pp. 227.

440

441