

## **Original Research Article**

Toxicological, Biochemical and Haematological Studies of Ethyl acetate Extract of *Persea americana* Leaf in Albino Rats

### **ABSTRACT**

The aim of this present study was to evaluate the effect of ethyl acetate extract of *P. americana* on liver and kidney function, lipid profile as well as haematological parameters in albino rats. A total of 20 albino rats were used for this experiment and they were divided into four groups of 5 (A-D) rats each. Group A served as normal control, group B-D served as experimental groups administered with different doses of ethyl acetate *Persea americana* leaf extract. This study was conducted in the Department of Biochemistry, Bayero University, Kano, in the month of May, 2018. The animals were administered with 100, 200 and 400mg/kg per day of the extract for 4 weeks. The activities of serum transaminases, alkaline phosphatase, total protein, albumin, globulin, electrolytes, urea and creatinine were determined. Biochemical parameters were evaluated by measuring the concentration of total cholesterol, triglycerides, low density lipoprotein, very low density lipoprotein and high density lipoprotein. Haemopoietic effect was evaluated by measuring the levels of packed cell volume, red blood cells, white blood cells, haemoglobin, mean corpuscular volume, mean corpuscular haemoglobin, mean corpuscular haemoglobin concentration and platelets. The levels of serum ALT and albumin were significantly ( $P < 0.05$ ) increased whereas total protein, globulin and potassium levels were significantly ( $P < 0.05$ ) decreased. Significant ( $P < 0.05$ ) decrease in total cholesterol, triglyceride, low density lipoprotein and very low density lipoprotein was observed. Haematological parameters were not significant affected. In conclusion, ethyl acetate extract of *P. americana* may be relatively safe and possesses hypolipidemic potentials but extremely high doses may not be advisable.

Keywords: *P. americana*, liver function test, kidney function test, lipid profile, haematological parameters

### **1. INTRODUCTION**

Many medicinal plants used in ethnomedical practices in Nigeria are known or little known to scientific world [1]. Many important drugs used in medicine today are directly or indirectly derived from plants [2].

Plants and herbs have been a tremendous source of food and folk remedies for mankind and have served as starting material for the development of new synthetic drugs. They have the ability to synthesize a wide variety of chemical compounds which are used to perform biological functions and to defend against attack from predators such as insects, fungi, yeasts, bacteria, virus and other pathogens [3]. Chemical compounds in plants mediate their effect on the human body through processes identical to those already well understood for the chemical compounds in conventional drugs [4, 5]. *Persea americana* belongs to the family *Lauraceae*, unflatteringly known in the past as alligator pear, midshipman's butter and vegetable butter, it is one of those plants currently used by indigenous persons for its nutritional value and to manage health problems. It is well known in many parts of the tropical world including Nigeria [6]. The fruit tree can attain a height up to 20m, with large spreading and flat topped crown. The plant is reported to possess antidiabetic, antihyperlipidemic potentials [2, 7], antioxidant [8, 9], cancer risk reduction [10], wound healing [11], hepatoprotection [12], analgesic and anti-inflammatory [13], anticonvulsant [14], vasorelaxant and blood pressure reducing [15, 16]. The use of traditional medicines as substitutes to orthodox medicines has been on the increase [17]. The reasons, which have given rise to this trend, include the cheapness, availability and accessibility of these natural medicines. Besides, there has been the erroneous belief that these medicines are free from adverse effects [18, 19]. On the other hand they have been rejected because many of the acclaimed medicinal values have not been scientifically evaluated and their safety profiles uncertain [19]. It is, therefore, pertinent that safety assessments should be conducted on natural products for which certain medicinal uses have been scientifically validated. The World Health Organization has also recommended the evaluations of traditional medicine, because of the modern drugs are not safe [20]. It is, therefore, pertinent that safety assessments should be conducted on ethyl acetate extract of *P. americana* for which certain medicinal uses have been scientifically validated.

## 2. MATERIALS AND METHODS

## 2.1 Sample Collection and Identification

The leaf of *Persea americana* were collected from Jos, Plateau state of Nigeria. The plant part was authenticated by a Botanist at Plant Science Department, B.U.K. with accession number BUKHAN 0305 and was deposited in Bayero University, Kano herbarium for reference.

## 2.2 Extraction of Plant Material

The sample was chopped into small pieces and then shades dried and ground into powdered form. Soxhlet extraction method was used for the extraction of the plant materials. Ethyl acetate was used as the extraction solvent and later concentrated in vacuum using rotary evaporator at 40°C.

## 2.3 Experimental animals

Experimental rats (80-100g) of either sex were obtained from the Physiology department, Faculty of Basic Medical Sciences, Bayero University, Kano. The animals were kept in cages and clean drinking water provided *ad libitum* while they were fed with standard commercial pelleted feed (Vital Feed® Nigeria). The temperatures varied between 27-30°C and relative humidity of about 55%-60% with 12-h light-dark cycle and adequate ventilation maintained in the animal house. Ethical conditions governing the conducts of experiments with life animals as stipulated were strictly observed. Also, the experimental protocol was approved by the College of Health Science ethical committee.

## 2.4 Experimental Design

A total of 20 rats were used for this experiment and they were divided into four groups of 5 rats each.

Group 1- normal rats

Group 2- treated with ethyl acetate extract of *P. americana* leaf.

Group 3- treated with ethyl acetate extract of *P. americana* leaf.

Group 4- treated with ethyl acetate extract of *P. americana* leaf.

The sub-chronic toxicity study lasted for a period of four weeks. The various doses (100, 200 and 400mg/kg b.w.) were administered to group 2, 3 and 4 respectively of normal rats. At the end of 4 weeks treatment the rats were sacrificed and blood samples collected for liver function (ALT, AST, ALP, ALB, TP

and GLO), kidney function indices [creatinine, urea, electrolytes ( Na<sup>+</sup>, K<sup>+</sup>, Cl<sup>-</sup> and HCO<sub>3</sub>)], lipid profile (CHOL, TG, HDL, LDL and VLDL) and hematological parameters (WBC, RBC, Hb, PCV, MCV, MCH, MCHC and PLT).

## **2.4 Assays**

Liver function test were assayed using the methods of; serum alanine and aspartate aminotransferase activity [21], serum alkaline phosphatase activity [22], serum albumin concentration [23], serum total protein concentration and serum globulin concentration [24]. Kidney function test were assayed using the methods of; serum urea concentration [25], serum creatinine concentration [26], serum potassium concentration [27], serum sodium concentration [28], serum chloride concentration [29] and serum bicarbonate concentration [30].

Lipid profile was analysed using the methods of; serum total cholesterol concentration [31], triglycerides concentration and serum HDL-cholesterol concentration [32], serum LDL and VLDL-cholesterol concentration [33]. Haematological parameters were analyzed using SYSMEX XE-200 (QBC Autoread Plus, UK).

## **2.5 Statistical Analysis**

The results were expressed as mean ± standard error of the mean and statistically analyzed by one-way analysis of variance (ANOVA). The level of significant was set at P < 0.05.

### 3. RESULTS

#### 3.1 Liver Function Test

Result of sub-chronic toxicity study showed no significant difference ( $P>0.05$ ) in AST and ALP activities in all the treated groups when compared with normal control. However, significant increase ( $P<0.05$ ) was observed in serum level of ALT of the entire treated group when compared with normal control (Figure 1). There was a significant decrease ( $P<0.05$ ) in serum total protein and globulin in all the treated groups compared with the normal control (Figure 2). The level of serum albumin was found to increase significantly ( $P<0.05$ ) in all the treated groups compared with normal control.

#### 3.2 Kidney Function Test

Kidney function tests carried out include serum urea, creatinine, sodium, potassium, chloride and bicarbonate (Figures 3 and 4). No significant difference ( $P>0.05$ ) was observed in serum urea, creatinine, sodium, chloride and bicarbonate in all the groups when compared with normal control group. Serum Potassium level was found to decrease significantly in group 2 (100mg/kg b.w.) When compared with normal and groups 3 and 4 (200 and 400mg/kg b.w.).

#### 3.3 Lipid Profile

The levels of total serum cholesterol, triglyceride and VLDL in the treated groups were found to decrease significantly ( $P<0.05$ ) when compared with normal control. No significant difference was observed ( $P>0.05$ ) in serum high density lipoprotein level. Serum low density lipoprotein level was reduced significantly ( $P<0.05$ ) in a dose dependent pattern within the treated groups when compared with normal control (Figure 5).

#### 3.4 Haematological Parameters

There was no significant difference ( $P>0.05$ ) between the normal control and the tested groups in all the haematological parameters assessed. These include, white blood cell, red blood cell, haemoglobin,

packed cell volume, mean corpuscular volume, mean corpuscular hemoglobin, mean corpuscular hemoglobin concentration and platelets (Table 1).


#### 4. DISCUSSION

Toxicity studies are essential tools in assessing the bioaccumulative effects of xenobiotics in biological systems. Assessment of liver and kidney function is very important in evaluating toxicity of modern and traditional medicines since these organs play major roles in metabolism of xenobiotics in the body. Elevated activities of liver enzymes are often diagnostic of underlying cellular injuries [34, 35]. From the biochemical result of this study, AST and ALP activities were not affected in all the groups administered with the extract. Serum concentration of ALT was found to increase in the treated groups, and this can be as a result of increase in hepatocytes proliferation or regeneration [36]. Chemicals often cause subclinical injury to the liver, which manifests only as abnormal liver enzyme tests. Certain medicinal agents, when taken in overdoses and sometimes even when introduced within therapeutic ranges may injure the organ [37]. Studies conducted by [38] showed that a large intake of tannins may cause liver damage. Reduction in total protein and globulin are indications of diminished synthetic function of the liver or might be due to impaired hepatocellular function [39, 40, 41, 42]. The increased level of albumin observed in this study depends on a number of factors such as the nutritional status, catabolism, hormonal factors, and urinary and gastrointestinal losses [42]. Ethyl acetate extract of *P. americana* was found to contain appreciable amount of saponins and tannins [8, 43] which can be related to the observed increase in ALT activity. Saponins may act by damaging cell membranes causing leakage of cellular materials, ultimately leading to cell death [44, 45].

Kidney function was evaluated by means of serum urea, creatinine and blood electrolyte concentrations. Urea is formed in the liver and is mainly excreted by the kidney, it is useful in evaluating kidney function in conjunction with creatinine which originates from the muscle and is filtered by the kidney. The serum urea and creatinine levels of treated groups in this study were not significantly changed. Increased blood urea and creatinine is a good indicator of compromised kidney function. These results suggest that the extract may not have altered the kidney function. In a similar study by [40] serum creatinine, urea and electrolytes such as sodium and calcium were not affected after *P. americana* extract treatment.


In this study, the administration of graded doses of EPAL significantly reduced serum levels of TC, TG, LDL and VLDL in treated rats. Saponins are known anti-nutritional factors, which lower cholesterol by binding with cholesterol in the intestinal lumen, preventing its absorption, and/or by binding with bile acids, causing a reduction in the enterohepatic circulation of bile acids and increase its faecal excretion [46, 47]. Increased bile acid excretion is offset by enhanced bile acid synthesis from cholesterol in the liver and consequent lowering of the plasma cholesterol; hence saponins have been reported to have hypocholesterolaemic effect [48]. Thus, the observed hypolipidemic effect of EPAL may be linked to the synergistic actions of phytochemicals like saponins and polyphenolic compounds contained in the plant extract.

The haematopoietic system is one of the most sensitive targets of toxic compounds and is an important index of physiological and pathological status in man and animals as it investigate the extent of damage to the blood [37]. It provides vital information regarding the status of bone marrow activity and intravascular effects such as haemolysis and anaemia [49]. Assessment of the haematological indices showed that the extract did not cause any significant effect on WBC, RBC, PCV, Hb, PLT, MCV, MCH and MCHC. The results on the haematological parameters indicate normal haemopoiesis and absence of anaemia confirming the non-toxic nature of the extract.



**Figure 1: Serum Levels of Aspartate Amino Transferase, Alanine Amino Transferase and Alkaline Phosphatase of Rats Administered with Ethyl acetate Extract of *P. americana* leaf (100, 200 and 400mg/kg b.w.) for 4 Weeks**


Results are presented as Mean  $\pm$  SD, n=5. Values with different superscripts are significantly different (P<0.05) with respect to normal control, NC= Normal control, EPAL= Ethyl acetate extract of *P. americana* leaf



**Figure 2: Serum levels of Total Protein, Albumin and Globulin of Rats Administered with Ethyl acetate Extract of *P. americana* Leaf (100, 200 and 400mg/kg b.w.) for 4 Weeks**


Results are presented as Mean  $\pm$  SD, n=5. Values with the different superscripts are significantly different (P<0.05) with respect to normal control, NC=Normal control, EPAL=Ethyl acetate extract of *P. americana* leaf






**Figure 3: Serum Levels of Urea and Creatinine Levels of Rats Administered with Ethyl acetate Extract of *P. americana* Leaf (100, 200 and 400mg/kg b.w.) for a Period of 4 Weeks**

Results are presented as Mean  $\pm$  SD, n=5. Values with the different superscripts are significantly different ( $P < 0.05$ ) with respect to normal control, NC= Normal control, EPAL= Ethyl acetate extract of *P. americana*



**Figure 4: Serum Levels of Sodium, Potassium, Chloride and Bicarbonate of Rats Administered with Ethyl acetate Extract of *P. americana* Leaf (100, 200 and 400mg/kg b.w.) For 4 Weeks**

Results are presented as Mean  $\pm$  SD, n=5. Values with the different superscripts are significantly different ( $P < 0.05$ ) with respect to normal control, NC= Normal control, EPAL= Ethyl acetate extract of *P. americana*



**Figure 5: Levels of Serum Total Cholesterol, Triglyceride, High Density Lipoprotein, Low Density Lipoprotein and Very Low Density Lipoprotein of Rats Administered with Ethyl acetate Extract of *P. americana* Leaf (100, 200 and 400mg/kg b.w.) for 4 Week**

Results are presented as Mean  $\pm$  SD, n=5. Values with the different superscripts are significantly different ( $P < 0.05$ ) with respect to normal control, NC= Normal control, EPAL= Ethyl acetate extract of *P. americana* leaf

**Table 1: Haematological Parameters of Rats administered with Ethyl acetate Extract of *P. americana* leaf (100, 200 and 400mg/kg b.w.) for 4 weeks**

<b>GROUPS</b>	<b>RBC(<math>10^3/\mu\text{l}</math>)</b>	<b>WBC(<math>10^6/\mu\text{l}</math>)</b>	<b>Hb(g/dl)</b>	<b>PCV(%)</b>	<b>MCV(FL)</b>	<b>MCHC(g/dl)</b>	<b>MCH(pg)</b>	<b>PLT(<math>10^3/\mu\text{l}</math>)</b>
<b>Normal Control</b>	6.45 ± .10 <sup>a</sup>	8.23 ± .20 <sup>b</sup>	7.60 ± .36 <sup>c</sup>	36.67 ± 1.86 <sup>d</sup>	55.30 ± .06 <sup>c</sup>	21.80 ± .23 <sup>f</sup>	11.23 ± .35 <sup>g</sup>	346.67 ± 3.22 <sup>h</sup>
<b>EE(100mg/kg)</b>	6.47 ± .14 <sup>a</sup>	8.73 ± .23 <sup>b</sup>	6.97 ± .27 <sup>c</sup>	37.67 ± 1.86 <sup>d</sup>	58.07 ± .54 <sup>e</sup>	22.93 ± .17 <sup>f</sup>	11.83 ± .19 <sup>g</sup>	336.67 ± 3.93 <sup>h</sup>
<b>EE (200mg/kg)</b>	6.73 ± .04 <sup>a</sup>	8.53 ± .35 <sup>b</sup>	8.03 ± .35 <sup>c</sup>	36.67 ± .88 <sup>d</sup>	55.20 ± .47 <sup>c</sup>	23.33 ± .98 <sup>f</sup>	11.40 ± .42 <sup>g</sup>	349.00 ± 8.50 <sup>h</sup>
<b>EE (400mg/kg)</b>	6.63 ± .22 <sup>a</sup>	8.77 ± .09 <sup>b</sup>	7.47 ± .35 <sup>c</sup>	37.67 ± 2.40 <sup>d</sup>	53.00 ± .78 <sup>e</sup>	22.23 ± .73 <sup>f</sup>	11.60 ± .40 <sup>g</sup>	357.33 ± 4.10 <sup>h</sup>

Results are presented as Mean ± SD, n=5. Values with the different superscripts in the same column are significantly different ( $p < 0.05$ ) with respect to normal control, EE= Ethyl acetate Extract of *P. americana* leaf

## CONCLUSION

While the medicinal importance of the *P. americana* is increasingly becoming of scientific interest, the consumption of high dosage should be discouraged, as oral administration of ethyl acetate extract of *P. americana* leaf results in an increase in the level of ALT. However, kidney function and haematological parameters were not affected with a significant decrease in the levels of total cholesterol, triglyceride, LDL and VLDL.

## ETHICAL APPROVAL

All authors hereby declare that; principle of laboratory animals care (NHI publication number 829 revised 1985) were followed, as well as all experiments have been examined and approved by the appropriate ethics committee.

## REFERENCES

1. Mashii JA, Atiku MK, Bala M, Sa'id AM, Idris RI, Babandi A and Babagana K et al. Hypoglycaemic and Hypolipidemic Properties of Ethyl acetate Fraction of *P. americana* Leaf in Alloxan-Induced Diabetic Rats. IJBBMB. 2018. In press. Doi: 10.11648/j.xxxx.2018xxxx.xx
2. Mashii JA, Atiku MK, Shehu D, Sa'id AM, Idris RI, Dangambo MA, Babagana K, Ya'u M, Babandi A. et al. Comparative study of different solvents extract of *Persea americana* leaf on alloxan induced hyperglycemic rats. Asian J. Biol. Sci. 2019a; 12: 67-72.
3. Tapsell LC, Hemphill I, Cobiac L. Health benefits of herbs and spices: the past, the present, the future. Med J Aust 2006; 185(4):S4–24.
4. Lai PK., Roy J. Antimicrobial and chemopreventive properties of herbs and spices. Curr Med Chem. 2004; 11(11):1451–1460.
5. Tene Tcheghebe O, Nyamen LD, Ngouafong Tatong F, Seukep AJ. Ethnobotanical uses, phytochemical and pharmacological profiles, and toxicity of *Persea americana* mill. An overview 2016; 3: 213-221.
6. Oluwole S, Yusuf K, Fajana O, Olaniyan D. Qualitative Studies on Proximate Analysis and Characterization of Oil from *Persea americana* (Avocado Pear). Journal of Natural Sciences Research. 2013; 3: 2.
7. Brai BI, Odetola AA, Agomo PU. Hypoglycemic and hypocholesterolemic potential of *Persea americana* leaf extracts. J Med Food. 2007; 10:356–360.
8. Mashii JA, Sa'id AM, Idris RI, Aminu I, Muhammad AA, Inuwa IM. Phytochemical, Radical Scavenging and Oxidative stress Potentials of Ethyl acetate Extract of *P. americana* Leaf in Alloxan-Induced Hyperglycemic Rats. Journal of Applied Sciences. 2019b; In press: 95063-JAS-ANSI

9. Mahadeva Rao US. Phytochemical screening and *in vitro* antioxidant and anti-diabetic potentials of *Persea americana* mill. (*lauraceae*) fruit extract. Universal Journal of Pharmaceutical Research. 2018; 3(5): 38-45
10. Lu QY, Arteaga JR, Zhang Q, Huerta S, Go VL, Heber D. Inhibition of prostate cancer cell growth by an avocado extract: role of lipid-soluble bioactive substances. J Nutr Biochem. 2005;16:23–30.
11. Nayak BS, Raju SS, Chalapathi Rao AV. Wound healing activity of *Persea americana* (avocado) fruit: a preclinical study on rats. J Wound Care. 2008; 17:123–126.
12. Kawagishi H, Fukumoto Y, Hatakeyama M. Liver injury suppressing compounds from avocado (*Persea americana*). Journal of Agriculture and Food Chemistry, 2001; 32(4): 128-131.
13. Adeyemi OO, Okpo SO, Ogunti OO. Analgesic and anti-inflammatory effects of the aqueous extract of leaves of *Persea americana* Mill (*Lauraceae*) Fitoterapia. 2002; 73:375–380.
14. Odetola AA, Akinloye O, Egunjobi C, Adekunle WA, Ayoola AO. Possible anti-diabetic and anti-hyperlipidemic effect of fermented *Parkia Biglobosa* extract in alloxan-induced diabetic rats. Clinical and Experimental Pharmacology and Physiology, 2006; 33(4):808-812.
15. Owolabi MA, Jaja SI, Coker HA. Vasorelaxant action of aqueous extract of the leaves of *Persea americana* on isolated thoracic rat aorta. Fitoterapia. 2005; 76:567–573.
16. Ojewole JA, Kamadyaapa DR, Gondwe MM, Moodley K, Musabayane CT. Cardiovascular effects of *Persea americana* Mill (*Lauraceae*) (avocado) aqueous leaf extract in experimental animals. Cardiovasc J Afr. 2007; 18:69–76.
17. Ozolua RI, Anaka ON, Okpo SO, Idogun SE. Acute and Sub-Acute Toxicological Assessment of the Aqueous Seed Extract of *Persea Americana* Mill (*Lauraceae*) in Rats. Afr J Tradit Complement Altern Med 2009; 6(4): 573–578.
18. Larrey D. Liver involvement in the course of phytotherapy. Presse Med. 1994; 23:691–693.
19. Ernst E. The efficacy of herbal medicine – an overview. Fundam Clin Pharmacol. 2005; 19:405–409.
20. Jagdish CN, Lalit SC. Antihyperglycemic and antihyperlipidemic activity of *Breynia vitis-idaea*. Pharmacognosy Journal. 2016; 8(3): 396-404.
21. Reitman S, Frankel SA. Colorimetric method for the determination of serum glutamic oxaloacetate and glutamic pyruvic transaminase. Amer.J . Clin . Patho. 1957; 28: 56.
22. Rec GSCC. Colorimetric method for serum alkaline phosphatase determination. J. Clin. Biochem, 1972; 10(2):182.
23. Grant GH. Amino acids and Proteins; *Fundamentals of Clinical Chemistry*. Tietz N.W. Editor, 3<sup>rd</sup> Edition, WB Saunders Company Philadelphia USA, Pp 328-329.
24. Tietz NW. *Clinical Guide to Laboratory Tests*. Second Edition W.B. Saunders Company, Philadelphia, USA, 1987; Pp 554-556.
25. Weatherburn, MW. Determination of serum urea. *Analytical Chemistry*, 1967; 39: 971.
26. Bartels H, Bohmer MA. Colorimetric Method for Determination of Serum Creatinine. *Clinica Chimica Acta*, 1972; 37: 193.
27. Henry JB. *Clinical Diagnosis and Management by Laboratory Methods*. 20<sup>th</sup> ed. Philadelphia, PA.: W. B. Saunders Company. 2001; Pp 186-193.
28. Kraut JA, Madias NE. Serum Anion Gap: Its Uses and Limitations in Clinical Medicine. Clinical Journal of the American Society of Nephrology, 2007; 2 (1): 162-174.
29. White WL. *Chemistry for Technologist*. 3<sup>rd</sup> Edition. The C.V. Mosby Co., St. Louis. 1970; Pp. 182.
30. Forrester RI, Wataji, IJ, Silverman DA, Pierre K.J. Enzymatic Method for Determination of CO<sub>2</sub> in Serum. *Journal of Clinical Chemistry*, 1976; 22: 243.
31. Lothar T. *Clinical Laboratory Diagnostic*. 1<sup>st</sup> edition TH-Books Verlagsgesellschaft mbh, Frankfurt/Main, Germany. 1998; Pp.169.
32. Jacobs D, Kasten BL, De Mott WR, Wolfson WL. Laboratory and Test Handbook. Lexi-company Inc: Hudson (Cleveland). 1990; Pp. 219.
33. Friedewald WT, Levy IR, Fredrickson DS. Estimation of ths concentration of low-density lipoprotein cholesterol in plasms, without use of the preparative ultracentrifuge. Clinical chemistry, 1972; 18(6): 206-228.

34. Karthikeyan, S., Gobianand, K., Pradeep, K., Mohan, C.V. and Balasubramanian, M.P. (2006). Biochemical changes in serum, lung, heart and spleen tissues of mice exposed to sub-acute toxic inhalation of mosquito repellent mat vapour. *Journal of Environmental Biology*, 27(5): 355-358.
35. Mosa RA. Some Bioactivity of Triterpenes from Stem Bark of *Protorhus longifolia* and their Derivatives. *Planta Med*, 2014; 6(2): 50-58.
36. Mathew Folaranmi Olaniyan, Effect of Liquid Extract of Pear Avocado Leaf (*Persea americana*) on Plasma Levels of Aminotransferases, Cholesterol and Total Bile Acids in Hypertensive Patients, *American Journal of Medicine and Medical Sciences*, 2014; 4(3): 87-91.
37. Omodamiro OD, Jimoh MA, Ewa IC. Hepatoprotective and Haemopoietic Activity of Ethanol Extract of *Persea americana* Seed in Paracetamol Induced Toxicity in Wistar Albino Rat. *Human Journals*, 2016; 5(3): 149-165.
38. Yamasaki T, Sato M, Mori T, Mohammed AM, Fujii K, Tsukioka J.. Toxicity of tannins towards the free-living nematode *Caenorhabditis elegans* and the brine shrimp (*Artemia salina*). *Journal of Natural Toxin*, 2000; 11(3): 165-171.
39. Haeri MR, Izaddoost M, Ardekani MR, Nobar MR, White KN. The effect of fenugreek 4-hydroxyisoleucine on liver function biomarkers and glucose in diabetic and fructose-fed rats. *Phytother Res*, 2009; 23(1): 61-64.
40. Olorunnisola, OS, Bradley G, Afolayan AJ. 2009. Acute and sub-chronic toxicity studies of methanolic extract of *Tulbaghia violacea* rhizomes in Wistar rats. *African Journal of Biotechnology*, 2009; 11(83): 14934-14940.
41. Yakubu MT, Bilbis, LS, Lawal M, Akanji MA. Effect of repeated administration of sildenafil citrate on selected enzyme activities of liver and kidney of male albino rats. *Niger. J. Pure Appl. Sci*, 2003; 18: 1395-4000.
42. Ilmie MU, Jaafar H, Mansor SM, Abdullah JM. Subchronic toxicity study of standardized methanolic extract of *Mitragyna speciosa* Korth in Sprague-Dawley Rats. *Front. Neurosci*, 2015; 9: 189.
43. Rahman A, Zareen S, Choudhary MI, Akhatar MN, Khan SN. A- Glucosidase inhibitory activity of triterpenoids from *Cichorium intybus*. *Journal of Natural Products*, 2008; 71: 910-913.
44. Aliyu MS, Lawal U, Tijjani MB, Doko MH, Garba I, Kokya HA. Phytochemical and Antibacterial Properties of Leaf Extracts of *Ipomoea asarifolia*. *Journal of Ethnopharmacology*, 2011; 19: 236-240.
45. Sreeshma LS, Nair BR. Brine Shrimp Lethality Assay in Two Species of *Biophytum Dc* (*Oxalidaceae*). *Innovare Academic Sciences*, 2014; 6(4): 5-7.
46. Nimenibo-uadia R. Effect of aqueous extract of *Canavalia ensiformis* seeds on hyperlipidemic and hyperketonaemia in alloxan-induced diabetic rats. *Biokemistri*, 2003; 15(2): 7-15.
47. Gouegni EF, Abubakar H. Phytochemical, Toxicological, Biochemical and Haematological Studies on Avocado (*Persea americana*) in Experimental Animals. 2013; NIFOJ; 31(1): 64 – 69.
48. James DB, Owolabi OA, Ibrahim AB, Folorunsho DF, Bwalla I, Akanta F. Changes in lipid profile of aqueous and Ethanolic extract of *Blighia sapida* in rats. *Asian Journal of Medical Sciences*. 2010; 2(4): 177- 180.
49. Adeneye AA, Ajagbonna OP, Adeleke TI, Bello SO. Preliminary Toxicity and Phytochemical Studies of the Stem Bark Aqueous Extract of *Musanga cecropioides* in Rats. *Journal of Ethnopharmacology*, 2006; 105(4): 374-379.