

Adoption of Agroforestry Practices in Katsina State, Nigeria

ABSTRACT

Unsustainable forest land use practices have resulted in land degradation in Nigeria leading to low crop yield. Agroforestry is a viable option for reversing dwindling crop yields through proper soil management practices. There is notably no sufficient published information on agroforestry practices (AP) in Katsina State. The aim of this study was to assess AP for sustainable land use in the study area. Multistage stratified sampling design was used to select respondents from the study. Three Local Government Areas (LGAs) were randomly selected from each of the agro-ecological zones (Sahel, Sudan and Guinea) of Katsina State. Within each of the selected LGAs, one community was randomly selected and forty respondents were randomly sampled from each community. Using structured questionnaire, information was sought on socio-economic and AP. Data were analysed using descriptive statistics. Based on the results, multipurpose trees on farmland (79%), windbreaks (50%), woodlots (49.7%), improved fallow in shifting cultivation (32%) and home gardens (24.7%) were the common AP in the study area. Benefits of AP in the area included preservation of the environment (98.5%), provision of fruits and leaves (98%), and improvement of soil fertility (98%), erosion control (98%) and improvement of farmers' income (96%). Agroforestry practices enrich the soil with important nutrients and prevent soil erosion. The adoption of multipurpose trees on farmland in the study area will help in preventing environmental degradation, desertification and enhance food crop production.

Keywords: Agroforestry, Environmental, Nutrients, Sustainable, Soil, Degradation

INTRODUCTION

Cropping system and anthropogenic activities are the major factors affecting sustainable land use in most developing countries. The result is continuous stress on the natural resources base with the conversion of forested areas into croplands, the cropping of marginal lands and the use of adverse agricultural methods including inappropriate and excessive application of agro-chemicals.

Environmental degradation has intensified during the worldwide population boom, just within about a quarter of a century the population has doubled (Repetto, 1987). The concomitant demands for living space, and higher food and energy production have resulted in some lands being converted to intrinsically unsuitable ones. Slash-and-burn farm plots cut out of rain forests, which at best will support only a few years of crops, are one example of land use practices, that were nearly harmless in an area of low population densities and resource demand, but are fast becoming unsuitable. It has been reported by various authors that shifting cultivation can no longer support the needs of the farmers in Nigeria because of the increasing population pressure and attendant short fallow periods resulting in soil deterioration and ecological imbalance (Kang *et al.*, 1984; Adeola and Ola, 1985; Kang and Wilson, 1987).

35 The need for more food has led to increased deforestation, shortened fallow periods in shifting cultivation
36 cycles, and set in motion a degradation spiral, leading to reduced productive capacity of the land and
37 decreased crop yields. In addition, indiscriminate fire wood gathering, timber harvesting and grazing have
38 aggravated land degradation in many parts of the tropics (Bene *et al.*, 1977, Poulsen, 1978, and Gorse,
39 1985).

40 Besides land degradation, deforestation also has implications for regional and global climate. On a global
41 basis, it contributes about 20% of the annual carbon dioxide added to the atmosphere (Anon, 1990). This
42 development is disturbing because the rising level of atmospheric carbon dioxide will bring about a global
43 warming through the so-called “green house effect”. Carbon dioxide taps the sun’s energy, thus causing
44 the temperature to rise. The accompanying increase in global temperature could directly affect agricultural
45 production (Swaminathan, 1987). Rainfall patterns are also disturbed by large-scale deforestation, and this
46 leads to unpredictable weather, which in turn affects crop yields.

47 Anon (1988) reported that a mixture of tree and annual crops of different heights (an agroforestry
48 practice) provide a more complete ground cover which again helps protect the soil from erosion and
49 makes maximum use of available sunlight. Gatherson (1982) opined that agroforestry seeks to develop
50 sustainable land-use systems that supply people’s needs for food and other basic necessities while
51 maintaining critical environmental stability. The problem of insufficiency of food production to which
52 successive governments in Nigeria have attempted to look for a probable solution, suggests adoption of
53 agroforestry options.

54 Population pressure today precludes practices that would enable the environment to recover from
55 extensive cropping systems or anthropogenic activities. It is expected that this study will assist
56 agricultural planners and policy makers to properly address the problem of environmental and soil
57 degradation. This paper is aimed at identifying the various agroforestry practices adopted by farmers; and
58 evaluate farmers’ use of agroforestry practices in the study area with a view to ascertaining the benefits
59 derived from planting trees along with food crops in the study area.

60 **MATERIALS AND METHODS**

61 **The Study Area**

62 Katsina State lies between latitude 12° N and longitude 8° E. The state is located in the upper Sudan and
63 lower Sahel regions of Nigeria. The entire landmass of Katsina State is composed of basement complex of
64 pre- Cambrian era, which consists of meta-sediments that have been transformed into anatectic

65 migmatites and granites. The hot, dry and dust laden North East Trade wind predominates in this area for
66 as long as 7 months of the year. Rainfall is experienced in the state from June to September; with mean
67 annual rainfall from 1016 mm to 1143mm in the south and less than 635 mm in the Northern part of the
68 state. The state on the whole has a mean annual rainfall of about 840 mm. Mean relative humidity is lower
69 than 50 % in January and February and could be as high as 80 % in June – July. Temperature range is
70 often from 38^o – 41^oC.

71 **Sampling techniques and Questionnaire surveys**

72 Multi-stage and stratified random sampling techniques were used for the study. Nine Local Government
73 Areas (LGAs) (three per each agro-ecological zone) were randomly selected out of the 34 LGAs in
74 Katsina state. Accurate data on the actual population of the rural farmers in Katsina state were not readily
75 available; therefore equal number of farmers (120 farmers) were randomly sampled from each of the
76 agro-ecological zones through the use of random numbers. This gave a sample size of 360. Questionnaire
77 was used to collect information on the study objectives. The questionnaire was sub-divided into: source of
78 farm land, farm location and size; other agricultural activities, types of crops and trees planted and
79 estimated farm yield, estimated agricultural income, source of information and the perceived benefits of
80 AP. Information was also obtained through personal observation.

81 **Statistical analyses and data presentation**

82 The data obtained from the study were collated and analysed. Descriptive statistics such as frequencies
83 and percentages were used to describe the variables and their occurrence among the respondents. Results
84 were presented in tables and graphs for clarity.

85 **RESULTS**

86 **Demographic Attributes of the Respondents in the Study Area**

87 The demographic characteristics of the respondents in the study area showed the males were more
88 involved in agroforestry practices than the females. Most of the respondents in the Sahel and Sudan
89 Savanna, were between 30 and 49 years; while in the Guinea savanna, majority of the respondents age
90 ranged from 60 years and above. The age distribution is an important factor in farming activities because
91 it affects the work force and decision-making in farming activities. The dominant age bracket among rural
92 dwellers (30 – 49 years) in Sahel and Sudan and ≥ 60 years in Guinea savanna was indication that this
93 was the age bracket that was actively involved in agroforestry practices.

94 Majority of the respondents in the three agro-ecological zones were married men and women. Most
95 marriages were polygamous and had an average of more than five children that provided labour force for
96 farming. The result showed that 93.3 % of the respondents were married, 4.2 % were single, 0.8 %
97 divorced and 1.7 % widowed. About 92 % of the respondents were Muslims, 4.6% were mixture of Islam
98 and Traditional religion, 2.6 % and 0.9 % Christians and traditional worshippers, respectively.

99 Based on these findings, majority of the respondents acquired Islamic education as
100 their highest educational attainment. On average, it represented 41.0 % across the
101 three zones. This was followed by primary education with 17.9 %.

102 Farming was the major occupation in the three agro-ecological zones of the study area.
103 The study revealed that an average of 66.1 % of the respondents as farmers. Other
104 occupation in the study area included trading, civil service, fishing and cattle rearing
105 among others.

106 **Involvement of Respondents in the Use of Agro-forestry practices (AP)**

107 Based on the list of agro-forestry practices identified by ICRAF, Nair₅ (1990), some agro-forestry
108 practices that were adopted in the study area were selected and the respondents were made to react to the
109 practices, that is, their involvement in the use of any of the practices. The responses were classified into
110 five: do not practise, practised but stopped, practise occasionally, practised but do not intend to continue,
111 and practise regularly (Table 1 to 2).

112 Multipurpose trees Agroforestry system on farmland was mostly adopted in the study area (Table 1). The
113 highest was in Sahel savanna with 91.7 %, followed by Guinea savanna (83.3%) while Sudan had 62.5%
114 adoption. Farmers who had never practised at all, practised but stopped, practised occasionally or
115 practised but did not intend to continue were negligible.

116 Also, Table 1 shows majority of the respondents in Sahel and Guinea savanna had never practised home
117 gardens, with 54.2% each. Those farmers had adopted the system were relatively few, (Sahel 35.0%,
118 Sudan savanna 20.8%, Guinea savanna 18.3%), while those who had practised but stopped, practiced
119 occasionally or practised but did not intend to continue recorded very low scores.

120 Table 1 indicates that majority of the respondents across the zones had not adopted *Taungya* system of
121 agro-forestry (Sahel savanna 91.7%, Sudan savanna 87.5% and Guinea savanna 81.7%). Only very few of
122 the respondents had adopted the *Taungya* system of agro-forestry (Sahel 5.0, Sudan savanna 8.3 and
123 Guinea Savanna 15.0%). No farmer had adopted the system but stopped practising it across the zones.

124 The Table also shows 55.0 % of the respondents in Sahel savanna had adopted woodlot system which was
 125 the highest across the zones, thus was followed by Sudan savanna with 50.0%, and Guinea savanna with
 126 44.2%. Those that had never adopted the system were 37.5 % in Guinea savanna, 35.0 % in Sahel
 127 savanna, and 26.7% in Sudan savanna.

128 **Table 1: Adoption of Multipurpose Trees, home-gardens, *Taungya* System and Woodlots on**
 129 **Farmland in Katsina State**

Variables	Sahel		Sudan		Guinea	
Multipurpose Trees	Freq.	%	Freq.	%	Freq.	%
Do not Practise	4	(3.3)	15	(12.5)	7	(5.8)
Practised but Stopped	0	(0)	10	(8.3)	3	(2.5)
Practise Occasionally	6	(5.0)	8	(6.7)	5	(4.2)
Practised but do not Intend to Continue	0	(0)	12	(10)	5	(4.2)
Practise Regularly	110	(91.7)	75	(62.5)	100	(83.3)
Total	120	(100)	120	(100)	120	(100)
Home-gardens						
Do not Practise	65	(54.2)	48	(40)	65	(54.2)
Practised but Stopped	0	(0)	32	(26.7)	0	(0)
Practise Occasionally	13	(10.8)	10	(8.3)	33	(27.5)
Practised but do not Intend to Continue	0	(0)	5	(4.2)	0	(0)
Practise Regularly	42	(35.0)	25	(20.8)	22	(18.3)
Total	120	(100)	120	(100)	120	(100)
<i>Taungya</i> system						
Do not Practise	110	(91.7)	105	(87.5)	98	(81.7)
Practised but Stopped	0	(0)	0	(0)	0	(0)
Practise Occasionally	4	(3.3)	0	(0)	0	(0)
Practised but do not Intend to Continue	0	(0)	5	(4.2)	4	(3.3)
Practise Regularly	6	(5.0)	10	(8.3)	18	(15.0)
Total	120	(100)	120	(100)	120	(100)
Woodlots						
Do not Practice	42	(35)	32	(26.7)	45	(37.5)
Practised but Stopped	0	(0)	18	(15.0)	7	(5.8)
Practise Occasionally	0	(0)	10	(8.3)	15	(12.5)
Practised but do not Intend to Continue	12	(10.0)	0	(0)	0	(0)
Practise Regularly	66	(55.0)	60	(50)	53	(44.2)
Total	120	(100)	120	(100)	120	(100)

130 **Adoption of border planting, windbreaks/shelterbelt, Alley Cropping System and Woody Perennial**
 131 **for Sustainable Soil Management**

132 A distribution of the respondents according to adoption of border planting as shown in Table 2, the Table
 133 indicates majority of the respondents had never adopted the system across the zones (Sahel savanna
 134 81.7%, Sudan savanna 74.2% and Guinea savanna 91.7%). No respondent adopted the system but stopped
 135 practising it. Sudan savanna recorded the highest number of adopters with 18.3 %, followed by Sahel
 136 savanna (10.0%), while Guinea savanna recorded only one adopter (0.8%). Majority of the respondents
 137 across the Sahel and Sudan savanna zones had adopted the windbreaks system of agro-forestry. The

138 respondents from the Sahel savanna, however, recorded the highest number of adopters with 65.0 %,
 139 followed by Sudan savanna with 55.0 %, while Guinea savanna recorded only 30.0 %. Conversely, for the
 140 respondents that had never adopted the system, Guinea savanna recorded the highest number with 52.5%
 141 (Table 2).

142 Table 2 indicates that majority of the respondents across the entire zones had never adopted the alley
 143 cropping system (Sahel savanna 95%, Sudan savanna 80% and Guinea savanna 97.5%). Very few of the
 144 respondents had adopted the system (Sahel savanna 3.3%, Sudan savanna 7.5% and Guinea savanna
 145 1.7%). No farmer from Sahel and Guinea savanna had adopted the system and stopped, but in the Sudan
 146 savanna 6.7% had. Similarly, for those that practised the system occasionally, the Sahel and Guinea
 147 savanna also recorded zero %, while Sudan had 2.5%. The Table also shows that respondents that had not
 148 adopted the system were highest in Guinea savanna with 87.5 %. This was followed by Sahel savanna
 149 with 81.7, while Sudan savanna recorded only 31.7%. On the other hand, the Sudan savanna recorded the
 150 highest number of adopters for this system, with 53.3 %; while Sahel and Guinea savanna had very low
 151 scores. No respondents had adopted the system but stopped practising it or did not intend to continue to
 152 do so across the zones.

153 **Table 2: Adoption of border planting, windbreaks/shelterbelt, Alley Cropping System and Woody**
 154 **Perennial in the Study Area**

border planting	Sahel		Sudan		Guinea	
	Freq.	%	Freq.	%	Freq.	%
Do not Practise	98	(81.7)	89	(74.2)	110	(91.7)
Practised but Stopped	0	(0)	0	(0)	0	(0)
Practise Occasionally	0	(0)	5.0	(4.2)	9	(7.5)
Practised but do not Intend to Continue	10	(8.3)	4	(3.3)	0	(0.0)
Practise Regularly	12	(10.0)	22	(18.3)	1	(0.8)
Total	120	(100)	120	(100)	120	(100)
windbreaks/shelterbelt						
Do not Practise	39	(32.5)	24	(20)	63	(52.5)
Practised but Stopped	0	(0)	10	(8.3)	8	(6.7)
Practise Occasionally	3	(2.5)	6	(5.0)	3	(2.5)
Practised but do not Intend to Continue	0	(0)	14	(11.7)	10	(8.3)
Practise Regularly	78	(65.0)	66	(55.0)	36	(30.0)
Total	120	(100)	120	(100)	120	(100)
Alley Cropping System						
Do not Practice	114	(95)	96	(80.0)	117	(97.5)
Practised but Stopped	0	(0)	8	(6.7)	0	(0)
Practise Occasionally	0	(0)	3	(2.5)	0	(0)
Practised but do not Intend to Continue	2	(1.7)	4	(3.3)	1	(0.8)
Practise Regularly	4	(3.3)	9	(7.5)	2	(1.7)
Total	120	(100)	120	(100)	120	(100)
Woody Perennial						
Do not Practised	98	(81.7)	38	(31.7)	105	(87.5)
Practised but Stopped	0	(0)	0	(0)	0	(0)
Practised Occasionally	9	(7.5)	18	(15.0)	5	(4.2)

Practised but do not Intend to Continue	0 (0)	0 (0)	0 (0)
Practised Regularly	13 (10.8)	64 (53.3)	10 (8.3)
Total	120 (100)	120 (100)	120 (100)

155 **Tree species combined with agricultural crops among the respondents in the Study Area**

156 Table 3 shows some of the commonest trees used in combination with agricultural crops among the
 157 respondents in the study area. In the Sahel savanna zone, those tree species with high scores include:
 158 *Azadirachta indica* with 25.8% *Parkia biglobosa* with 14.2% and *Adansonia digitata* having 13.3%. In
 159 Guinea savanna, those species recorded in decreasing number include: *Adansonia digitata*, with 9.2%,
 160 *Parkia biglobosa*, with 6.7% and *Azadirachta indica* having 5.8%. Across the three zones, the highest
 161 score is recorded by *Azadirachta indica*, having 13.3%, while *Adansonia digitata* and *Parkia biglobosa*
 162 recording 9.7% respectively. Other tree species recorded include: *Borassus aethiopum*, *Anacardium*
 163 *occidentale* and *Tamarindus indica*.

164 **Table 3: Identified Tree Species Combined with Agricultural Crops Among the Respondents.**

Tree Species	Sahel		Sudan		Guinea		Mean %
	Freq.	%	Freq.	%	Freq.	%	
<i>Adansonia digitata</i>	16	(13.3)	8	(6.7)	11	(9.2)	9.7
<i>Azadirachta indica</i>	31	(25.8)	10	(8.3)	7	(5.8)	13.3
<i>Anacardium occidentale</i>	2	(1.7)	-	-	3	(2.5)	1.4
<i>Borassus aethiopum</i>	2	(1.7)	-	-	01	(0.8)	0.8
<i>Parkia biglobosa</i>	17	(14.2)	6	(5.0)	8	(6.7)	8.6
<i>Tamarindus indica</i>	-	-	-	-	6	(5.0)	1.7
Combination/others	52	(43.3)	96	(80)	84	(70)	64.4
Total	120	(100)	120	(100)	120	(100)	(100)

165 **Constraints to adoption of agro-forestry practices in the study area**

166 Table 4 reveals some of the problems militating against the adoption of agroforestry practices in the study
 167 area. The most serious problem preventing adoption of AP in Sahel savanna zone was scanty rainfall,
 168 which accounted for (30.8 %) of the problems. This was followed by lack of land (6.7 %) and inadequate
 169 labour (5.0 %). In the Guinea savanna zone, the most serious problem was high labour demand (8.3 %),
 170 followed by lack of land (6.7 %). Lack of required seedlings and scanty rainfall had 5.0 % respectively.
 171 The problems, in decreasing order of severity across the three zones were: scanty rainfall, lack of land,
 172 high labour demand, inadequate extension personnel and lack of transportation recording the same
 173 percentage, lack of required seedlings and lack of incentives recorded the same percentage.

174
 175
 176

177 **Table 4: Identified constraints to adoption of agro-forestry practices in the study area**

Identified Problems	Sahel		Sudan		Guinea		Mean %
	Freq.	%	Freq.	%	Freq.	%	

High Labour Demand	6	(5.0)	-	-	10 (8.3)	4.4
Lack of Required Tree Seedlings	3	(2.5)	-	-	1 (0.3)	1.1
Inadequate Extension Personnel	2	(1.7)	-	-	4 (3.3)	2.5
Lack of Land	8	(6.7)	1	(0.8)	8 (6.7)	4.73
Scanty Rainfall	37	(30.8)	3	(2.5)	6 (5.0)	12.8
Lack of Transportation	4	(3.3)	1	(0.8)	4 (3.3)	2.5
Lack of Incentive	2	(1.7)	-	-	2 (1.7)	1.1
Combination	58	(48.3)	115	(95.8)	80 (66.7)	70.3
Total	120	(100)	120	(100)	120 (100)	(100)

178 DISCUSSION

179 Socio-economic Attributes of the Respondents in the Study Area

180 Respondents' gender showed that there were more male than female farmers across the zones. This
 181 implies that the male gender is more involved in agroforestry practices as compared to their female
 182 counterparts. Farming generally, is almost an exclusive business for the male in the study area. This may
 183 be as a result of the strenuous nature of most farming activities in general and agroforestry practices in
 184 particular. These activities are not attractive to women who often engage in household activities. The
 185 paucity of women in agroforestry practices might also be attributed to culture and religion which made
 186 access to women by male extension agents difficult since there were very few women extension agents.

187 In Sahel and Sudan savanna zones, majority of the respondents fell between 30 and 49 years. Except in
 188 Guinea savanna, where majority of the respondents fell from ≤ 60 years.

189 This dominant age bracket among the rural dwellers implied that this is the age bracket that is actively
 190 involved in agroforestry practices. This also meant that, the farming population is mainly made up of both
 191 the old and the middle-aged people. The young ones might have migrated to urban areas in search of
 192 white collar jobs while others might have been in school or too young to have farms. This has an adverse
 193 effect on the economy of the rural people because at old age, farmers ~~can not~~ cannot have optimum
 194 productivity.

195 Also, majority of the respondents across the three zones were married men and women. Most marriages
 196 were polygamous and have an average of more than five children that provided labour force for farming.
 197 This is because agroforestry practices are not only capital intensive but also labour intensive.

198 On religious affiliation, most of the respondents were Muslims while other religious included Christians
 199 and traditional worshipers.

200 As revealed in the study, educational level of the rural dwellers was low. However, on average, Islamic
201 education recorded the highest %_age, which was followed by primary education. The study also revealed
202 that inspite of the low level of western education, they had indigenous/traditional knowledge and high
203 level of awareness about farming systems, tree species, shrubs, herbs and other agroforestry practices.
204 This indigenous knowledge affects their perception and willingness to participate in agroforestry
205 practices. However, they still needed more enlightenment and training on modern agroforestry techniques
206 as a means of sustainable land management.

207 Frequency analysis of the respondents' occupation revealed that farming is the major occupation in the
208 three agro-ecological zones of the study area. The study reveals on average, a high % of the respondents
209 as farmers. Other occupation in the study area includes: trading, civil servants, fishing and cattle rearing
210 among others.

211 **Adoption of Agro-forestry by the Respondents in the study Area**

212 Agro-forestry entails combination of agricultural crops with tree crops together with pastures or animals
213 on the same piece of land either in sequence or at the same time. This means that agroforestry facilitates
214 multiple land use. The study shows that majority of the respondents were involved in food crop-
215 production, while others were involve in rearing of animals, and planting of tree crops such as
216 *Azadirachta indica*, *Parkia biglobosa*, *Adansonia digitata* among others.

217 Respondents' involvement in AP varied from zone to zone. The differences in adoption could be that an
218 innovation which was appropriate for a given zone might not necessarily be accepted in another zone. It
219 might also be due to socio-economic reasons, complexity and incompatibility of the innovation with the
220 existing practices. Thus majority of respondents across the zones could not adopt *Taungya*, border
221 planting and alley cropping. Very few respondents across the zones adopted these systems of agro-
222 forestry. Conversely, multipurpose trees on farmland, improved fallow in shifting cultivation, home-
223 garden, woodlots and windbreaks or shelterbelt were much more adopted by farmers. The reason for low
224 adoption of *Taungya* system of agroforestry might be that food crop might compete with tree crop. Sahel
225 and Sudan zones adopted woodlots practices more than Guinea savanna.

226 Majority of the respondents adopted multipurpose trees on farmland across the zones. The findings
227 therefore reveal that this agroforestry system was popular among the farmers across the zones hence the
228 massive adoption. This might be due to the blend of the system with indigenous or traditional farming
229 practice across the zones.

230 | The study also revealed that majority of the respondents in ~~sahel-Sahel~~ and ~~guinea-Guinea~~ savanna had
231 not practiced home-gardens, as the two zones recorded high scores each. Those farmers that had adopted
232 the system were relatively few. While those that had practiced but stopped, practice occasionally or
233 practiced but do not intend to continue recorded very low scores.

234 The reason for their adoption might be to stem the environmental degradation in the Sahel and Sudan
235 savanna zones. Farmers could only take fuel-wood from these plantations and no other place.
236 Indiscriminate felling of trees for timber, fuel-wood and other domestic uses and clearing of land for
237 agricultural purposes and industrial development help to remove the forest cover; thereby exposing the
238 soil to wind erosion (Repetto, 1988). Adeola (2001) observed that the system is used for various purposes
239 such as provision of wood, fodder, electric-poles, fencing poles, roofing poles, soil protection, soil
240 reclamation etc.

241 Sudan Savanna had the highest adoption of woody perennials for soil conservation across the zones. The
242 farmers' interest and adoption of the system could be to check the menace of annual flooding of this zone
243 which leads to soil and gully erosion. Plants help to stabilize the soil and other conservation works
244 thereby fulfilling one of the environmental functions of agroforestry (Baumer, 1990).

245 Majority of the respondents adopted multipurpose trees on farmland across the zones;
246 this could be due to good yield obtained if tree species are combined with agricultural
247 crops in the study area. This implies that native tree species enhance high yield of
248 agricultural crops when combined; agroforestry system was popular among the
249 farmers across the zones hence the massive adoption. This might be due to the blend
250 of the system with indigenous or traditional farming practice across the zones.

251 On sources of information/awareness on some sustainable land use practices, the study revealed that
252 extension agents recorded high scores for all the land use practices in the study area. This may be due to
253 the availability of the agricultural development programme in the area. This study therefore agreed with
254 the findings of Onumadu (2002) who observed that agricultural agents were the most important source of
255 agricultural information to farmers. This view was also supported by Azeez (2002).

256 Majority of the respondent were of the view that they obtained information on sustainable land use
257 practices/agro-forestry practices through radio. This implies that farmers in these zones had several
258 options of other sources of information that could enhance or stimulate their use of AP. Although,

259 radio/mass media and extension agents were the principal means of the awareness, these two sources of
260 information could as well be responsible for the significance of the AP.

261 Mass media therefore plays an important role in the dissemination of information on agricultural activities
262 as it enables even the cattle rearers that roam about in the bush to have access to the information on
263 agricultural activities through their radio. This was also supported by Onumadu (2002), who observed that
264 mass media was one of the most important sources of seeking information on agro-forestry practices.

265 Other sources such as traditional and a combination of one or more of these sources also recorded
266 relatively high scores, whereas sources such as relatives and neighbour recorded very low scores; while
267 some of the respondents reported that they had no information at all. This may be due to lack of adequate
268 publicity or enlightenment. This calls for an increase in agricultural extension agents who should take up
269 the responsibility of educating, training and monitoring of farmers in the areas of food crop production.

270 Farmers in the study area may have adopted AP because of the various benefits they derive from it. These
271 benefits range from social, economic and environmental benefits. The social benefits in the study area
272 include provision of fruits and leaves, provision of shade, provision of fuel-wood, provision of fodder and
273 medicinal herbs.

274 **CONCLUSION**

275 Based on the findings of this study, windbreaks, scattered trees on farmland, woodlots, improved fallow
276 and home-gardens were the various AP that were common in the study area. There was difference in
277 adoption of AP among the three agro-ecological zones. The differences in adoption could be that an
278 innovation in which was appropriate for a given zone might not necessarily be accepted in another zone.
279 This could also be due to soil and climate type and socio-economic reasons. In order to sustain and even
280 increase our agricultural productivity and to reduce, to the barest minimum, the effects of desertification
281 and environmental degradation, the following recommendations are made: Government should encourage
282 the adoption of agroforestry as a system of multiple land use to increase wood and food production
283 thereby ensuring the optimum use of land. Provide incentives such as seedlings, transportation, inorganic
284 fertilizers and tractors to farmers so as to encourage them to participate actively in agroforestry activities.
285 The use of more indigenous tree species that can improve soil fertility and at the same time more
286 adaptable to the environment should be promoted (eg. *Parkia biglobosa*). Application of organic
287 fertilizers and planting of leguminous trees will help to resuscitate the soil for high productivity. A study
288 to re-examine the factors limiting the adoption of some AP that have low adoption in the study area such
289 as alley cropping and *Taungya* is recommended.

290

291

292 **REFERENCES**

293 Adeola, A.O. (2001). Farmer's perception of social forestry impact in the semi-arid zone of Nigeria.
294 *Unpublished Ph.D. Thesis*, Department of Forest Resources Management, University of Ibadan,
295 275 Pp.

296 Adeola, A.O. and Ola – Adams, B.A. (1985). Agroforestry and small scale farmers In: proceedings of the
297 15th Annual conference of the Forestry Association of Nigeria, Yola 408 – 417.

298 Anon (1990): Environment: The danger ahead. Global Edition. Making the world and oven. June 11
299 publication of Newswatch Organization, Lagos, Nigeria.

300 Anon, I. (1988). *Modernization of Agriculture in Developing Countries*, England: A whiley Inter-Science
301 Publication, 565 Pp.

302 Azeez I.O. (2002): Evaluation of media mix for disseminating conservation information in South Western
303 Nigeria. Ph.D. Thesis (unpublished), Department of Forest Resources Management, University of
304 Ibadan, Nigeria, 230 Pp.

305 Baumer, M. (1990): Agroforestry and Desertification. Technical Centre for Agricultural and Rural
306 Cooperation. *The Netherlands* 250 Pp.

307 Bene, J.G., Beall, H.W. and Cote, A. (1977). *Trees, food and people; Land management in the Tropics*.
308 Ottawa, IDRC. 89 Pp.

309 Eboh, E. (1995). Poverty, Population Growth and Environmental Degradation: A vicious cycle of human
310 misery: In: Eboh, E., Okoye and Ayidu (eds). Rural development in Nigeria Auto Century
311 publication publishing co. pp 274.

312 FAO (2007): Adaptation to Climate Change in Agriculture, Forestry and Fisheries: Perspective
313 Framework and Priorities FAO, United Nations, Rome 79 Pp.

314 Gatherson, T.M. (1982): Agroforestry production systems: putting them into action. In: L.H. Mac. Donald
315 (ed). *Agroforestry in African Humid Tropics*, United Nations University, Tokyo. Pp 128 – 133.

316 Gorse, J. (1985): Fuel wood and Domestic Energy. *The fuel wood crisis in tropical West Africa*,
317 Washington D.C. World Bank 46 Pp.

318 Kang, B.T. and Wilson, G.F. (1987). The development of alley cropping as a promising agroforestry
319 technology. Reprint from: *Agroforestry: a Decade of Development*. Stepler, H.A. and Nair,
320 P.K.R. (eds). ICRAF, Nairobi, Kenya.

321 Kang, B.T., Wilson, G.F. and Lawason, T.L. (1984). Alley cropping: *A stable alternative to shifting*
322 *cultivation*, Ibadan, Nigeria., IITA. 23 pp.

323 Nigerian Environmental study/action team, NEST (1995). *The challenge of sustainable development in*
324 *Nigeria*. Edited by Tade Aina and Ademola, T. Salawu. 80 Pp.

- 325 Okali, D.U.U. (1997): Environment and Resources Development: Towards Sustainable Forestry
326 Development in Nigeria. In: Oduwaiye E.A, Obiaga, P.C. and Abu, J.E. (eds). Proceedings of 25th
327 Annual Conference of FAN held in Ibadan, Nigeria Pp 2 – 4.
- 328 Okojie, J.A. (1991): Misuse of renewable natural resources and environmental degradation. An invited
329 paper presented at the symposium to mark the African year of the environment in Ogun State,
330 Abeokuta, June 25, 1991.
- 331 Okojie, J.A. (1997): Forestry and the Environment. Paper presented at the Department of Forest
332 Resources Management, University of Ibadan. Symposium, 10th July, 1997.
- 333 Onumadu, F.N. (2002): Determinants of Adoption of Agroforestry Practices by Small Scale Farmers in
334 Katsina State, Nigeria. Unpublished P.h.D. Thesis, University of Ibadan 259 Pp.
- 335 Poulsen, G. (1978). *Man and tree in tropical Africa*. Ottawa IDRC 125 Pp.
- 336 Repetto, R. (1987). Population, Resources, Environment: An uncertain future. *Population Bulletin* 42(2):
337 3 - 9.
- 338 Swaminathan, M.S. (1987). The promise of agroforestry for ecological and nutritional security., In:
339 Stepler, H.A. and P.K.R. Nair (eds). *Agroforestry; A decade of development*. International
340 council for research in Agroforestry, Nairobi, Kenya Pp 15 - 18.
- 341 Umeh, A. (1989) Compulsory acquisition of land and compensation in Nigeria. Sweet and Maxwell,
342 London 73 Pp.
- 343
- 344
- 345
- 346
- 347
- 348