

Original Research Article

Growth, Carbon Assimilation and Quality of Kesum (*Persicaria Minor*) as Exposed to Zinc Oxide Nanoparticles

ABSTRACT

Aims: This study was conducted to investigate the effect of zinc oxide nanoparticles towards the *Persicaria minor* that can be used as a guidance for further toxicity investigation of ZnO-NPs.

Study design: A Completely Randomized Block Design (RCBD) was used with three replication. Each unit was consisted with eight plants and the total of 96 plants were used in this study.

Place and Duration of Study: This study was conducted in plot 1, Vegetables Field plot for Teaching and Research, Taman Pertanian Universiti, Universiti Putra Malaysia (UPM) Selangor, Malaysia, from May 2018 until August 2018.

Methodology: *Persicaria minor* were exposed to four different concentration of zinc oxide nanoparticles (ZnO-NPs) which were (50,100 and 150 mg/L) and 0 mg/L as a control. The ZnO-NPs was dissolved in distilled water before being applied to plants. 40 mL of ZnO-NPs solution was applied to each plant. The growth, carbon assimilation and also secondary metabolites were measured in this experiment.

Results: The results showed that the treatment of zinc oxide nanoparticles enhanced growth of the *Persicaria minor* as the plant treated with zinc oxide nanoparticles have higher plant height and total biomass when compared to control treatment. However, the analysis revealed that the treatment of zinc oxide nanoparticles highly and significantly influenced the carbon assimilation and quality of this plant as the treated plants showed reduction in chlorophyll content, photosynthesis rate, stomatal conductance and transpiration rate but increased in production of secondary metabolites. The increased in production of plant secondary metabolites may be attributed by the plant protection mechanism due to metabolic stress caused by high concentration of zinc oxide nanoparticles.

Conclusion: This research will progressively help in contributing some reliable and valid data on the effect of zinc oxide nanoparticles (ZnO-NPs), towards the *Persicaria minor* that can be used as guidance for further experimental investigation regarding this field.

Keywords: *Persicaria minor*, zinc- oxide nanoparticles, growth, carbon assimilation, secondary metabolites, toxicity

1. INTRODUCTION ()

According to National Health Portal [1], the World Health Organization (WHO) estimated that 80 percent of the world's population consuming herbal medicine for their health care and there are about 21000 plants species have potential to be utilized as medicinal plants. *Persicaria minor* is one of the plants that gained great attention in this field of study. According to Christopher et al. [2], *Persicaria minor* have gained great attention in scientific

Comment [d1]: Rewrite this section again in a more meaningful manner, first by identifying the need to carry out this study

23 study due to its high content of antioxidant. This plant possesses variety of pharmacological
24 properties such as antioxidant activity, antiulcer activity, anti-inflammatory activity,
25 antimicrobial activity, anticancer activity and can enhance the digestive properties and
26 cytotoxic activity [2]. According to Mikram et al. [3], *Polygonum minus* has been used
27 traditionally in herbal medicine to treat digestive disorder, remove dandruff and the essential
28 oil that extracted from *Persicaria minor* leaves is used as aroma therapy and also in perfume
29 industry.

Comment [d2]: Treid to include latest references after 2016 atleast.

30
31 Recently, nanoparticles (NPs) are widely studied because its beneficial properties in
32 agriculture and allied sector [4].According to them, zinc oxide nanoparticles (ZnO-NPs) is
33 one of NPs that used widely as it been utilized in variety of industrial sector including
34 medication, cosmetic materials, opposed microorganisms and textile industries. As it has
35 been commercially used, the toxicity effect of these ZnO-NPs to the environment and also
36 soil ecosystem are mainly concerned [5]. Sabir et al. [6] stated that ZnO-NPs possess
37 significant characteristics which have antimicrobial, optical and physical properties therefore
38 it have great potential to enhance agriculture. The presence of ZnO-NPs may enhance the
39 antioxidant mechanism that helps to stabilize the plants and improve the photosynthetic
40 efficiency [7]. However, the effect is depend on the concentration of ZnO-NPs and it is varies
41 from plant to plant [8].

Comment [d3]: Add more rfences to indicate the impotence of nano-partricle. If possible stress on its application in different fields

42
43 Secondary metabolites are the natural compound that produced by the plants and some of
44 the compound are utilized as medicines, flavoring and drugs. According to Biology
45 Reference [9], the simple classification of plants' secondary metabolites includes three main
46 groups which are terpenes, phenolics and nitrogen- containing compounds. Secondary
47 metabolites does not involve in the plant growth and development but required for plant to
48 survive in the environment because they give negatives impact on other organisms such as
49 pathogen and herbivores that can harm the plants [9]. Secondary metabolites possess
50 significant biological properties and also medicinal importance that can improve
51 pharmaceuticals field [10].

Comment [d4]: Don't use the word may when you are putting the references

52
53 There is no previous study being conducted on the effect of ZnO-NPs on physical and
54 biochemical response of *Persicaria minor* and also only few research being carried out to
55 discover about the effect of ZnO-NPs on carbon assimilation and production of secondary
56 metabolite of this plant, so this present study will reveal about the effect of ZnO-NPs towards
57 the *Persicaria minor* that can be the guidance to improve the field of agriculture in planting
58 this important medicinal plant and also other herb that being importantly used recently in
59 medicinal field and also be used as guidance for further toxicity investigation of ZnO-NPs.
60 Hence, the objectives of this study were to study the growth, carbon assimilation and quality
61 of *Persicaria minor* as affected by zinc oxide nanoparticles, to determine the optimum
62 concentration dose of zinc oxide nanoparticles that can enhance the optimum growth and
63 secondary metabolites of *Persicaria minor* and to recognize the relationship between
64 secondary metabolites and growth of *Persicaria minor* as exposed by zinc oxide
65 nanoparticles application.

Comment [d5]: Put more related rfrences and also explain the negative side of the metabolites if any????

Comment [d6]: Split this lines into two or more lines

67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117

2. MATERIAL AND METHODS

2.1 Experimental site

This study was conducted in plot 1, Vegetables Field plot for Teaching and Research, Taman Pertanian Universiti, Universiti Putra Malaysia (UPM) Selangor. The research site was set up with net shading and black plastic to reduce the absorption of water by sunlight since *Persicaria minor* require high amount of water and also to reduce the competition with grasses and other plants. This experiment was conducted from the month of May 2018 until August 2018.

Comment [d7]: Must explain different physico-chemical properties of the site of experiment...

2.2 Planting material

Persicaria minor was obtained from Jabatan Pertanian, Serdang. The first internodes at the bottom of the shoots with five number of leaves was cut in 5cm in height for the shoot stem cutting preparation. The shoot stem cuttings were immersed in the tap water overnight to increase their turgidity which increases the speed of the germination. Then, the propagation step were done in the trays with peat moss as the medium and the shoots stem cutting were left for two weeks for the development of the root. Then, the plants were transferred into standard polybag (16cm x 30cm) which was filled with top soil as the medium [11].

Comment [d8]: Also delineates the weather parameters during the course of investigation...

2.3 Soil preparation

The soil medium was obtained from Unit Herba, Taman Pertanian Universiti and top soil was used as the medium for the *Persicaria minor* planting. The top soil was transferred into the polybag until it occupied three- quarter of the polybag

2.4 Synthesis and properties of Zinc-Oxide Nanoparticles (ZnO-NPs)

The ZnO-NPs was synthesized chemically in the Laboratory of BioPhysics of Physics Department, Faculty of Science UPM by using sol gel method. ZnO-NPs is characterized as solid, white odorless powdery. In room temperature, 200ml of ethanol was added 0.2M of zinc acetate dehydrate. Then, the mixture was stirred for two hours to obtain clear solution. Then, 1.0 M sodium hydroxide (NaOH) was titrated into the mixture until the pH 9 is reach. The milky white slurry was obtained from the titration. To allow homogenous mixing, the white slurry was stirred for one hour more. The sample was left for 24 hours to allow the complete hydrolysis and gelation. The sample was separated from its solution. Then, the sample was filtrated to obtain white precipitate. The sample is dried in an oven for 48 hours at 100°C. The dried sample was grinded by mortar and pestle to yield ZnO powder.

Comment [d9]: Method is not very much clear, do rewrite it in a simple language as this is most important aspect of your study.

2.5 Experimental design

A Completely Randomized Block Design (RCBD) was used in this study with three replication. Each unit was consisted with eight plants and the totals of 96 plants were used in this study. After a month, *Persicaria minor* was exposed to four different concentration of zinc oxide nanoparticles (ZnO-NPs) which were (50,100 and 150 mg/L) and 0 mg/L as a control. The ZnO-NPs was dissolved in distilled water before being applied to plants. 40 mL of ZnO-NPs solution was applied to each plant.

118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152

153
154
155

156
157
158
159
160
161
162
163
164
165
166
167
168

2.6 Plant maintenance

The maintenance steps are very crucial to ensure the plant develops healthy and to avoid the plants from wilting or attacked by any disease that can cause the plants to die. At the early phase of cultivation, *Persicaria minor* were watered two times daily. The watering was unnecessary only when the heavy rain occur. This to avoid the over watering to the plants that can interfere the plants growth. The common insects that can interfere the plants growth were removed quickly from the planting area.

2.7 Collection of data

The growth data collection was conducted once a week after the application of treatment for the plants growth parameter. The destructive analysis and leaf gas exchange of the experiment were conducted at the end of the experiment.

2.7.1 Plant growth measurements

The plant growth measurements were conducted to obtain data about height, number of leaf and stem, diameter of stem, root to shoot ratio and the chlorophyll content.

2.7.1.1 Plant height

The plant height was measured starting from the stem on the soil surface until the highest shoot growth using measuring tape.

2.7.1.2 Plant basal diameter

The plant basal diameter was measured by using vernier caliper at the tips of the plants.

2.7.1.3 Plant leaves number

The leaves of the *Persicaria minor* were counted manually in every three weeks.

2.7.1.4 Chlorophyll content measurement

The total chlorophyll content of the leaves were measured by using chlorophyll meter (SPAD 502). The leaves of the plants in each treatment for each replication were clipped by chlorophyll meter clipper to obtain the reading.

2.7.1.5 Plant fresh weight measurement

The plants were removed first from the soil and all the dirt were removed under the flowing tap water. Then, the shoot and the root parts were separated for further analysis and all the plants parts were weighted separately using analytical balance.

2.7.1.6 Dry weight (biomass) measurement

The plants were dried in the oven at 60°C for 48 hours. Then, the measurements were recorded by using electronic weighing scale.

Comment [d10]: Delineates the different weather parameters during the course of investigation.....

169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221

2.7.1.7 Root to shoot ratio

The root to shoot ratio was determined by dividing the weight of the roots part to the shoot part after the oven dried process.

3.7.1.8 Plant leaf temperature determination

The Infrared (IR) thermometer was used to measure the plant leaves temperature.

2.7.2 Leaf gas exchange measurement

LI-6400XT (Li-COR Inc; Nebraska; USA) portable photosynthesis system was used to measure the leaf gas exchange. This equipment was warmed and was calibrated with ZERO IRGA mode for 30 minutes. The measurement was set at optimum condition which were 400 $\mu\text{mol mol}^{-1} \text{CO}_2$, 30°C cuvette temperature, 60% relative humidity with the rate of air flow set at 500 $\text{cm}^3 \text{min}^{-1}$ and then the cuvette condition was modified at 800 $\mu\text{molm}^{-2}\text{s}^{-1}$ photosynthetically photon flux density (PPFD). The measurement process of gas exchange was carried out between 9.00 am to 11.00 am by using fully expanded young leaves that give the measurement of net photosynthesis (A), stomata conductance (gs) and transpiration rate (E). Water use efficiency (WUE) was measured by using the formula of net photosynthesis dividing with transpiration rate. This is automatic operation and the results were saved in the the LI-6400XT console and Photosyn Assistant Software (Dundee Scientific, Dundee, UK) was used to analyze it. Precautions were taken to avoid mistakes during taking the measurements.

2.7.3 Total Phenolics and Flavonoids Quantification

Firstly, grounded plant tissue samples (0.1g) were extracted with 80% ethanol (10mL) on an orbital shaker for 120 minutes at 50°C. The mixture was filtered and the filtrate was used for the measurement of total phenolics and total flavonoids content. The total phenolic content in the leaves sample was measured by the Follin-Ciocalteu reagent (SigmaAldrich, Missouri, USA; diluted 10-fold). The absorbance was measured at 725 nm. The data were expressed as mg g^{-1} gallic acid equivalent (mg GAE g^{-1} dry sample). The determination process of total falvonoids content was measured by mixing a sample (1 mL) with NaNO_3 (Sigma Aldrich, Missouri, USA; 0.3 mL) in a test tube that covered with aluminium foil. The mixture then was left for 5 minutes. Then, 10% AlCl_3 (Wako Pure Chemical Industries Ltd; Tokyo, Japan; 0.3 mL) was added followed by the addition of 1.0 NaOH (Kanto Chemical Co. Inc.; Hokkaido, Japan; 2 mL). Then, the absorbance was measured at 510 nm using a spectrophotometer with rutting as a standard (data were expressed as mg g^{-1} rutting dry sample).

2.7.4 Chlorophyll fluorescence determination

The chlorophyll fluorometer was used to measure the chlorophyll florescence of the *Persicaria minor*. The mature leaf tissue was obtained from the *Persicaria minor* plant that cultivated at 20°C in glasshouse exposed with artificial light to give minimum photon flux density of 550 $\mu\text{mol m}^{-2} \text{S}^{-1}$ for 16 h photoperiod and phothosynthetically active radiation were supplied at 250 $\mu\text{mol m}^{-2} \text{S}^{-1}$ during 16 h photoperiod.

Comment [d11]: Wrote in running style why wrote as per thesis style looks like cut and paste from the thesis as such.....

Comment [d12]: Wrote some lines regarding infrared thermometer do quote its reference

Comment [d13]: No reference, write in a simple language as some budding scientists may consult it for their studies.....

Comment [d14]: Same as above

Comment [d15]: Same as above

222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274

2.8 Statistical Analysis

Statistical Package for Social Sciences (SPSS) version 24 was used to analyze the recorded data. A two-way ANOVA Test was conducted to analyze data for all the parameters used in the experiment. Results were significant if the p-value level ≤ 0.05 .

3. RESULTS AND DISCUSSION

3.1 Plant Height

Fig.1 depicted the effect of three months treatment of zinc oxide nanoparticles (ZnO-NPs) on the plant height of *Persicaria minor*. The result from analysis of variance showed that there was a significant effect between different concentration of zinc oxide nanoparticles treatment toward the height of *Persicaria minor* ($P=0.05$). From the figure, the increasing of concentration of zinc oxide nanoparticles treatment increased the plant height of *Persicaria minor*. The highest plant height was recorded in 100 mg/L on 12 weeks after treatment with mean 26.017 cm that might indicates the optimum concentration for the plant. Meanwhile, the control treatment recorded the shortest plant height with mean 22.083 cm.

The appropriate concentration of zinc oxide nanoparticles plays significant role in plant growth and promotion [12]. From the result, on twelfth week after harvesting, the plants treated with zinc oxide nanoparticles have higher plant height as compared to plants in control treatment. This finding indicates that the application of zinc oxide nanoparticles can induced the growth of the plants. Kouhi et al. [13] explained that the zinc oxide nanoparticles possess plant growth promoting effects and were used as micronutrient fertilizer as the presence of these nanoparticles triggered the physiological processes, acting as growth regulating compound that increased the plant growth such as the plant height and biomass. In addition, Prasad et al. [14] also supported that zinc oxide nanoparticles possess beneficial effects in enhancing plant growth and development. The presence of zinc can enhance the biochemical, physiological and anatomical responds of the plants thus increased the plant growth such as plant height and biomass [15]. Therefore, it can be concluded that the zinc oxide nanoparticles treatment induced the plant growth and 100 mg/L can be considered as the best concentration in promoting the height of *Persicaria minor* plant.




Fig.1. The effect of zinc oxide nanoparticles treatment (0, 50, 100 and 150) mg/L on plant height of *Persicaria minor*. Data are means with standard error of mean (SEM) of 24 replicates.

Comment [d16]: Well 0, 50 and 150 are same while only 100 shows some difference.....

Comment [d17]: What happened if only control and 100 con. Is there. I think scientist must select some more concentration.....

Comment [d18]:

Comment [d19R18]: Change this bar diagram into line graph


275
276
277
278
279

3.2 Plant Leaf Temperature

280 Fig. 2 highlighted the effect of three months treatment of zinc oxide nanoparticles (ZnO-NPs)
281 on the total leaf temperature of *Persicaria minor*. The result from the analysis variance
282 showed that there was a significant effect between different concentration of zinc oxide
283 nanoparticles treatment toward the total leaf temperature of *Persicaria minor* ($P=0.05$).
284 Based on the figure, the trend shows that the plant leaf temperature increased linearly with
285 the concentration of zinc oxide nanoparticles treatment.
286

287 The plants maintaining their most important physiological process (photosynthesis) by
288 maintaining their average leaf temperature at around 21 degree celsius [16]. The plants leaf
289 temperature depends on the stomatal conductance and transpiration rates of the plants [17].
290 Transpiration is one of the best mechanism used by plants to cool themselves by 'pumping'
291 out water from leaves through stomata [18]. From this study, the increasing of plant leaves
292 temperature can be explained through the reduction of the stomatal conductance and
293 transpiration rates of the plants due to the increasing the concentration of the treatment. This
294 high temperature in turns will give negative effect to the photosynthesis process thus affect
295 the plant yields. Therefore, it can be concluded that the zinc oxide nanoparticles treatment
296 increased the plant leaf temperature due to the reduction of stomatal conductance and also
297 transpiration rate of *Persicaria minor*.
298

299
300
301
302
303
304
305
306
307
308
309
310
311
312
313



314 **Fig.2.**The effect of zinc oxide nanoparticles treatment (0, 50, 100 and 150) mg/L on
315 plant leaf temperature of *Persicaria minor*. Data are means with standard error of
316 mean (SEM) of 24 replicates.
317

3.3 Total Biomass

320 Fig.3 shows the effect of three months treatment of zinc oxide nanoparticles (ZnO-NPs) on
321 the total biomass of *Persicaria minor*. The result from the analysis variance showed that
322 there was a significant effect between different concentration of zinc oxide nanoparticles
323 treatment toward the total biomass of *Persicaria minor* ($P=0.05$). From the result, it showed
324 that higher concentration of zinc oxide nanoparticles increased the total biomass of
325 *Persicaria minor* as plants treated with zinc oxide nanoparticles treatment have higher total
326 biomass as compared to the control treatment and 100 mg/L have the highest value of total
327 biomass with mean 2.675g.

Comment [d20]: If treatment raises the temperature then why control mplots has higher temperature than 50. Convert this fig to line graph.

328
329
330
331
332
333
334
335
336
337
338
339
340
341

The highest value of total biomass indicates that 100 mg/L was the optimum concentration of zinc oxide nanoparticles for *Persicaria minor*. This study showed that the treatment of zinc oxide nanoparticles increased the plant biomass so the treatment might be effective in boosting the plant growth and yield. Similar finding was observed from study conducted by Venkatachalam et al. [19] that revealed total biomass significantly increased in the zinc oxide nanoparticles treated plants as compared to control. This is supported by Munir et al. [20] that stated the treatment of zinc oxide nanoparticles increased the shoot and root dry weight thus increased the plant biomass. The presence of zinc can enhanced the biochemical, physiological and anatomical responds of the plants thus increased the plant growth such as plant height and biomass [15]. Hence, it can be concluded that the presence of zinc oxide nanoparticles can boost the *Persicaria minor* growth resulting in increasing of the plant total biomass.

342
343
344
345
346
347
348
349
350
351
352
353
354
355
356




Fig.3.The effect of zinc oxide nanoparticles treatment (0, 50, 100 and 150) mg/L on total biomass of *Persicaria minor*. Data are means with standard error of mean (SEM) of 24 replicates.

361
362
363
364
365
366
367
368
369
370
371

3.4 Net photosynthesis rate (A)

Fig.4 illustrated the effect of three months treatment of zinc oxide nanoparticles (ZnO-NPs) on the net photosynthesis rate of *Persicaria minor*. The exposure of zinc oxide nanoparticles toward the *Persicaria minor* was highly and significantly affected the net photosynthesis rate of the plant ($P=0.05$). From the result, the highest photosynthesis rate was recorded in control treatment while the lowest photosynthesis rate was recorded in 150 mg/L of zinc oxide nanoparticles treatment with mean 6.32 and 4 respectively. From the figure, the photosynthesis rate of the plant reduced with the increasing concentration of zinc oxide nanoparticles treatment.


372
373
374
375
376
377
378
379
380

Photosynthesis is the perfect measurement to access plant performance. From this study, the net photosynthesis of *Persicaria minor* was reduced with the increasing concentration of zinc oxide nanoparticles treatment. Photosynthesis is highly affected in plants that exposed to excess heavy metal where higher level of zinc oxide nanoparticles inhibits the photosynthetic apparatus and caused critical changes to chlorophyll structure and amount [21]. This finding is also similar with Wang et al. [22] that revealed the presence of zinc oxide nanoparticles reduced the chlorophyll content in leaves thus reduced the photosynthetic efficiency in plants. In addition, plants exposed to high concentration of zinc oxide nanoparticles have low photosynthetic efficiency due to the reduction of chlorophyll content

Comment [d21]: Must indicate the clear dose with proper reasoning. And why it is on lower side when treated with 150 mg/L

Comment [d22]: Change it to line graph

381 and also damaged to the photochemical system. Therefore, it can be concluded that the
382 presence of zinc oxide nanoparticles reduced the photosynthetic efficiency of *Persicaria*
383 *minor* plants.
384



Comment [d23]: Explain your result with reference to each and every treatment.....

385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402 **Fig.4: The effect of zinc oxide nanoparticles treatment (0, 50, 100 and 150) mg/L net**
403 **photosynthesis rate of *Persicaria minor*. Data are means with standard error of mean**
404 **(SEM) of 24 replicates.**

405 3.5 Transpiration rate (E)

406
407
408 Fig.5 depicted the effect of three months treatment of zinc oxide nanoparticles (ZnO-NPs) on
409 the transpiration rate of *Persicaria minor*. The exposure of different concentration of zinc
410 oxide nanoparticles toward the *Persicaria minor* was highly and significantly affected the
411 transpiration rate of the plant ($P=0.05$). From the figure, the transpiration rate of the plant
412 reduced with the increasing concentration of zinc oxide nanoparticles treatment. The
413 transpiration rate of 50 mg/L treatment was significantly higher with mean 2.342 while the
414 lowest transpiration rate of *Persicaria minor* was observed in 150 mg/L of zinc oxide
415 nanoparticles treatment with mean 1.42.
416

Comment [d24]: Why is it so? Provide reasoning/.....


Comment [d25]: Water vapors from the stomata.....

417 Transpiration is a process of the movement of water through plant and this process mainly
418 take place in leaves. Transpiration process is mainly controlled by the opening and closing of
419 the stomata. From the result, the transpiration rate reduced with the increasing concentration
420 of zinc oxide nanoparticles treatment. This finding is similar with Xiaoping et al. [23] that
421 revealed both transpiration rate and stomatal conductance reduced in plants treated with
422 zinc oxide nanoparticles. The stomatal closure reduced the transpiration rate of the plants.
423 According to Vankova et al. [24], the presence of zinc oxide nanoparticles induced the
424 production of plant stress hormone, abscisic acid and this hormone mainly accumulated in
425 leaves. The higher level of abscisic acid triggered the stomatal closure which in turn reduced
426 the transpiration rate in plants. Hence, the application of zinc oxide nanoparticles reduced
427 transpiration rate of *Persicaria minor* due to the high level of stress hormone that cause the
428 closure of stomata.
429

Comment [d26]: Meaning not clear, rewrite these lines

430
431
432
433

434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486



Comment [d27]: Convert this figure into the line graph

Fig.5. The effect of zinc oxide nanoparticles treatment (0, 50, 100 and 150) mg/L transpiration rate of *Persicaria minor*. Data are means with standard error of mean (SEM) of 24 replicates.

3.6 Stomatal conductance (gs)


Fig.6 shows the effect of three months treatment of zinc oxide nanoparticles (ZnO-NPs) on the transpiration rate of *Persicaria minor*. The exposure of different concentration of zinc oxide nanoparticles toward the *Polygonum minus* was highly and significantly affected the stomatal conductance of the plant ($P=0.05$). From the figure, the stomatal conductance of the plant reduced as the concentration of the zinc oxide nanoparticles treatment increasing. The highest stomatal conductance was recorded in control treatment while the lowest stomatal conductance was observed in 150 mg/L of zinc oxide nanoparticles treatment with mean 0.428 and 0.07 respectively.

Comment [d28]: Why? Provide reasoning>>

Stomatal conductance is a measure of the degree of stomatal opening and a good indicator in accessing plant water status [25]. The finding showed reduction in stomatal conductance due to the increasing of zinc oxide nanoparticles concentration is similar with Xiaoping et al. [23] that proved the higher concentration of zinc oxide nanoparticles reduced the stomatal conductance resulting in low photosynthetic efficiency of the plants. Singh and Bhati [26] also stated that high amounts of zinc oxide nanoparticles can restrict the stomatal conductance. This might due to the toxicity of the treatment disturbed the cell mechanism thus alters the stomatal function. Tsonev and Lidon [21] explained that the stomatal response to high concentration of zinc oxide nanoparticles is related to the changes in carbonic anhydrase (CA) activity. Carbonic anhydrase is enzyme that responsible for the stomatal activity and the presence of zinc oxide nanoparticles influenced the CA activity that triggered the stomatal closure thus reduced the stomatal conductance of the plants. Hence, it can be concluded that the presence of zinc oxide nanoparticles alter the stomatal mechanism thus reducing the stomatal conductance of *Persicaria minor* plants.

Comment [d29]: Put introductory lines in the strat of the discussion

487



Comment [d30]: Change to line graph

488
489

Fig.6. The effect of zinc oxide nanoparticles treatment (0, 50, 100 and 150) mg/L on stomatal conductance of *Persicaria minor*. Data are means with standard error of mean (SEM) of 24 replicates.

491

492

493

494

3.7 Maximum efficiency of photosystem II

495

496

497

498

499

500

501

502

503

504

505

Fig.7 depicted the effect of three months treatment of zinc oxide nanoparticles (ZnO-NPs) on the maximum efficiency of photosystem II of *Persicaria minor*. The exposure of different concentration of zinc oxide nanoparticles toward the *Persicaria minor* was highly and significantly affected the maximum efficiency of photosystem II of the plant ($P=0.05$). Based on the figure, increasing the concentration of zinc oxide nanoparticles caused the maximum efficiency of photosystem II to decrease. The highest value of maximum efficiency of photosystem II was observed in control treatment while the lowest value of maximum efficiency of photosystem II was recorded in 150 mg/L treatment with mean 0.758 and 0.522 respectively.

Comment [d31]: Why???????

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524


525

526

The phytotoxicity of zinc oxide nanoparticles can be accessed through the efficiency of photosynthetic mechanism (chlorophyll florescence) that act as indicator in phytotoxicity assays. The finding of this study revealed that increasing the zinc oxide nanoparticles concentration resulting in lower maximum efficiency of photosystem II of *Polygonum minus*. Wang et al. [22] stated that the treatment of zinc oxide nanoparticles reduced the chlorophyll florescence parameter and damaged the photochemical system. This finding can be explained further that the presence of zinc oxide nanoparticles induced the oxidative stress in plants and increase the production of reactive oxygen species (ROS) which alter the gene expression pathway thus reduced the chlorophyll florescence in plants. Therefore, it can be concluded that the zinc oxide nanoparticles treatment reduced the chlorophyll florescence parameters of *Persicaria minor* plants.

Comment [d32]: Brought thes line in the start

527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579



Comment [d33]: Change it too

Fig.7. The effect of zinc oxide nanoparticles treatment (0, 50, 100 and 150) mg/L on maximum efficiency of photosystem II of *Persicaria minor*. Data are means with standard error of mean (SEM) of 24 replicates.

3.8 Maximum yield of photosystem II

Fig.8 shows the effect of three months treatment of zinc oxide nanoparticles (ZnO-NPs) on the maximum yield of photosystem II of *Persicaria minor*. The exposure of different concentration of zinc oxide nanoparticles toward the *Persicaria minor* was highly and significantly affected the maximum yield of photosystem II of the plant ($P=0.05$). The trend shows that increasing the concentration of zinc oxide nanoparticles caused the maximum yield of photosystem II to decrease. Based on the figure, 50 mg/L of zinc oxide nanoparticles treatment shows the highest value of maximum yield of photosystem II with mean 3.166 when compared with other treatment.

Comment [d34]: Why???????

The photosynthetic pigments (chlorophyll fluorescence) act as indicator in phytotoxicity assays in accessing the phytotoxicity of zinc oxide nanoparticles towards the plant. From this study, it was observed that the treatment of zinc oxide nanoparticles reduced the maximum efficiency of photosystem II which in turn reduced the maximum yield of photosystem II of *Persicaria minor*. According to Tsonev and Lidon [21], inside the chloroplast lamellae, the presence of zinc oxide nanoparticles caused the inhibition of photosynthetic electron transport and implicates the water evolving complex of photosystem II thus inhibits the photolysis and oxygen emission that disturb the conformation of photosystem II core complex. This mechanism explained how the zinc oxide nanoparticles treatment reduced the efficiency and yield of photosystem II in plants. Hence, the treatment of zinc oxide nanoparticles reduced the maximum yield and efficiency of photosystem II which in turn disturbed the photosynthetic process of *Persicaria minor* plant.

Comment [d35]: Introductory lines

580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632




Fig.8. The effect of zinc oxide nanoparticles treatment (0, 50, 100 and 150) mg/L on maximum yield of photosystem II of *Persicaria minor*. Data are means with standard error of mean (SEM) of 24 replicates.

3.9 Minimal fluorescence

Fig.9 shows the effect of three months treatment of zinc oxide nanoparticles (ZnO-NPs) on the minimal fluorescence of *Persicaria minor*. The exposure of different concentration of zinc oxide nanoparticles toward the *Polygonum minus* was highly and significantly affected the maximum yield of photosystem II of the plant ($P=0.05$). The result showed that higher concentration of zinc oxide nanoparticles treatment resulted in higher value of minimal fluorescence of *Persicaria minor*. The highest value of minimal inflorescence was recorded in 150 mg/L with mean 627.2 while the lowest value was observed in 50 mg/L with mean 462.6.

Comment [d36]: Why?????

Higher minimal fluorescence indicates higher heat dissipation of plants. This might due to the presence of zinc oxide nanoparticles that induced stress in plants thus caused plants to produce high amount of heat. From this study, the treatment of zinc oxide nanoparticles reduced the transpiration rate of *Persicaria minor*. This reduction might related with the increasing of minimal fluorescence of the plants. The high minimal fluorescence can cause heat stress to the plants. Heat stress is defined as the increase temperature beyond the threshold level that cause damage to plant growth and development [27]. Therefore, it can be deduced that the treatment of zinc oxide nanoparticles increased the minimal fluorescence of *Persicaria minor* due to the reduction in transpiration rate of the plants.

633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685




Fig.9. The effect of zinc oxide nanoparticles treatment (0, 50, 100 and 150) mg/L on minimal fluorescence of *Persicaria minor*. Data are means with standard error of mean (SEM) of 24 replicates.


3.10 Performance index (PI)

Fig.10 indicated the effect of three months treatment of zinc oxide nanoparticles (ZnO-NPs) on the performance index of *Persicaria minor*. The exposure of different concentration of zinc oxide nanoparticles toward the *Polygonum minus* was highly and significantly affected the performance index of the plant ($P=0.05$). Based on the figure, the highest and lowest performance index were observed in control treatment and 150 mg/L of zinc oxide nanoparticles treatment with mean 2.422 and 1.504 respectively. The performance index of *Persicaria minor* reduced when treated with higher concentration of zinc oxide nanoparticles that indicates zinc oxide nanoparticles increased the plant stress.

Comment [d37]: Why???????

Nanoparticles such as zinc oxide and silver were located on the surface of plants cells and induced the oxidative stress to the cells by the activation of oxidative stress signaling [28]. From this study, it was observed that the treatment of zinc oxide nanoparticles reduced the plants performance index. Zahed et al. [28] stated that the generation of reactive oxygen species (ROS) due to the zinc oxide nanoparticles treatment alter the gene expression and cell mechanism which in turn reduced the performance index of the plants. Wang et al. [22] explained that the toxicity of zinc oxide nanoparticles reduced chlorophyll content plants, resulted in low photosynthesis efficiency thus reduced the plants performance. Hence, it can be concluded that the presence of zinc oxide nanoparticles induced stress in *Persicaria minor* resulting in low performance index of the plants.

686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738



Comment [d38]: Covert to line graph

Fig.10. The effect of zinc oxide nanoparticles treatment (0, 50, 100 and 150) mg/L on performance index of *Persicaria minor*. Data are means with standard error of mean (SEM) of 24 replicates.

3.11 Total phenolics content

Fig.11 shows the effect of three months treatment of zinc oxide nanoparticles (ZnO-NPs) on the total phenolics production of *Persicaria minor*. The exposure of different concentration of zinc oxide nanoparticles toward the *Polygonum minus* was highly and significantly affected the total phenolics production of the plant ($P=0.05$). Based on the figure, the total phenolics production of the plant was directly proportional with the concentration of zinc oxide nanoparticles treatment. The lowest total phenolics production was recorded in control treatment while the highest total phenolics production was recorded in 150 mg/L treatment with mean 1.444 and 3.82 respectively. This result indicates that zinc oxide nanoparticles treatment induced stress and increased the secondary metabolites production of *Persicaria minor*.

Phenolics are compound that produced by plants to protect plants against stress. These compound play significant role in plant development (lignin and pigment biosynthesis) and also provided structural integrity for plant's support [29]. From this study, the greater production of total phenolics content in *Polygonum minus* with the increasing of concentration treatment revealed that the presence of zinc oxide nanoparticles induced stress towards the plants. This finding is supported by Rastogi et al. [30] that stated zinc oxide nanoparticles treatment induced the Reactive Oxygen Species (ROS) production in plants thus increased plants stress. They also stated that higher concentration of zinc oxide nanoparticles lead to the damage of plant cell wall and plasma membrane thus induced the production of plant secondary metabolites for plants defense against disease and threat. Therefore, it can be concluded that higher concentration of zinc oxide nanoparticles lead to plant stress and boost the plants secondary metabolites production which in turn can enhance the defense response of *Persicaria minor* plants.

Comment [d39]: Rewrite and put importance lines to initial paragraph.

739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791




Fig.11. The effect of zinc oxide nanoparticles treatment (0, 50, 100 and 150) mg/L on total phenolics production of *Persicaria minor*. Data are means with standard error of mean (SEM) of 24 replicates

3.12 Total flavonoids content

Fig. 12 shows the effect of three months treatment of zinc oxide nanoparticles (ZnO-NPs) on the total flavonoids production of *Persicaria minor*. The exposure of different concentration of zinc oxide nanoparticles toward the *Persicaria minor* was highly and significantly affected the total flavonoids production of the plant ($P=0.05$). Based on the figure, the total flavonoids production of the plant was directly proportional with the concentration of zinc oxide nanoparticles treatment. The lowest total flavonoids production was recorded in control treatment while the highest total flavonoids production was recorded in 150 mg/L treatment with mean 0.43 and 1.478 respectively. This result indicates that zinc oxide nanoparticles treatment induced stress and increased the secondary metabolites production of *Persicaria minor*.

Flavonoids are a wide group of plants chemicals (phytonutrients) that found mostly in fruits and vegetables. Flavonoids plays significant role in pharmacological field since this compound is a good source of antioxidant and anti-inflammatory, protect skin, enhanced brain function and also good for blood pressure regulation [31]. From this study, the greater production of total flavonoids content in *Polygonum minus* with the increasing of concentration treatment revealed that the presence of zinc oxide nanoparticles induced stress towards the plants. This finding is similar with Zafar et al. [32] the higher treatment concentration of zinc oxide nanoparticles generates oxidative stress of plants thus increasing the plant secondary metabolites production to protect plants against stress. The initial response of plants towards the presence of nanoparticles involved the increasing level of reactive oxygen species (ROS), cytoplasmic Ca^{2+} and up regulation of nitrogen activated protein kinase (MAPK) cascades thus activates the plants secondary metabolites that act against stress to protect the plants [33]. In addition, the presence of zinc oxide nanoparticles enhanced the expression of genes related to antioxidant capacity thus boost the defense mechanism of the plants by enhancing the production of plants secondary metabolites. Hence, it can be concluded that the presence of zinc oxide nanoparticles enhanced the *Persicaria minor* secondary metabolites production by increasing the total flavonoids production of the plants.

Comment [d40]: Rewrite discussion.....

792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844




Fig.12. The effect of zinc oxide nanoparticles treatment (0, 50, 100 and 150) mg/L on total flavonoids production of *Persicaria minor*. Data are means with standard error of mean (SEM) of 24 replicates.

4. CONCLUSION

Overall, the treatment of zinc oxide nanoparticles increased the growth parameter of the plants as the treated plants showed higher value of plant height and total biomass when compared to plants in control treatment. However, the treatment of zinc oxide nanoparticles reduced the photosynthesis rate, transpiration rate and stomatal conductance thus reduced the performance index of *Persicaria minor* plants. The treatment also might induced the plants stress as it was significantly observed that the production of secondary metabolites (total phenolics and flavonoids production) were directly proportional with the treatment concentration that used mainly for plants production against stress. From this study, it can be concluded that the optimum concentration of zinc oxide nanoparticles for enhancing the *Persicaria minor* growth was 100 mg/L because it recorded the highest value for most of the plants growth parameters.

5. REFERENCES

1. National Health Portal. Introduction and Importance of Medicinal Plants and Herbs. 2016. Accessed 8 July 2018. Available: https://www.nhp.gov.in/introduction-and-importance-of-medicinal-plants-and-herbs_mtl.
2. Christopher PV, Parasuraman S, Christina JMA, Asmawi MZ, Vikneswaran M. Review on Polygonum minus. Huds, a commonly used food additive in Southeast Asia. Pharmacognosy Research. 2015; 7(1): 1–6.
3. Vikram P, Chiruvella KK, Husna IAR, Mohammed A. A recent review on phytochemical constituents and medicinal properties of kesum (Polygonum minus Huds.). Asian Pacific Journal of Tropical Biomedicine. 2014; 4(6): 430–435.
4. Mirzaei H, Darroudi M. Zinc oxide nanoparticles: Biological synthesis and biomedical applications. Journal of Ceramics International. 2016; 43(1): 907-914. doi: <https://doi.org/10.1016/j.ceramint.2016.10.051>.
5. Vishnu DR, Minkina T, Yaning C, Sushkova S, Chaplugin VA, Mandzhieva S. A review on salinity adaptation mechanism and characteristics of Populus euphratica, a boon for arid ecosystems. Acta Ecológica Sinica. 2016; 36(6): 497-503.
6. Sabir S, Arshad M, Sunbal KC. Zinc Oxide nanoparticles for revolutionizing agriculture: synthesis and applications. Scientific World Journal. 2014; 8.

Comment [d41]: Convert this to line graph and if possible took the into power point and then prepare one plate from the four figure. In the present form it is very difficult for us to adjust so much of the figures.....

Comment [d42]: Conclusion needs to be rewritten based on the results reported.....

- 845 Doi:<https://dx.doi.org/10.1155%2F2014%2F925494>
846 7. Faizan M, Faraz A, Yusof M, Khan M, Hayat S. Zinc oxide nanoparticle-mediated
847 changes in photosynthetic efficiency and antioxidant system of tomato plants. *Journal of*
848 *Springer*. 2017; 56(2): 678-686. doi <https://doi.org/10.1007/s11099-017-0717-0>
849 8. Manzer HS, Waibi MH, Firoz M, Mutahhar YAK. Role of nanoparticles in plants. 2015.
850 Accessed 18 February 2018. Available: [file:///C:/Users/User/Downloads/9783319145013-](file:///C:/Users/User/Downloads/9783319145013-c2.pdf)
851 [c2.pdf](file:///C:/Users/User/Downloads/9783319145013-c2.pdf).
852 9. Biology Reference. Secondary Metabolites in Plants. 2018. Accessed 13 July 2018.
853 Available:[http://www.biologyreference.com/Re-Se/Secondary-Metabolites-in Plants.html](http://www.biologyreference.com/Re-Se/Secondary-Metabolites-in%20Plants.html).
854 10. Schater H, Wink M. Medicinally important secondary metabolites in recombinant
855 microorganisms or plants: progress in alkaloid biosynthesis. *Journal of Biotechnol.*
856 2009; 4(12): 1684-1703. doi: 10.1002/biot.200900229.
857 11. Ibrahim MH, Nurul NM, Zain NAM. Growth, carbon assimilation and biochemical changes
858 of *Polygonum minus* Huds. as affected by nitrogen fertilization. *Journal of Annual*
859 *Research and Review in Biology*. 2018; 26(1): 1-17.
860 12. Paripark R, Sutichai S, Sutee C. ZnO nanoparticles affect differently the morphological
861 and physiological responses of riceberry plants (*Oryza sativa* L.). *SNRU Journal of*
862 *Science and Technology*. 2018; 10(1): 75-81.
863 13. Kouhi SMM, Lahouti M, Ganjeali A, Entezari MH. Comparative phytotoxicity of ZnO
864 nanoparticles, ZnO microparticles and Zn²⁺ on rapeseed (*Brassica napus* L.):
865 investigating a wide range of concentration. *Toxicology and Environmental Chemistry*.
866 2014; 96(6): 861-868.
867 14. Prasad TNVKV, Sudhakar Y, Sreenivasulu P, Latha V, Munaswamy K, Raja RJS,
868 Sreepasad PR, Sajanlal Pradeep T. Effect of nanoscale zinc oxide nanoparticles on
869 germination, growth and yield of peanut. *Journal of Plant Nutrition*. 2012; 35: 905-927.
870 15. Vishnu DR, Tatiana MM, Arvind B, Svetlana NS, Saglar M, Ritu S, Andrey G, Viktoriia
871 ST, William OP, Karen AG, Hamshik SM. Effects of zinc-oxide nanoparticles on soil,
872 plants, animals and soil organisms: a review. *Environmental Nanotechnology, Monitoring*
873 *and Management*. 2018; 9: 76-84.
874 16. Ledford H. Leaves keep their cool. 2008. Accessed 13 November 2018. Available:
875 <https://www.nature.com/news/2008/080611/full/news.2008.884.html>.
876 17. Urban J, Miles WJ, McGuire MA, Teskey RO. Increase in leaf temperature opens stomata
877 and decouples net photosynthesis from stomatal conductance in *Pinus taeda* and
878 *Populus deltoides* x *nigra*. *Journal of Experimental Botany*. 2017; 68(7): 1757-1767.
879 18. Harlequin G. Plants in high temperature. 2016. Accessed 13 November 2018. Available:
880 <http://www.harlequinsgardens.com/mikls-articles/plants-in-high-temperatures/>.
881 19. Venkatachalam P, Priyanka N, Manikandan K, Ganeshbaru I, Indiraarulsevi P, Geetha
882 N, Muralikrishna K, Bhattacharya RC, Tiwari M, Sharma N, Sahi SV. Enhanced plant
883 growth promoting role of phycocyanin coated zinc oxide nanoparticles with P
884 supplementation in cotton (*Gossypium hirsutum* L.). *Plant Physiology and*
885 *Biochemistry*. 2017; 110: 118-127.
886 20. Munir T, Rizwan M, Kashif M, Shahzad A, Ali S, Amin N, Zahid R, Alam MFE, Imran M.
887 Effect of zinc oxide nanoparticles on the growth and zinc uptake in wheat
888 (*Triticum aestivum* L.) by seed priming method. *Digest Journal of Nanomaterials and*
889 *Biostructures*. 2018; 13(1): 315-32.
890 21. Tsonev T, Lidon FJC. Zinc in plants- an overview. *American Journal Food Agriculture*.
891 2012; 24(4): 322-333.
892 22. Wang XP, Li QQ, Pei ZM, Wang SC. Effects of zinc oxide nanoparticles on the growth,
893 photosynthetic traits and antioxidative enzymes in tomato plants. *Biological Plantarum*.
894 2018; 1-8.
895 23. Xiaoping W, Chen S, Xiyu Y, Shucai W. Zinc oxide nanoparticles affect the biomass
896 accumulation and photosynthesis in *Arabidopsis*. *Frontiers Plant Science*. 2016). doi
897 10.3389/fpls.2015.01243

- 898 24. Vankova R, Lamda P, Podlipna R, Dobrey PT, Prerostova S, Langjansova L, Gaudinova
899 A, Motkova K, Knirsch V, Vanek T. ZnO nanoparticles effects on hormonal pools in
900 Arabidopsis Thaliana. *Science Total Environment*. 2017; 593-594.
- 901 25. Gimenez C, Gallardo M, Thompson RB. Plant-water relations. *Earth System and*
902 *Environmental Science*. 2013; 231-238.
- 903 26. Singh G, Bhati M. Mineral toxicity and physiological functions in tree seedling irrigated
904 with effluents of varying chemistry in sandy soil of dry region. *Journal of Environmental*
905 *Science*. 2003; 21(1): 45-63.
- 906 27. Nahla SH, Taher ASED, Mohamed HH, Ibrahim HB, Asmaa AM. Magnetic and zinc oxide
907 nanoparticles alleviated heat stress in wheat plants. *Current Nanomaterials*. 2018; 3(1):
908 32-43.
- 909 28. Zahed H, Ghazala M, Komatsu S. Plant's responses to nanoparticles stress. *International*
910 *Journal of Molecular Science*. 2015; 16(11): 26644-26653.
- 911 29. Bhattacharya A, Sood P, Citovsky V. The roles of plant phenolics in defense and
912 communication during Agrobacterium and Rhizobium infection. *Molecular Plant*
913 *Pathology*. 2010; 11(5): 705-719.
- 914 30. Rastogi A, Zivcak M, Sytar O, Kalaji HM, Xiaolan H, Mbarki S, Brestic M. Impact of metal
915 and metal oxide nanoparticles on plant: a critical review. *Frontiers in Chemistry*. 2017; 5:
916 78.
- 917 31. Szalay S. What are flavonoids?. 2015. Accessed 13 November 2018.
918 <https://www.livescience.com/52524-flavonoids.html>.
- 919 32. Zafar H, Ali A, Ali JS, Haq IU, Zia M. Effect of ZnO nanoparticles on Brassica seedlings
920 and stem explants: growth dynamics and antioxidant response. *Frontiers Plants Science*.
921 2016; 7: 535.
- 922 33. Marslin G, Sheeba CJ, Franklin G. Nanoparticles alter secondary metabolism in plants
923 via ROS burst. *Frontiers in Plant Science*. 2017; 8: 832.
- 924