

Original Research Article

1

2

Effect of silicon and phosphorus additions and their interactions on wheat plants grown on a clay soil

4

5

Abstract

7 A pot experiment was conducted in clay soil collected from Agricultural Research Center farm, Giza governorate, Egypt. Wheat grains (*Triticumaestivum* L., Giza 168) were cultivated to study the effect of silicate and phosphate ions as well as their interactions on the growth and nutritional status of the growing plants, beside their availability in the studied soil. Silicon (Si) in the form of sodium metasilicate penta-hydrate ($\text{Na}_2\text{SiO}_3 \cdot 5\text{H}_2\text{O}$) was added at a rate of 0, 200, 300 and 400 mg Si kg^{-1} soil, and phosphorus (P) in the form of calcium super phosphate was given at a rate of 0, 3.5, 7.0, 10.0 and 13.0 mg P kg^{-1} soil to represent 0, 25, 50, 75 and 100% of the recommended rate of P fertilization by the Egyptian Ministry of Agriculture for wheat cultivation. Also, the experiment included combinations between all these concentrations of Si and P. Obtained results showed that Si and P availability increased in the studied soil with increasing either Si or P concentrations added. This means that P availability in soil as an essential element for plant growth can be improved by addition of Si. Also Si increased in plant with increasing applied Si concentrations. Interaction between Si and P generally increased all parameters of plant growth; such responses were significant for fresh and dry weights of wheat plants at booting stage. It could be recommended that selecting good P fertilization design, including time and rate of addition, goes along with values of available Si in the soil.

Keywords: Silicon, Phosphorus, Interactions, Wheat plant, Clay soil.

25

Introduction

26 Silicon (Si) is a basic mineral formatting element, which is considered the second most abundant element in the earth's crusts (Tubana et al., 2016). It is known as non-essential element for higher plants, while lately it confirmed its role in enhancing crop yield by promoting several desirable plant physiological processes. Silicon fertilization has been reported to result in increased soil exchange capacity, improved soil aggregation; consequently improved water and air regimes, transformation of P-containing minerals and

Delete the comma :[1G]Comment

formation of aluminosilicates and heavy metal silicates (Matichenkov and Bocharnikova, 2003). The amount of literature documenting the benefits of Si on plants is vast and primarily highlights the value of Si fertilization in maintaining plant productivity under stressed conditions (Epstein, 1999; Li et al., 2007). The established Si-induced mechanisms to improve plants' resistance to biotic and abiotic stresses take place in the soil, the root system, and inside the plant. Silicon deposition of silica in shoots and leaf epidermis, also known as the mechanical barrier hypothesis (Rodrigues et al., 2015), enhances the plant's mechanical strength and protective layer. This barrier is believed to help promote the plant's resistance to pathogens, improve plants' tolerance to abiotic stresses and increase photosynthetic rates (Danoff et al., 2009; Rizwan et al., 2012).

42 Phosphorus (P) is a macro-nutrient that has the most influence on plant growth and productivity, and, according to statistics of Egyptian Ministry of Agriculture, more than 50% of the cultivated areas are in P-deficient soils (Academy of Scientific Research and Technology, 2017). The low availability of P in those soils is one of the greatest challenges in the fertilization of those soils. There are various factors that interfere with the availability of P in the soil, including the environmental factors that control the activity of microorganisms which immobilize or liberate the orthophosphate ions, the physiochemical and mineralogical properties of the soil (Santos et al., 2008) and the silicon-phosphorus interactions (Carvalho et al., 2001; Tokura et al., 2011). One of the most effective solutions to raise the P availability is to exploit the relationship between it and Si concentration in the soil. Ma and Takahashi (1992) studied the effect of Si on the growth and P uptake of rice and found that addition of Si raised the optimum P level for rice and enhanced plant growth and yield. Alovisei et al. (2014) studied the Si-P interactions in some soils cultivated with bean plants in Brazil and found that application of silicate could result in the increase of P availability to plants, as based on the fact that the silicate anion occupies the adsorption sites of the phosphate anion, which reflect on enhancing plant growth and yield and increasing the content of its different parts with P.

58 Wheat (*Triticum aestivum* L.) is the most important cereal crop in Egypt where it provides more than 30% of the population's calorie intake (Academy of Scientific Research and Technology, 2017). Works related to the use of Si to enhance P fertilization efficiency and uptake in wheat plant are rare in Egypt to date. Thus, the main objective of this investigation is to study the effect of silicate and phosphate ions as well as their interactions on the nutritional status of growing wheat plants, beside their availability in the studied clay soil.

64

65

67 A pot experiment was conducted in clay soil classified as *Entisol*, collected from Agricultural Research Center farm, Giza governorate, Egypt (29° 58' 55.046" N and 31° 12' 49.292" E). Soil samples (0-30 cm) were air-dried, crushed and finely ground, then sieved to pass through a 2 mm sieve. Some physical and chemical characteristics of the studied soil samples were achieved as described by Klute (1986) and Page et al. (1982) and the results are presented in Table 1. The study was conducted by growing wheat grains (*Triticum aestivum* L., Giza 168) under open field conditions. Twenty grains were sown per pot (15.5 cm inter-diater and 21 cm height, containing 4 holes); each pot had 5 kg of the studied soil samples after preparation.

76 Phosphorus treatments were performed using calcium super phosphate (15% P₂O₅) supplied at a rate of 0, 3.5, 7.0, 10.0 and 13.0 mg P kg⁻¹ soil to represent 0, 25, 50, 75 and 100% of the recommended rate of P fertilization by the Egyptian Ministry of Agriculture for wheat cultivation. It was added before plant cultivation and mixed uniformly with the studied soil in pots. Silicon in the form of sodium metasilicate penta-hydrate (Na₂SiO₃.5H₂O) was added at a rate of 200, 300 and 400 mg Si kg⁻¹ soil, in addition to the control treatment (without any Si additions). Silicon concentrations were applied to the soil once, after sowing the grains of wheat plants. Also, the experiment included combinations between all these concentrations of Si and P. The experimental design was laid out in a split plot one with three replications. The main plots were assigned to the P additions and the subplots were occupied by the Si applications.

87 Nitrogen was added in the form of ammonium nitrate (33.5% N) at a rate of 100 kg N fed⁻¹ for wheat cultivation. It was added in three split of equal doses after 2, 4 and 7 weeks from sowing. Potassium was added in the form of potassium sulfate (48% K₂O) at a rate of 50 kg K₂O fed⁻¹ for wheat plants. It was added once after 4 weeks from cultivation. The moisture content in the studied soil was maintained at field capacity using tap water till the end of the experiment.

93 Plants were harvested at booting growth stage (60 days from sowing), divided into root and shoot, and weighted the fresh and dry weights. Plant samples representing each studied treatment were oven dried at 70°C for 24 h, then ground in a stainless mill to be finally digested using a mixture of sulfuric acid and hydrogen peroxide according to the method described by Black et al. (1965). Aliquots were finally taken out and analyzed for total Si and P using a Spectrophotometer (Page et al., 1982).

See comments on separate :[2G]Comment sheet

Why do you dry for 24 hours :[3G]Comment only?

99 After plant harvest, soil samples were subjected to analyses of the studied elements to determine their availability. The available Si was extracted by 0.5 M acetic acid as described by Kondorfer et al. (2001) and determined colorimetrically by yellow silico-molybdic acid according to Page et al. (1982). Available P was extracted by using sodium bicarbonate (NaHCO₃) 0.5 M at pH 8.5 according to Watanabe and Olsen (1965) and determined using Spectrophotometer by using ascorbic acid method.

105 The obtained data of soil and plant were statistically analyzed according to the method described by Gomez and Gomez (1984) using the LSD at 5% level of significance. Correlation coefficients were calculated between available Si and P in soil and their contents in plants using linear equation.

109

Table 1. Some physical and chemical characteristics of the studied soil.

Particle size distribution (%)		Soluble ions (meq L ⁻¹)	
C. Sand	6.22	Ca ²⁺	4.18
F. Sand	2.66	Mg ²⁺	2.98
Silt	26.1	Na ⁺	2.75
Clay	65.0	K ⁺	1.91
Textural class	Clay	HCO ₃ ⁻	5.61
FC (%)	42.8	Cl ⁻	2.29
WP (%)	4.12	SO ₄ ²⁻	3.92
AW (%)	38.7	Available elements (mg kg ⁻¹)	
OM (%)	0.96	N	55.2
CaCO ₃ (%)	3.33	P	28.5
pH (1: 2.5 soil: water suspension)	7.80	K	189
EC (dS m ⁻¹ , in paste extract)	1.12	Si	268

Carbonate ions were not detected. Note:

112

Results and Discussion

Availability of silicon and phosphorus in soil

114 Data presented in Table 2 shows the available concentrations of Si and P in the studied clay soil at booting growth stage of wheat plants as affected by their ground application. Results revealed that available Si increased with increase in applied dose in soil, with significant increases as compared to the control treatment. The highest increase (56%) occurred due to application of 400 mg Si kg⁻¹, as compared to control. This result agreed with

See comments on separate sheet :[4G]Comment

Please briefly define the abbreviations used in the table e.g. C. Sand, FC, etc. :[5G]Comment

Indicate the level of significance :[6G]Comment

those of Lee and Kim (2007) who found that available Si in soil significantly increased with increasing the rate of silicate application and consequently affected the absorbed Si by plants. Also, the available P increased with increase in applied P concentration in soil, with significant differences as compared to the control treatment. The application of 3.5 mg P kg⁻¹ soil gave the lowest increase (10%) in available P, while the highest increase (22.9%) was given by the treatment having 13 mg P kg⁻¹ soil as compared to control. These findings agreed with those obtained by Opala (2017) who reported that soil available P was positively and significantly affected by increasing P application in the soil.

Regarding the interaction between the applied Si and P concentrations in soil, data illustrated in Fig. 1 showed that Si application increased the Si availability in soil under conditions of various P concentrations (Fig. 1A), as compared to the control treatment. Phosphorus application increased the P availability in soil under various Si concentrations (Fig. 1B), as compared to control. Phosphorus availability increased with increasing the concentration of either Si or P (Fig. 1C), as compared to control. Silicon availability increased with increasing either P or Si (Fig. 1D), as compared to control. Ma and Takahashi (1990) reported that Si can improve soil P bioavailability through sorptive interaction between the two elements and enhancement of phosphate solubility.


136

Table 2. Available Si and P in the studied soil at booting growth stage of wheat plants as affected by their ground applications.

Treatment mg Si kg ⁻¹	Available Si mg Si kg ⁻¹	Treatment mg P kg ⁻¹	Available P mg P kg ⁻¹
0.00	300	0.00	23.0
200	438	3.50	30.8
300	450	7.00	31.9
400	468	10.0	33.6
		13.0	34.4
LSD_{0.05}	13.4	LSD_{0.05}	5.90

139

Indicate the significance level : [7G]Comment



140
 141 Fig.11. Available concentration (mg kg^{-1}) of Si and P as affected by their interactions in the
 142 studied soil at booting growth stage of wheat plants.
 143

144 ***Vegetative growth parameters of wheat plants as affected by Si and P additions and their interactions***

146 Data in Table 3 shows the effect of either Si or P on fresh and dry weights along with moisture content in shoots and roots of wheat plants at booting growth stage. Results showed that application of $400 \text{ mg Si kg}^{-1}$ soil gave the highest increase percent for fresh and dry weights of either shoots (38.1 and 48.4%, respectively) or roots of wheat plants (24.7 and 54.5% respectively), as compared to the control treatment. These findings agreed with those obtained by Zia et al. (2017) who reported that shoot and root dry weight of wheat plants increased with Si applied. In that sense, Soratto et al. (2012) stated that Si played a favorable role in plant growth, mineral nutrition, and mechanical strength. The increase in plant growth may be attributed to the changes in physiological and morphological properties which are facilitated by the presence of Si. On the contrary, the treatment of $400 \text{ mg Si kg}^{-1}$ soil gave the lowest value of moisture content in shoots and roots of wheat plants at booting growth stage (after 60 days from cultivation), and this confirmed increasing the dry matter content by applying this treatment. Regarding the effect of P, results in Table 3 showed that, the treatment of 13 mg P kg^{-1} soil gave the highest % increase in the fresh and dry weights of

shoots (31 and 67%, respectively) or roots of wheat plants (78 and 154%, respectively), as compared to control. These results agreed with those obtained by Zhang et al. (2016) who reported that dry weight of shoots and roots of wheat plants increased substantially as the P application rate increased. Increasing the P application rate to the soil greatly increased root dry weight, root length and density, and consequently root surface area of the cultivated plants. Additionally, Marschner (1986) stated that the increment in growth and yield due to P fertilization may be attributed to the activation of metabolic processes, where its role in building phospholipids and nucleic acid is known. Moreover, this treatment (13 mg P kg⁻¹ soil) gave the lowest value of moisture content in shoots and roots of wheat plants.

169

Table 6. Fresh and dry weights of shoots and roots of wheat plants at booting growth stage as affected by Si and P applications to the studied soil.

Treatment mg kg ⁻¹	Fresh weight g plant ⁻¹		Dry weight g plant ⁻¹		Moisture content %	
	Shoots	Roots	Shoots	Roots	Shoots	Roots
Control	6.99	0.85	1.26	0.22	82.0	74.0
Si Concentrations						
Si ₂₀₀	7.25	0.88	1.36	0.25	81.0	72.0
Si ₃₀₀	8.28	0.95	1.50	0.27	82.0	72.0
Si ₄₀₀	9.65	1.06	1.87	0.34	81.0	68.0
LSD_{0.05}	0.55	0.16	0.33	0.02	3.08	7.05
P Concentrations						
P _{3.5}	8.47	1.17	1.77	0.23	79.0	80.0
P _{7.0}	8.61	1.23	1.81	0.29	79.0	76.0
P _{10.0}	9.00	1.33	2.01	0.32	78.0	70.0
P _{13.0}	9.18	1.52	2.11	0.56	77.0	63.0
LSD_{0.05}	0.90	0.22	0.30	0.06	4.81	5.35

172

Data in Table 4 shows that the interaction between Si and P applications generally increased the growth parameters of wheat plants. This may be due to some sort of synergistic interaction between them. Also, the treatment of 400 mg Si kg⁻¹ in combination with 13 mg P kg⁻¹ soil gave the highest % increase in fresh and dry weights of shoots (50 and 62%, respectively) and roots of wheat plants (109 and 186%, respectively), as compared to the

See comments on separate sheet :[8G]Comment sheet

control treatment. Regarding the moisture content in shoots and roots of wheat plants at booting growth stage, there are no significant differences among the studied treatments as compared to the control treatment or the control treatment has the highest values. In this same light Yang et al. (2008) reported that Si promoted the growth of maize seedling, dry matter accumulation of different organs and root/shoot ratio under low P stress. Also, Owino-gerroin and Mascho (2005) found that application of silicate and phosphate increased the shoot and root dry weights of maize plants and the greatest increases were observed when silicate was applied together with phosphate.

186

Table 7. Effect of Si and P interactions on fresh and dry weights of shoots and roots of wheat plants at booting growth stage as affected by their applications to the studied soil.

Treatment mg kg ⁻¹	Fresh weight g plant ⁻¹		Dry weight g plant ⁻¹		Moisture content %		
	Shoots	Roots	Shoots	Roots	Shoots	Roots	
P₀×Si₀	6.99	0.85	1.26	0.22	82.0	74.0	
P _{3.5}	Si ₂₀₀	7.80	0.91	1.26	0.26	82.2	71.3
	Si ₃₀₀	7.77	1.07	1.48	0.31	80.8	71.4
	Si ₄₀₀	8.41	1.25	1.67	0.36	80.2	71.4
P _{7.0}	Si ₂₀₀	7.78	1.08	1.49	0.31	80.8	71.1
	Si ₃₀₀	8.12	1.24	1.55	0.34	80.9	72.9
	Si ₄₀₀	9.77	1.48	1.82	0.45	81.4	69.5
P _{10.0}	Si ₂₀₀	8.43	1.21	1.67	0.37	79.8	69.3
	Si ₃₀₀	8.83	1.38	1.76	0.39	80.0	71.6
	Si ₄₀₀	10.0	1.51	1.97	0.49	80.4	67.3
P _{13.0}	Si ₂₀₀	8.66	1.37	1.87	0.39	78.4	71.6
	Si ₃₀₀	9.71	1.47	1.94	0.41	80.0	72.3
	Si ₄₀₀	10.5	1.78	2.04	0.63	80.6	64.7
LSD_{0.05}	0.96	0.11	0.14	0.05	2.77	3.69	

189

Silicon and phosphorus concentrations in wheat plants at booting growth stage

Data in Table 5 shows that Si concentration and its total content in shoots and roots of wheat plants at booting growth stage were increased by increasing its applied concentrations in the studied soil. This response was clearer with roots than shoots. Also, data illustrated in

Indicate the confidence levels : [9G]Comment
(i.e. the t and p values)

Rewrite this statement. It is : [10G]Comment
not clear

Fig. 19A and B showed strong correlations between the available Si in soil and its concentration in both shoots and roots ($R^2 = 0.93$ and 0.99) of wheat plants, respectively. The results were in coincidence with those obtained by Mali and Aery (2008) who reported that there was a significant direct relationship between tissue (shoot and root) Si content and its applied rate in the soil. Sarto et al. (2014) and Zia et al. (2017) added that plant supplied with exogenous Si had high concentration of Si in shoot. Also, the accumulation of Si in shoots and roots of wheat plants increased with increasing the applied rate as compared to the control treatment. Regarding the P concentration and its total content in shoots and roots of wheat plants, data in Table 5 showed that, there were significant increases in plant organs with which related positively with its applied concentration in the studied soil. Data illustrated in Fig. 3A and B confirmed these results, which show the correlation between available P in soil and its concentration in shoots of grown wheat plants was equal 0.83 and in roots 0.85. These results were in concordance with those obtained by Zhang et al. (2016) who reported that the concentration of P in different plant organs increased with increasing its applied rate in the soil.

See comments on separate sheet :[11G]Comment

Indicate the p value :[12G]Comment


209

Table 5. Concentration and total content of Si and P in shoots and roots of wheat plants at booting growth stage as affected by their applications to the studied soil.

Treatment mg kg ⁻¹	Concentration mg g ⁻¹		Total content mg plant ⁻¹	
	Shoots	Roots	Shoots	Roots
Si Concentrations				
Si ₀	3.50	4.60	4.41	1.01
Si ₂₀₀	5.72	6.73	7.78	1.68
Si ₃₀₀	6.35	6.93	10.9	1.87
Si ₄₀₀	7.30	7.07	11.9	2.40
LSD_{0.05}	0.97	0.40	1.50	0.37
P Concentrations				
P ₀	4.05	3.90	5.10	0.86
P _{3.5}	5.80	4.40	10.3	1.01
P _{7.0}	6.00	4.70	10.9	1.36
P _{10.0}	6.80	4.90	13.7	1.57
P _{13.0}	7.50	5.30	15.8	2.97

LSD_{0.05}	0.66	0.60	2.30	0.27
---------------------------	-------------	-------------	-------------	-------------

212




213

214

Fig.215 The relationship between available Si concentration (mg kg^{-1}) in the studied soil and its concentration (mg g^{-1}) in shoots and roots of the growing wheat plants.

217



218

219

Fig.220 The relationship between available P concentration in the studied soil and its concentration in shoots and roots of the growing wheat plants.

222

223 Data in Table 6 shows the effect of the interaction between Si and P on concentration and content of Si in shoots and roots of wheat plants at booting growth stage. Results show that the treatment of $400 \text{ mg Si kg}^{-1}$ in combination with 13 mg P kg^{-1} soil gave the highest concentration and total content of Si in shoots and roots of wheat plants. Similar results were found with P concentration and total content per plant as affected by the same

interaction in the soil (Table 7). Data in Table 8 showed positive and, in general, strong correlation between available concentration of Si and P in the soil, and their concentrations in shoots and roots of the grown wheat plants. These data agreed with those obtained by Roy et al. (1971) who found that Si application tended to increase P content in the green tops. These data suggested that Si enhanced P mobilization from metabolically less active to metabolically more active tissue. In addition, Zia et al. (2017) found that Si application markedly increased the concentration of P in shoots. The increase in Si concentration in root medium may increase the uptake of P. It may be due to desorption of P from adsorption sites with the soil. Bai et al. (2013) reported that the critical levels of soil Olsen-P for optimal crop yield ranged from 10.9 to 21.4 mg kg⁻¹, and these levels were highly affected by crop species, soil type, pH and soil organic matter content. While the critical levels of soil available Si extracted by 0.5 M acetic acid for rice plants ranged from 44 to 116 ppm, according to Naranaswamy (2007). Considering these levels into account, the applied P in the studied soil till 13 mg kg⁻¹ considered adequate, while applying Si to the studied soil till 400 mg kg⁻¹ considered high but not within the toxic range according to the previous studies of Alovise et al. (2014).

244

Table 5. Concentration and total content of Si in shoots and roots of wheat plants at booting growth stage as affected by Si and P interactions in the studied soil.

Treatment mg kg ⁻¹	Concentration of Si mg g ⁻¹		Total content of Si mg plant ⁻¹		
	Shoots	Roots	Shoots	Roots	
P₀×Si₀	3.50	4.60	4.41	1.01	
P _{3.5}	Si ₂₀₀	3.64	6.07	4.59	1.58
	Si ₃₀₀	4.11	6.72	6.08	2.08
	Si ₄₀₀	5.67	7.32	9.47	2.64
P _{7.0}	Si ₂₀₀	3.53	6.43	5.26	1.99
	Si ₃₀₀	4.60	6.77	7.13	2.28
	Si ₄₀₀	5.89	7.63	10.7	3.43
P _{10.0}	Si ₂₀₀	3.60	6.90	6.01	2.55
	Si ₃₀₀	4.02	7.13	7.08	2.78
	Si ₄₀₀	6.04	7.69	11.9	3.77
P _{13.0}	Si ₂₀₀	6.02	6.98	11.3	2.72

Si ₃₀₀	6.07	7.31	11.8	3.00
Si ₄₀₀	6.33	8.64	12.9	5.44
LSD_{0.05}	1.23	0.73	1.93	0.52

247

Table 147. Concentration and total content of P in shoots and roots of wheat plants at booting growth stage as affected by Si and P interactions in the studied soil.

Treatment mg kg ⁻¹	Concentration of P mg g ⁻¹		Total content of P mg plant ⁻¹		
	Shoots	Roots	Shoots	Roots	
P₀ × Si₀	4.05	3.90	5.10	0.86	
P _{3.5}	Si ₂₀₀	4.15	4.33	5.23	1.13
	Si ₃₀₀	5.37	4.54	7.95	1.41
	Si ₄₀₀	5.88	4.51	9.82	1.62
P _{7.0}	Si ₂₀₀	5.10	4.77	7.60	1.48
	Si ₃₀₀	5.53	5.49	8.57	1.87
	Si ₄₀₀	6.12	5.22	11.1	2.35
P _{10.0}	Si ₂₀₀	5.66	4.80	9.45	1.78
	Si ₃₀₀	5.78	5.48	10.2	2.14
	Si ₄₀₀	6.01	5.72	11.8	2.80
P _{13.0}	Si ₂₀₀	5.82	5.18	10.9	2.02
	Si ₃₀₀	6.05	5.07	11.7	2.08
	Si ₄₀₀	6.41	5.08	13.1	3.20
LSD_{0.05}	0.62	0.94	1.29	0.40	

250

Table 148. Relationship between Si and P concentrations in shoots and roots of wheat plant, and their available concentrations in the studied soil.

Treatment	R ² (Si)		R ² (P)	
	Shoots	Roots	Shoots	Roots
Si + P _{3.5}	0.97	0.80	0.97	0.80
Si + P _{7.0}	0.97	0.99	0.97	0.50
Si + P _{10.0}	0.99	0.98	0.80	0.93
Si + P _{13.0}	0.98	0.96	0.99	0.93

P + Si ₂₀₀	0.50	0.90	0.80	0.63
P + Si ₃₀₀	0.20	0.97	0.96	0.20
P + Si ₄₀₀	0.90	0.62	0.50	0.30

253

254

Conclusion

255 It could be concluded that application of Si to the clay soil can release P into the soil
 256 solution and make it readily available to the plant absorption. So, it could be recommended
 257 that selecting good P fertilization design, including time and rate of addition, goes along with
 258 value of available Si in the studied soil. In addition, Si fertilization with suitable rate is very
 259 important to encourage wheat plant growth and productivity. The treatment of 400 mg Si kg⁻¹
 260 in combination with 13 mg P kg⁻¹ soil gave the highest growth parameters and total content of
 261 both Si and P in the different wheat plants organs.

262

263

264

References

- 265 Academy of Scientific Research and Technology (2017). The national campaign for upraising
 266 wheat production. [http://www.asrt.sci.eg/index.php/food-agriculture/132-](http://www.asrt.sci.eg/index.php/food-agriculture/132-the-national-campaign)
 267 [the-national-campaign](http://www.asrt.sci.eg/index.php/food-agriculture/132-the-national-campaign)
- 268 Alovero, A.M.T., Neto, A.E.F., Carneiro, L.F., Curi, N. and Alovero, A.A. (2014). Silicon-
 269 phosphorus interactions in soils cultivated with bean plants.
 270 *Acta Scientiarum Agron.* **36**, 79-86.
- 271 Bai, Li, H., Yang, X., Zhou, B., Shi, X., Wang, B., Li, D. et al. (2013). The critical soil P
 272 levels for crop yield, soil fertility and environmental safety in different soil types.
 273 *Plant Soil*, **372**, 27-37.
- 274 Black, C.A., Evans, D.D., White, J.L., Ensminger, L.E., Clark, F.E. and Dinauer, R.C. (1965).
 275 *Methods of Soil Analysis*. Am. Soc. Agron., Madison, Wisconsin, USA.
- 276 Carrijo, R., Furtini Neto, A.E., Santos, C.D., Fernandes, L.A., Curi, N. and Rodrigues, D.C.
 277 (2001). Interação silício-fósforo em solos cultivados com eucalipto em casa de
 278 vegetação. *Pesquisa Agropecuária Brasileira*, **36**, 557-565.
- 279 Datnoff, L.E., Rodrigues, F.A. and Seebold, K.W. (2009). Silicon and plant disease. In:
 280 *Mineral Nutrition and Plant Disease*. Datnoff, L.E., Elmer, W.H. and Huber, D.M.
 281 (Eds.). The American Phytopathological Society Press, St. Paul, MN, pp. 233-246.
- 282 Epstein, E. (1999). Silicon. *Annual Rev. Plant Physiol. Plant Mol. Biol.* **50**, 641-664.

- Gomez, K.A. and Gomez, A.A. (1984). *Statistical Procedures for Agricultural Research*.
284 John Wiley and Sons, Inc., New York, USA.
- Klute, A. (1986). *Methods of Soil Analysis*, part I, 2nd ed., Madison, Wisconsin, USA.
- Korfer, G.H., Snyder, M., Ulloa, G. and Datnoff, L.E. (2001). Calibration of soil and plant
286 silicon analysis for rice production. *J. Plant Nutr.* **24**, 1071-1084.
- Lee, B. and Kim, P.J. (2007). Reduction of phosphate adsorption by ion competition with
288 silicate in soil. *Korean J. Environ. Agric.* **26**, 286-293.
- Li, F., Ma, C.C. and Shang, Q.L. (2007). Effects of silicon on photosynthesis and
290 antioxidative enzymes of maize under drought stress. *Ying Yong Sheng Tai*
292 *Xue Bao.* **18**, 531-536.
- Ma, Takahashi, E. (1990). Effect of silicon on the growth and phosphorus uptake of rice.
293 *Plant Soil*, **126**, 115-119.
- Malhotra, M. and Aery, N.C. (2008). Influence of silicon on growth, relative water contents and
294 uptake of silicon, calcium and potassium in wheat grown in nutrient solution. *J.*
296 *Plant Nutr.* **31**, 1867-1876.
- Marschner, H. (1986). *Mineral Nutrition of Higher Plants*. Inst. Plant Nutr., Univ.
298 Hohenheim, FRG Academic Press, Harcourt Brace & Company, London, UK.
- Matvienkov, V.V. and Bocharnikova, E.A. (2001). The relationship between silicon and soil
300 physical and chemical properties. *Studies Plant Sci.* **8**, 209-219.
- Naraswamy, C. (2007). Calibration and categorization of plant available soil silicon in
302 different rice ecosystems for evaluating drought and disease resistance in rice
304 genotypes. Ph. D. Thesis, Department of Soil Science and Agricultural Chemistry,
305 University of Agricultural Sciences, Gkvk, Bangalore.
- Opabode, P.A. (2017). Influence of lime and phosphorus application rates on growth of maize in
306 an acid soil. *Adv. Agric.* **2017**, 5 pages. <https://doi.org/10.1155/2017/7083206>
- Owens, Gerroh, C. and Gascho, G.J. (2005). Effect of silicon on low pH soil phosphorus
307 uptake and on uptake and growth of maize. *Commun. Soil Sci. Plant Anal.* **35**, 2369-
309 **2378**.
- Pagano, A.L., Miller, R.H. and Keeney, D.R. (1982). *Methods of Soil Analysis*, part II, 2nd ed.,
310 Wisconsin, USA.
- Rizwan, M., Meunier, J., Miche, H. and Keller, C. (2012). Effect of silicon on reducing
312 cadmium toxicity in durum wheat (*Triticum turgidum* L. cv. Claudio W.) grown in a
314 soil with aged contamination. *J. Hazard. Mater.* **209-210**, 326-334.

- Rodrigues, F.A., Resende, R.S., Dallagnol, L.J. and Datnoff, L.E. (2015). Silicon potentiates host defense mechanisms against infection by plant pathogens. In: *Silicon and Plant Disease*. Rodrigues, F.A. and Datnoff, L.E. (Eds.). Springer International Publishing, Switzerland, p. 109–138.
- Roy, A.C., Ali, M.Y., Fox, R.L. and Silva, J.A. (1971). Influence of calcium silicate on phosphate solubility and availability in Hawaiian Latosols. In: *International Symposium on Soil Fertility Evaluation*, New Delhi, p. 757-768.
- Sanchez, D.R., Gatiboni, L.C. and Kaminski, J.K. (2008). Fatores que afetam a disponibilidade e o manejo da adubação fosfatada em solos sob sistema plantio direto. *Ciência Rural*, **38**, 576-586.
- Sartorius, M.V.M., Lana, M.C., Rampim, L., Rosset, J.S., Wobeto, J.R., Ecco, M., Bassegio, D. and da Costa, P.F. (2014). Effect of silicate on nutrition and yield of wheat. *African J. Agric. Res.* **9**, 956-962.
- Soriano, R.P., Crusciol, C.A.C., Castro, G.S.A., Costa, C.H.M. and Ferrari, J. (2012). Leaf application of silicic acid to white oat and wheat. *R. Bras. Ci. Solo*, **36**, 1538-1544.
- Tokimatsu, A.M., Furtini Neto, A.E., Carneiro, L.F., Curi, N., Santos, J.Z.L. and Alovisei, A.A. (2011). Dinâmica das formas de fósforo em solos de textura e mineralogia contrastantes cultivados com arroz. *Acta Scientiarum Agron.* **33**, 171-179.
- Tubana, B.S., Babu, T. and Datnoff, L.E. (2016). A review of silicon in soils and plants and its role in us agriculture: History and future perspectives. *Soil Sci.* **181**, 1-19.
- Wardrobe, F.C. and Olsen, S.R. (1965). Test of an ascorbic acid method for determining phosphorus in water and NaHCO₃ extracts from soils. *Soil Sci. Soc. Amer. Proc.* **29**, 677- 678.
- Yang, X., Roonasi, P. and Holmgren, A. (2008). A study of sodium silicate in aqueous solution and sorbed by synthetic magnetite using in situ ATR-FTIR spectroscopy. *J. Colloid Interface Sci.* **328**, 41-47.
- Zhang, W., Liu, D., Liu, Y., Cui, Z., Chen, Z., and Zou, C. (2016). Zinc uptake and accumulation in winter wheat relative to changes in root morphology and mycorrhizal colonization following varying phosphorus application on calcareous soil. *Field Crop Res.* **197**, 74-82.
- Zia, Bakhat, H.F., Saqib, Z.A., Mustafa, G., Muhammad, S. and Ashraf, R. (2017). Effect of water management and silicon on germination, growth, phosphorus and arsenic uptake in rice. *Ecotoxicol. Environ. Safety*, **144**, 11-18.